

Cosmological consequences of MSSM flat directions

KARI ENQVIST¹

Department of Physical Sciences, University of Helsinki,

and Helsinki Institute of Physics

P.O. Box 9, FIN-00014 University of Helsinki, Finland.

AND

ANUPAM MAZUMDAR^{2,3}

The Abdus Salam, International Centre for Theoretical Physics,

Strada Costiera-11, 34100, Trieste, Italy.

February 1, 2008

Abstract

We review the cosmological implications of the flat directions of the Minimally Supersymmetric Standard Model (MSSM). We describe how field condensates are created along the flat directions because of inflationary fluctuations. The post-inflationary dynamical evolution of the field condensate can charge up the condensate with B or L in a process known as Affleck-Dine baryogenesis. Condensate fluctuations can give rise to both adiabatic and isocurvature density perturbations and could be observable in future cosmic microwave experiments. In many cases the condensate is however not the state of lowest energy but fragments, with many interesting cosmological consequences. Fragmentation is triggered by inflation-induced perturbations and the condensate lumps will eventually form non-topological solitons, known as Q -balls. Their proper-

ties depend on how supersymmetry breaking is transmitted to the MSSM; if by gravity, then the Q -balls are semi-stable but long-lived and can be the source of all the baryons and LSP dark matter; if by gauge interactions, the Q -balls can be absolutely stable and form dark matter that can be searched for directly. We also discuss some cosmological applications of generic flat directions and Q -balls in the context of self-interacting dark matter, inflatonic solitons and extra dimensions.

¹ kari.enqvist@helsinki.fi

² a.mazumdar@ictp.trieste.it, anupamm@hep.physics.mcgill.ca

³ Physics Department, McGill University, Canada, from October 2002

Contents

1	Introduction	6
2	Baryogenesis	11
2.1	Baryon asymmetric Universe	11
2.2	Thermal history of the Universe	12
2.2.1	Expanding Universe	12
2.2.2	Entropy	14
2.2.3	Nucleosynthesis	16
2.3	Requirements for baryogenesis	17
2.3.1	Non-conservation of baryonic charge	17
2.3.2	C and CP violation	18
2.3.3	Departure from thermal equilibrium	18
2.3.4	Sphalerons	20
2.3.5	Washing out $B + L$	22
2.4	Alternatives for baryogenesis	23
2.4.1	GUT-baryogenesis	23
2.4.2	Electroweak baryogenesis	25
2.4.3	Electroweak baryogenesis in MSSM	27
2.4.4	Leptogenesis	27
2.4.5	Baryogenesis through field condensate decay	29
2.5	Old Affleck-Dine baryogenesis	30
2.5.1	Classical motion of the order parameter	30
2.5.2	Condensate decay	32
3	Field fluctuations during inflation	34
3.1	Fluctuation spectrum in de Sitter space	34
3.2	Slow roll inflation	36
3.3	Primordial density perturbations	37
3.3.1	Adiabatic perturbations and the Sachs-Wolfe effect	38

3.3.2	Spectrum of adiabatic perturbations	40
3.3.3	Gravitational waves	42
3.4	Multi-field perturbations	43
3.4.1	Adiabatic and isocurvature conditions	44
3.4.2	Adiabatic perturbations due to multi-field	44
3.4.3	Isocurvature perturbations and CMB	45
3.5	Inflation models	46
3.5.1	Non-supersymmetric inflation	46
3.5.2	F-term inflation	47
3.5.3	D-term inflation	50
3.5.4	Supergravity corrections	51
3.6	Reheating of the Universe	52
3.6.1	Perturbative inflaton decay	52
3.6.2	Non-perturbative inflaton decay	53
3.6.3	Gravitino and inflatino problems	54
4	Flat directions	57
4.1	Degenerate vacua	57
4.2	MSSM and its potential	58
4.2.1	F-and D-renormalizable flat directions of MSSM	60
4.2.2	An example of F-and D-flat direction	61
4.3	Lifting the flat direction	63
4.3.1	Lifting by non-renormalizable operators	63
4.3.2	Lifting by soft supersymmetry breaking	65
4.4	Supersymmetry breaking in the MSSM	65
4.4.1	Soft supersymmetry breaking Lagrangian	66
4.4.2	Gravity mediated supersymmetry breaking	67
4.4.3	Gauge mediated supersymmetry breaking	68
4.5	Supersymmetry breaking in the early Universe	70
4.5.1	Inflaton-induced terms	70
4.5.2	Supergravity corrections	71

4.6	The potential for flat direction	74
4.6.1	F-term inflation	74
4.6.2	D-term inflation	75
5	Dynamics of flat directions	76
5.1	Running of the couplings	76
5.1.1	Running of gravitational coupling	76
5.1.2	Renormalization group equations in the MSSM	78
5.2	Hubble induced radiative corrections	79
5.2.1	The case with $C_I \approx -1$	80
5.2.2	The case with $C_I \approx +1$	83
5.2.3	Running of the flat direction field in no-scale supergravity . . .	85
5.3	Post-inflationary running of the flat direction	85
5.3.1	Gravity mediated supersymmetry breaking	86
5.3.2	Gauge mediated supersymmetry breaking	88
5.4	Density perturbations from the flat direction condensate	89
5.4.1	Energetics of flat direction and the inflaton field	89
5.4.2	Adiabatic perturbations during D-term inflation	92
5.4.3	Adiabatic perturbations during F-term inflation	93
5.4.4	Isocurvature fluctuations in D-term inflation	94
5.4.5	Isocurvature fluctuations in F-term inflation	97
5.4.6	MSSM flat directions as a source for curvature perturbations . .	98
5.5	Baryon number asymmetry	99
5.6	Thermal effects	102
5.6.1	Thermal corrections to the flat direction potential	102
5.6.2	Thermal evaporation of the flat direction	104
5.7	Baryosynthesis and neutrino mass	104
5.8	Trajectory of a flat direction	106
5.9	Instability of the coherent condensate	107
5.9.1	Negative pressure	107
5.9.2	Growth of perturbations in the AD condensate	109

5.9.3	The true ground state	110
5.10	Numerical studies of fragmentation	112
5.10.1	Perturbation theory	112
5.10.2	Lattice simulations	115
5.11	Equilibrium ensembles	120
6	Q-balls	122
6.1	Q -ball as a non-topological soliton	122
6.1.1	Proofs of existence	122
6.1.2	Beyond thin wall solution	123
6.2	Varieties of Q -balls	124
6.2.1	Thin wall Q -balls	124
6.2.2	Thick wall Q -balls in the gauge mediated case	125
6.2.3	Thick wall Q -balls in the gravity mediated case	126
6.2.4	Hybrid case: gauge and gravity mediated Q -ball	128
6.2.5	Effect of gravity on Q -balls	129
6.2.6	Q -balls and local gauge invariance	129
6.3	Q -ball decay	131
6.3.1	Surface evaporation to fermions	131
6.3.2	The decay temperature	133
6.3.3	Q -ball decay into a pair of bosons	133
6.4	Cosmological formation of Q -balls	136
6.4.1	In gravity mediated case	136
6.4.2	In gauge mediated case	137
6.5	Q -ball collisions	138
6.6	Q -balls in a thermal bath	140
6.6.1	Dissociation	141
6.6.2	Diffusion	142
6.6.3	Evaporation at finite T	142

7	Cosmological consequences of Q-balls	145
7.1	L -ball cosmology	145
7.2	B -ball cosmology	146
7.3	B -balls in gravity mediated supersymmetry breaking	148
7.3.1	B -ball Baryogenesis	149
7.3.2	LSP dark matter from B -ball decay	150
7.3.3	The LSP abundance	152
7.3.4	Which direction?	154
7.3.5	Direct LSP searches and B -balls	155
7.4	Q -balls and gauge mediated supersymmetry breaking	157
7.4.1	Baryogenesis and gauge mediation	157
7.4.2	Generic gauge mediated models	159
7.4.3	Late formation of gauged Q -balls	160
7.5	Q -balls as self-interacting dark matter	161
7.6	Direct searches for gauge mediated Q -balls	162
8	Flat directions other than MSSM	166
8.1	Fragmentation of the inflaton condensate	166
8.1.1	Reheating as a surface effect	166
8.1.2	Q -balls from the inflaton condensate	168
8.2	Affleck-Dine baryogenesis without MSSM flat directions	170
8.2.1	Leptogenesis with sneutrino	170
8.2.2	AD baryogenesis in theories with low scale quantum gravity . .	171
8.3	Solitogenesis	174
8.3.1	Accretion of charge by Q -balls	174
8.3.2	Phase transition via solitogenesis	176

1 Introduction

The interplay between particle physics and cosmology plays an increasing role in understanding the physics beyond the Standard Model (SM) [1] and the early Universe before the era of Big Bang Nucleosynthesis (BBN) [2, 3]. On both fronts we currently lack hard data. Above the electroweak scale $E \sim \mathcal{O}(100)$ GeV, the particle content is largely unknown, while beyond the BBN scale $T \sim \mathcal{O}(1)$ MeV, there is no direct information about the thermal history of the Universe. However, there are some observational hints, as well as a number of theoretical considerations, which seem to be pointing towards a wealth of new physics both at small distances and in the very early Universe. Perhaps most importantly, new data is expected soon from accelerator experiments such as LHC and from cosmological measurements carried out by satellites such as MAP [4] and Planck [5].

In cosmology the recent observations of the cosmic microwave background (CMB) radiation, which has a temperature $\sim 2.728 \pm 0.004$ K [6], have given rise to an era of precision cosmology. The Cosmic Background Explorer (COBE) satellite [7] first detected in a full-sky map a temperature perturbation of one part in 10^5 at scales larger than 7 degrees [8]. The irregularities are present at a scale larger than the size of the horizon at the time when the microwave photons were generated and cannot be explained within the traditional hot Big Bang model [9]. The recent balloon experiments BOOMERANG [10] and MAXIMA [11], together with the ground-based DASI [12] experiment have established the existence of the first few acoustic peaks in the positions predicted by cosmic inflation [13, 14, 15, 9]. Inflation, a period of exponential expansion in the very early Universe, is a direct link to physics at energy scales that will not be accessible to Earth-bound experiments for any foreseeable future. Inflation could occur because a slowly rolling scalar field, the inflaton, dynamically gives rise to an epoch dominated by a false vacuum. During inflation quantum fluctuations are imprinted on space-time as energy perturbations which then are stretched outside the causal horizon. These primordial fluctuations eventually re-enter our horizon, whence their form can be extracted from the CMB (for a review, see [16, 9]).

Inflation can be considered as a model for the origin of matter since all matter arises from the vacuum energy stored in the inflaton field. However the present models do not give clear predictions as to what sort of matter there is to be found in the Universe. From observations we know that baryons constitute about 3% of the total mass [3], whereas relic diffuse cosmic ray background virtually excludes any domains of anti-baryons in the visible Universe [17]. Almost 30% of the total energy density is in non-luminous, non-baryonic dark matter [18]. Its origin and nature is unknown, although various simulations of large scale structure formation suggest that there must be at least some *cold dark matter* (CDM), comprising of particles with negligible velocity, although there may also be a component of *hot dark matter* (HDM), comprising of particles with relativistic velocities [19]. The rest of the energy density is in the form of dark energy [20, 21].

The striking asymmetry in the baryonic matter has existed at least since the time of BBN and plays an important role in providing the right abundances for the light elements. The present Helium (3He), Deuterium (D) and Lithium abundances suggest a baryon density and an asymmetry relative to photon density of order 10^{-10} [3]. Such an asymmetry is larger by a factor of 10^9 than what it should have been by merely assuming a initially baryon symmetric hot Big Bang [22]. Therefore baryon asymmetry must have been created dynamically in the early Universe.

The origin of baryon asymmetry and dark matter bring cosmology and particle physics together. Within SM all the three Sakharov conditions for baryogenesis [23] are in principle met; there is baryon number violation, C and CP violation, and an out-of-equilibrium environment during a first-order electroweak phase transition. However, it has turned out that within SM the electroweak phase transition is not strong enough [24, 25, 26, 27], and therefore the existence of baryons requires new physics. Regarding HDM, light neutrinos are a possible candidate [19, 28], but there is no candidate for CDM in the SM. HDM alone cannot lead a successful structure formation because of HDM free streaming length [29, 19]. Therefore one must resort to physics beyond the SM also to find a candidate for CDM [18].

The tangible evidence for small but non-vanishing neutrino masses as indicated by

the neutrino oscillations observed by the Super-Kamiokande [30] and SNO collaborations [31] is definitely another indication for new physics beyond the SM. The main sources of neutrino mass could be either Dirac or Majorana. A Dirac neutrino would require a large fine tuning in the Yukawa sector (one part in 10^{11}) while a Majorana mass would appear to require a scale much above the electroweak scale together with an extension of the SM gauge group $SU(3)_C \times SU(2)_L \times U(1)_Y$. In the Majorana case the lightness of the neutrino could be explained via the see-saw mechanism [32, 33].

A theoretical conundrum is that the mass scale of SM is $\sim \mathcal{O}(100)$ GeV, much lower than the scale of gravity $M_P = (8\pi G_N)^{-1/2} = 2.436 \times 10^{18}$ GeV, and not protected from quantum corrections. The most popular remedy is of course supersymmetry (for a review, see [34, 35, 37]), despite the fact that so far supersymmetry has evaded all observations [38]. The minimal supersymmetric extension of the SM is called the MSSM. Supersymmetry must be broken at a scale $\sim \mathcal{O}(1)$ TeV, presumably in some hidden sector from which breaking is transmitted to the MSSM, e.g., by gravitational [34, 35] or gauge interactions [39].

In the MSSM the number of degrees of freedom are increased by virtue of the supersymmetric counterparts of the SM bosons and fermions. One of them, known as the lightest supersymmetric particle (LSP), could be absolutely stable with a mass of the order of supersymmetry breaking scale. LSP would be a natural candidate for CDM (see e.g. [18]). In addition, because of the larger parameter space, electroweak baryogenesis in MSSM in principle has a much better chance to succeed. However, there are a number of important constraints, and lately Higgs searches at LEP have narrowed down the parameter space to the point where it has all but disappeared [40, 41, 42].

Electroweak baryogenesis within MSSM thus appears to be heading towards deep trouble. Moreover, although MSSM can provide CDM, there is no connection between dark matter and electroweak baryogenesis. On the other hand, by virtue of supersymmetry, MSSM has the intriguing feature that there are directions in the field space which have virtually no potential. They are usually known as *flat directions*, which are made up of squarks and sleptons and therefore carry baryon number and/or lepton

number. The MSSM flat directions have been all classified [43].

Because it does not cost anything in energy, during inflation squarks and sleptons are free to fluctuate along the flat directions and form scalar condensates. Because inflation smoothes out all gradients, only the homogeneous condensate mode survives. However, like any massless scalar field, the condensate is subject to inflaton-induced zero point fluctuations which impart a small, and in inflation models a calculable, spectrum of perturbations on the condensate. After inflation the dynamical evolution of the condensate can charge the condensate up with a large baryon or lepton number, which can then be released into the Universe when the condensate decays, as was first discussed by Affleck and Dine [44].

The potential along the MSSM flat direction is not completely flat because of supersymmetry breaking. In addition to the usual soft supersymmetry breaking, the non-zero energy density of the early Universe also breaks supersymmetry, in particular during inflation when the Hubble expansion dominates over any low energy supersymmetry breaking scale [45, 46]. Flatness can also be spoiled by higher-order non-renormalizable terms, and the details of the condensate dynamics depend on these.

In most cases, the MSSM condensate along a flat direction is however not the state of lowest energy. The condensate typically has a negative pressure, which causes the inflation-induced perturbations to grow. Because of this the condensate fragments, usually when the Hubble scale equals the supersymmetry breaking scale, into lumps of condensate matter which eventually settle down to non-topological solitons dubbed as Q -balls by Coleman [47]. Q -balls carry a global charge, which in the case of MSSM is either B or L .

The properties of Q -balls depend on supersymmetry breaking. If transmitted to MSSM by gravity, the Q -balls turn out to be only semistable but nevertheless long-lived compared with the time scales of the very early Universe [48]. When they decay, they may provide not only the baryonic matter but also dark matter LSPs [49]. If supersymmetry breaking is transmitted from the hidden sector to MSSM by gauge interactions, the resulting Q -balls would be stable and could exist at present as a form of dark matter [50]. In this case one can make direct searches for their existence

[51]. In both cases there is a prediction for the relation between the baryon and dark matter densities. Moreover, the condensate perturbations are inherited by the Q -balls, and can thus be a source of both isocurvature and adiabatic density perturbations [52, 53, 54].

This review is organized as follows. In Section 2, we recapitulate some basic cosmology, and in particular baryogenesis. We briefly discuss various popular schemes of baryogenesis and describe the original Affleck-Dine baryogenesis. In Section 3, we present some background material for inflation, mainly concentrating on supersymmetric models. Quantum fluctuations and reheating are also discussed. In Section 4, we present the MSSM flat directions and discuss their properties. Various contributions to the flat direction potential in the early Universe are listed. Low energy supersymmetry breaking schemes, such as gravity and gauge mediation, are also discussed. In Section 5, we discuss the dynamical properties of flat directions and the running of the flat direction potential due to gauge and Yukawa interactions. Leptogenesis along LH_u flat direction, and the condensate evaporation in a thermal bath, is also described. We discuss fragmentation of the condensates for both gravity and gauge mediated supersymmetry breaking and present the relevant numerical studies. In Section 6, Q -ball properties are presented in detail. We describe various types of Q -balls, their interactions and their behavior at finite temperature. We discuss surface evaporation, diffusion, and dissociation of charge from Q -balls in a thermal bath. In Section 7, we focus on the cosmological consequences of Q -balls. We consider Q -ball baryogenesis and non-thermal dark matter generation through charge evaporation for different types of Q -balls. We discuss Q -balls as self-interacting dark matter and present experimental and astrophysical constraints on stable Q -balls. In Section 8, we briefly survey beyond-the-MSSM-condensates by considering inflatonic Q -balls and Affleck-Dine mechanism without MSSM flat directions. We also describe solitosynthesis, a process of accumulating large Q -balls in a charge asymmetric Universe.

2 Baryogenesis

2.1 Baryon asymmetric Universe

There are only insignificant amounts of anti-particles within the solar system. Cosmic ray showers contain $\sim 10^{-4}$ anti-protons for each proton [55], but the anti-protons are by-products of the interaction of the primary beam with the interstellar dust medium. This strongly suggest that galaxies and intergalactic medium is made up of matter rather than anti-matter, and if there were any anti-matter, the abundance has to be smaller than one part in 10^4 . The absence of annihilation radiation from the Virgo cluster shows that little anti-matter is to be found within a 20 Mpc sphere, and the relic diffuse cosmic ray background virtually excludes domains of anti-matter in the visible Universe [17].

The best present estimation for the baryon density comes from BBN [56] combined with the CMB experiments and it is given by [57]

$$0.010 \leq \Omega_b h^2 \leq 0.022, \quad (1)$$

where $\Omega_b \equiv \rho_b/\rho_c$ defines the fractional baryon density ρ_b with respect to the critical energy density of the Universe: $\rho_c = 1.88 h^2 \times 10^{-29} \text{ g cm}^{-3}$. The observational uncertainties in the present value of the Hubble constant; $H_0 = 100 h \text{ km} \cdot \text{s}^{-1} \cdot \text{Mpc}^{-1} \approx (h/3000) \text{Mpc}^{-1}$ are encoded in h . Various considerations such as Hubble Space Telescope observations and type Ia supernova data suggest that $h = 0.70$ [58]. However, from the age of the globular cluster which comes out to be 11 Gyr, h seems to take lower value of about 0.5 [59]. The present convention is to take $0.5 \leq h \leq 0.8$. In terms of the baryon and photon number densities we may write

$$\eta \equiv \frac{n_b - n_{\bar{b}}}{n_\gamma} = 2.68 \times 10^{-8} \Omega_b h^2, \quad (2)$$

where n_b is the baryon number density and $n_{\bar{b}}$ is for anti-baryons. The photon number density is given by $n_\gamma \equiv (2\zeta(3)/\pi^2)T^3$. Observations of the deuterium abundance in quasar absorption lines suggest [60]

$$4(3) \times 10^{-10} \leq \eta \leq 7(10) \times 10^{-10}. \quad (3)$$

The conservative bounds are in parenthesis.

Often in the literature the baryon asymmetry is given in relation to the entropy density $s = 1.8g_*n_\gamma$, where g_* measures the effective number of relativistic species which itself a function of temperature. At the present time $g_* \approx 3.36$, while during BBN $g_* \approx 10.11$, rising up to 106.75 at $T \gg 100$ GeV. In the presence of supersymmetry at $T \gg 100$ GeV, the number of effective relativistic species are doubled to 213.30.

The baryon asymmetry B , defined as the difference of baryon and anti-baryon number densities relative to the entropy density, is bounded by

$$5.7(4.3) \times 10^{-11} \leq B \equiv \frac{n_b - n_{\bar{b}}}{s} \leq 9.9(14) \times 10^{-11}, \quad (4)$$

where the numbers in parenthesis are conservative bounds [60]. If at the beginning $\eta = 0$, then the origin of this small number can not be understood in a CPT invariant Universe by a mere thermal decoupling of nucleons and anti-nucleons at $T \sim 20$ MeV. The resulting asymmetry would be too small by at least nine orders of magnitude, see [22].

2.2 Thermal history of the Universe

2.2.1 Expanding Universe

The hot Big Bang cosmology assumes that the Universe is spatially homogeneous and isotropic and can be described by the Friedmann-Robertson-Walker (FRW) metric

$$ds^2 = g_{\mu\nu}dx^\mu dx^\nu = dt^2 - a^2(t) \left[\frac{dr^2}{1 - Kr^2} + r^2(d\theta^2 + \sin^2\theta d\phi^2) \right], \quad (5)$$

where $a(t)$ is the scale factor that determines the expansion or the contraction of the Universe; the constant K defines the spatial geometry. If $K = 0$, the Universe is flat and has Euclidean geometry, otherwise there is a spatial curvature corresponding either to a closed elliptic ($K = +1$) or an open hyperbolic ($K = -1$) geometries. The value of K cannot however fix the global topology; for instance, an Euclidean topology can be flat and infinite $\mathbb{R}^3$, or a surface of a 3-torus $\mathbb{T}^3$. However, topology has other observable consequences, e.g., for the pattern of CMB temperature fluctuations [61].

There are two characteristic scales corresponding to the homogeneous and isotropic Universe: the curvature scale $r_{\text{curv}} = a(t)|K|^{-1/2}$, and the Hubble scale

$$H^{-1} = \left[\frac{\dot{a}(t)}{a(t)} \right]^{-1}, \quad (6)$$

where dot denotes derivative w.r.t. t . The Hubble time is denoted by

$$t_{\text{Hub}} = \int_i^f \frac{dt}{H^{-1}} = \ln \left(\frac{a_f}{a_i} \right). \quad (7)$$

The behavior of the scale factor depends on the energy momentum tensor of the Universe. For a perfect fluid

$$T_{\mu\nu} = -pg_{\mu\nu} + (\rho + p)u_\mu u_\nu, \quad (8)$$

where ρ is the energy density and p is the pressure of a fluid and the four velocity $u_\mu \equiv dx_\mu/ds$. For the FRW metric and for the perfect fluid the equations of motion gives the Friedmann-Lemaitre equation

$$H^2 = \frac{\rho}{3M_{\text{P}}^2} - \frac{K}{a(t)^2}, \quad (9)$$

also known as the Hubble equation. The acceleration equation is given by

$$\frac{\ddot{a}(t)}{a(t)} = -\frac{1}{6M_{\text{P}}^2}(\rho + 3p), \quad (10)$$

and the conservation of the energy momentum tensor $T_{;\nu}^{\mu\nu} = 0$ gives

$$\frac{d(\rho a^3)}{da} = -3pa^2. \quad (11)$$

Note that ρa^3 is constantly decreasing in an expanding Universe for a positive pressure.

The early Universe is believed to have been radiation dominated with $p = \rho/3$ and $a(t) \propto t^{1/2}$, followed by a matter dominated era with $p = 0$ and $a(t) \propto t^{2/3}$. The early Universe might also have had an era of acceleration, known as the inflationary phase, which could have happened only if

$$\ddot{a} > 0 \Leftrightarrow \rho + 3p < 0. \quad (12)$$

A geometric way of defining inflation is [9]

$$\frac{d(H^{-1}/a(t))}{dt} < 0, \quad (13)$$

which states that the Hubble length as measured in comoving coordinates decreases during inflation. We will use this particular definition of inflation very often while discussing the number of e-foldings and density perturbations.

The Hubble expansion rate is related to the temperature by

$$H = \sqrt{\frac{\rho}{3M_{\text{P}}^2}} = 1.66 \times g_*^{1/2} \frac{T^2}{M_{\text{P}}}, \quad (14)$$

where g_* is the total number of relativistic degrees of freedom and it is given by

$$g_*(T) = \sum_{i=b} g_i \left(\frac{T_i}{T} \right)^4 + \frac{7}{8} \sum_{i=f} g_i \left(\frac{T_i}{T} \right)^4. \quad (15)$$

Here T_i denotes the effective temperature of species i , which decouples at a temperature $T = T_{\text{D}}$.

During the radiation era when $H = (1/2t)$, one finds

$$\frac{t}{1 \text{ s}} \approx 2.42 g_*^{-1/2} \left(\frac{1 \text{ MeV}}{T} \right)^2. \quad (16)$$

2.2.2 Entropy

An ideal gas of particles respects the Fermi-Dirac or Bose-Einstein distributions

$$f_i(\mathbf{p}, \mu, T) = [\exp((E_i - \mu_i)/T) \mp 1]^{-1}, \quad (17)$$

where $E_i^2 = |\mathbf{p}|^2 + m^2$, μ_i represents the chemical potential of the species i , $-/+$ corresponds to Bose/Fermi statistics. The value of μ is equal and opposite for particles and anti-particles. Therefore in the early Universe a finite net chemical potential is proportional to the particle anti-particle asymmetry. The bound on charge asymmetry relative to the photon number density is severe, less than one part in 10^{43} at temperatures close to BBN [62], while baryon asymmetry is comparatively larger, but still small enough for $\mu_e, \mu_b \approx 0$ to be an excellent approximation. Neutrinos may however carry a net $B - L$ charge which need not be vanishingly small at early times, although a large enough neutrino chemical potential can affect nucleosynthesis, for example, see [28].

The number density n , energy density ρ , and pressure p can be expressed in terms of temperature, and g_i is the number of internal degrees of freedom [63]

$$n_i(T) = \frac{g_i}{(2\pi)^3} \int f_i(\mathbf{p}, \mu, T) d^3\mathbf{p} = \frac{g_i}{2\pi^2} T^3 I_i^{11}(\mp),$$

$$\begin{aligned}\rho_i(T) &= \frac{g_i}{(2\pi)^3} \int E_i(\mathbf{p}) f_i(\mathbf{p}, \mu, T) d^3\mathbf{p} = \frac{g_i}{2\pi^2} T^4 I_i^{21}(\mp), \\ p_i(T) &= \frac{g_i}{(2\pi)^3} \int \frac{|\mathbf{p}|^2}{3E_i(\mathbf{p})} f_i(\mathbf{p}, \mu, T) d^3\mathbf{p} = \frac{g_i}{6\pi^2} T^4 I_i^{03}(\mp),\end{aligned}\quad (18)$$

where

$$I_i^{ab}(\mp) \equiv \int_{x_i}^{\infty} \frac{y^a (y^2 - x_i^2)^{b/2}}{(e^y \mp 1)} dy, \quad x_i \equiv \frac{m_i}{T}. \quad (19)$$

For a relativistic case with $x_i \ll 1$,

$$\begin{aligned}I_r^{11}(-) &= 2\zeta(3), & I_r^{21}(-) = I_r^{03}(-) &= \frac{\pi^4}{15}, & \text{for bosons,} \\ I_r^{11}(+) &= \frac{3\zeta(3)}{2}, & I_r^{21}(+) = I_r^{03}(+) &= \frac{7\pi^4}{120}, & \text{for fermions,}\end{aligned}\quad (20)$$

where ζ denotes the Riemann Zeta function and $\zeta(3) = 1.202$. Thus the energy density of radiation reads

$$\rho_r = \frac{\pi^2}{30} g_* T^4. \quad (21)$$

For non-relativistic particles with $x_i \gg 1$, one obtains for both bosons and fermions

$$n_{\text{nr}}(T) = \frac{\rho_{\text{nr}}}{m} = g_i \left(\frac{m_i T}{2\pi} \right)^{3/2} e^{-m_i/T}, \quad p_{\text{nr}} = 0. \quad (22)$$

If the chemical potential is non-zero, the exponential Eq. (22) also includes a factor $e^{+\mu_i/T}$.

The entropy density is defined as

$$s \equiv \frac{S}{T} = \frac{\rho_i + p_i}{T}, \quad (23)$$

where $d(sa^3) = 0$ is a thermodynamically conserved quantity. The decoupling temperature can be expressed as [64]

$$\frac{T_D}{T} = \left(\frac{g_{*S_A}(T_D)}{g_{*S_A}(T)} \frac{g_{*S-S_A}(T)}{g_{*S-S_A}(T_D)} \right)^{1/3}, \quad (24)$$

where S is the total entropy and S_A the entropy in the degrees of freedom that have decoupled at T_D .

2.2.3 Nucleosynthesis

According to BBN (for reviews see [3, 57]) the light elements 2H , 3He , 4He , and 7Li have been synthesized during the first few hundred seconds. The abundances depend on the baryon-to-photon ratio

$$\eta \equiv \frac{n_B}{n_\gamma}. \quad (25)$$

All the relevant physical processes take place essentially in the range from a few MeV ~ 0.1 sec down to $60 - 70$ KeV $\sim 10^3$ sec. During this period only photons, $e^\pm$ pairs, and the three neutrino flavors contribute significantly to the energy density. Any additional energy density may be parameterized in terms of the effective number of light neutrino species N_ν , so that

$$g_* = 10.75 + \frac{7}{4}(N_\nu - 3). \quad (26)$$

Nucleosynthesis starts off with a freezing out of the weak interaction between neutron and proton at $T_D \approx 0.8$ MeV. Free neutrons keep decaying until deuterium begins to form through $n + p \rightarrow d + \gamma$. Deuterium synthesis is over by $T_D \approx 0.086$ MeV (assuming $\eta = 5 \times 10^{-10}$). At T_D , neutron abundance has been depleted to $X_n(t_D) \equiv n/(n+p) \approx 0.122$. All the surviving neutrons are now captured through $n + D \rightarrow (^3H, ^3He)$, and subsequently by virtue of the process $(^3H, ^3He) + n \rightarrow ^4He$, which has a binding energy of 28.3 MeV. The total mass fraction of primordial helium, which is denoted by $Y_P(^4He)$, is given by

$$Y_P(^4He) \approx 2X_n(T_D) = 0.245. \quad (27)$$

Adopting the experimentally allowed range of $0.22 < Y_P < 0.26$, one can constraint that number of light neutrino species by [56]

$$N_\nu \leq 4. \quad (28)$$

The four LEP experiments combined give the best fit as [65]

$$N_\nu = 2.994 \pm 0.12. \quad (29)$$

Nucleosynthesis also constrains many non-conventional ideas, for instance alternative theories of gravity such as scalar-tensor theories [66].

Besides ${}^4\text{He}$, D and ${}^3\text{He}$ are produced at the level of 10^{-5} , and ${}^7\text{Li}$ at the level of 10^{-10} . The theoretical prediction has some slight problems in fitting the observed ${}^4\text{He}$ and ${}^7\text{Li}$ abundances. Both seem to indicate $1.7 \times 10^{-10} < \eta < 4.7 \times 10^{-10}$, corresponding to $0.006 < \Omega_b h^2 < 0.017$ with a central value $\Omega_b h^2 = 0.009$ [57]. The abundance ratio D/H is comparable with ${}^4\text{He}$ and ${}^7\text{Li}$ abundances at the 2σ level in the range $4.7 \times 10^{-10} < \eta < 6.2 \times 10^{-10}$, which corresponds to $0.017 < \Omega_b h^2 < 0.023$. The likelihood analysis which includes all the three elements (D, ${}^4\text{He}$, and ${}^7\text{Li}$) yields [57]

$$4.7 \times 10^{-10} < \eta < 6.2 \times 10^{-10}, \quad 0.017 < \Omega_b h^2 < 0.023. \quad (30)$$

Despite the uncertainties there appears to be a general concordance between theoretical BBN predictions and observations, which is now being bolstered by the CMB data from several different experiments. The results from the ground based DASI experiment indicates $\Omega_b h^2 = 0.022^{+0.004}_{-0.003}$ [12], while the results from the BOOMERANG balloon-borne experiment imply $\Omega_b h^2 = 0.021^{+0.004}_{-0.003}$ [10]. MAXIMA, another balloon experiment, quotes a somewhat larger value $\Omega_b h^2 = 0.0325 \pm 0.006$ [67].

2.3 Requirements for baryogenesis

As pointed out by Sakharov [23], baryogenesis requires three ingredients: (1) baryon number non-conservation, (2) C and CP violation, and (3) out-of-equilibrium condition. All these three conditions are believed to be met in the very early Universe¹.

2.3.1 Non-conservation of baryonic charge

In the SM, baryon number B is violated by non-perturbative instanton processes [70, 71]. At the quantum level both baryon number current J_B^μ and the lepton number current J_L^μ are not conserved because of chiral anomalies [72]. However the anoma-

¹There have been attempts [68, 69] for baryogenesis via a repulsive interaction between baryons and anti-baryons which would lead to their spatial separation before thermal decoupling of nucleons and anti-nucleons. However at such early times the causal horizon contained only a very small fraction of the solar mass so that the asymmetry could not be smooth at distances greater than the galactic size.

lous divergences of J_B^μ and J_L^μ come with an equal amplitude and an opposite sign. Therefore $B - L$ remains conserved, while $B + L$ may change via processes which interpolate between the multiple non-Abelian vacua of $SU(2)$. The probability for the $B + L$ violating transition is however exponentially suppressed [70, 71]. As was first pointed out by Manton [73], at high temperatures the situation is different, so that when $T \gg M_W$, baryon violating transitions are in fact copious (see Sect. 2.4.2).

In addition to baryogenesis, B violation also leads to proton decay in GUTs. For instance, the dimension 6 operator $(QQQL)/\Lambda$ generates observable proton decay unless $\Lambda \geq 10^{15}$ GeV. In the MSSM the bound is $\Lambda \geq 10^{26}$ GeV because the decay can take place via a dimension 5 operator. In the MSSM superpotential there are also terms which can lead to $\Delta L = 1$ and $\Delta B = 1$. Similarly there are other processes such as neutron-anti-neutron oscillations in SM and in supersymmetric theories which lead to $\Delta B = 2$ and $\Delta B = 1$ transitions [74]. These operators are constrained by the measurements of the proton lifetime, which yield the bound $\tau_p \geq 10^{33}$ years [65].

2.3.2 C and CP violation

Weak interactions ensures maximum C violation while neutral Kaon is an example of CP violation in the quark sector which has a relative strength $\sim 10^{-3}$ [65]. CP violation could also expected to be found in the neutrino sector. Beyond the SM there are many sources for CP violation. An example is the axion proposed for solving the strong CP problem in QCD [75]. Quantum fluctuations of light scalars in the early Universe, in particular during inflation, can create different domains of various C and CP phases. C and CP can also be spontaneously broken during a phase transition, so that domains of broken phases form with different CP -charges [76].

2.3.3 Departure from thermal equilibrium

Departure from a thermal equilibrium cannot be achieved by mere particle physics considerations but is coupled to the dynamical evolution of the Universe. If B -violation processes are in thermal equilibrium, the inverse processes will wash out the pre-existing asymmetry $(\Delta n_b)_0$ [77]. This is a consequence of S -matrix unitarity and

CPT -theorem [78]. However there are several ways of obtaining an out-of-equilibrium process in the early Universe.

- *Out-of-equilibrium decay or scattering:*

The Universe in a thermal equilibrium can not produce any asymmetry, rather it tries to equilibrate any pre-existing asymmetry. If the scattering rate $\Gamma < H$, the process can take place out-of-equilibrium. Such a situation is appropriate for e.g. GUT baryogenesis [78, 79].

- *Phase transitions:*

Phase transitions are ubiquitous in the early Universe. The transition could be of *first*, or of *second* (or of still higher) order. First order transitions proceed by barrier penetration and subsequent bubble nucleation resulting in a temporary departure from equilibrium. Second order phase transitions have no barrier between the symmetric and the broken phase. They are continuous and equilibrium is maintained throughout the transition.

Prime examples of first order phase transitions in the early Universe are the QCD and electroweak phase transitions. The nature and details of QCD phase transition is still very much an open debate [80, 25, 81], and although a mechanism for baryogenesis during QCD phase transition has been proposed [82], much more effort has been devoted to the electroweak phase transition [24, 26] (see Sect. 2.4.2).

- *Non-adiabatic motion of a scalar field:*

Any complex scalar field carries C and CP , but the symmetries can be broken by terms in the scalar potential. This can lead to a non-trivial trajectory of a complex scalar field in the phase space. If a coherent scalar field is trapped in a local minimum of the potential and if the shape of the potential changes to become a maximum, then the field may not have enough time to readjust with the potential and may experience completely non-adiabatic motion. This is similar to a second order phase transition but it is the non-adiabatic classical motion which prevails over the quantum fluctuations, and therefore, departure

from equilibrium can be achieved. If the field condensate carries a global charge such as the baryon number, the motion can charge up the condensate. This is the basis for the Affleck-Dine baryogenesis [44] (see Sect. 2.5).

2.3.4 Sphalerons

In the SM $B + L$ is very weakly violated in the vacuum [70]. At finite temperatures violation is large [73, 83, 84, 85, 86] by virtue of the sphaleron configurations, which mediate transitions between degenerate gauge vacua with different Chern-Simons numbers related to the net change of $B + L$. Thermal scattering produces sphalerons which in effect decay in $B + L$ non-conserving ways below 10^{12} GeV [87], and thus can exponentially wash away $B + L$ asymmetry. Sphalerons and associated electroweak baryogenesis has been reviewed in [88, 24, 26, 89, 90, 91]. Let us here just give a brief summary of the main ingredients.

- *Chiral anomalies*

An anomaly means that a classical current conservation no longer holds at the quantum level; an example is the chiral anomaly [72]. In the SM there is classical conservation of the baryon and lepton number currents J_B^μ and J_L^μ , but because of chiral anomaly the currents are not conserved. Instead [70],

$$\begin{aligned}\partial_\mu J_B^\mu &= -\frac{\alpha_2}{8\pi} N_g W_i^{\mu\nu} \tilde{W}_{i\mu\nu} + \frac{\alpha_1}{8\pi} N_g \left(\frac{4}{9} + \frac{1}{9} - \frac{2}{36} \right) F^{\alpha\beta} \tilde{F}_{\alpha\beta}, \\ \partial_\mu J_L^\mu &= -\frac{\alpha_2}{8\pi} N_g W_i^{\mu\nu} \tilde{W}_{i\mu\nu} + \frac{\alpha_1}{8\pi} N_g \left(1 - \frac{1}{2} \right) F^{\alpha\beta} \tilde{F}_{\alpha\beta},\end{aligned}\tag{31}$$

where N_g is the number of generations, α_2 and α_1 ($W_{i\mu\nu}$ and $F_{\mu\nu}$) are respectively the $SU(2)$ and $U(1)$ gauge couplings (field strengths), and the various numbers inside the brackets correspond to the squares of the hypercharges multiplied by the number of states. Note that while at the quantum level $B + L$ is violated, $B - L$ is still conserved.

- *Gauge theory vacua*

The vacuum structure of the gauge theories is very rich [71, 92]. In case of $SU(2)$, the vacua are classified by their homotopy class $\{\Omega_n(\mathbf{r})\}$, characterized by the

winding number n which labels the so called θ -vacua [71, 92]. A gauge invariant quantity is the difference in the winding number (Chern-Simons number)

$$N_{CS} \equiv n_+ - n_- = \frac{\alpha_2}{8\pi} \int d^4x W_a^{\mu\nu} \tilde{W}_{a\mu\nu} . \quad (32)$$

In the electroweak sector the field density $W\tilde{W}$ is related to the divergence of $B + L$ current. Therefore a change in $B + L$ reflects a change in the vacuum configuration and is determined by the difference in the winding number

$$\Delta(B + L) = \int d^4x \partial_\mu J_{B+L}^\mu = -\frac{\alpha_2}{4\pi} N_g \int d^4x W_a^{\mu\nu} \tilde{W}_{a\mu\nu} = -2N_g N_{CS} . \quad (33)$$

For three generations of SM leptons and quarks the minimal violation is $\Delta(B + L) = 6$. Note that the proton decay $p \rightarrow e^+ \pi^0$ requires $\Delta(B + L) = 2$, so that despite B -violation, proton decay is completely forbidden in the SM.

The probability amplitude for tunneling from an n vacuum at $t \rightarrow -\infty$ to an $n + N_{CS}$ vacuum at $t \rightarrow +\infty$ can be estimated by the WKB method [70]

$$P(N_{CS})_{B+L} \sim \exp\left(\frac{-4\pi N_{CS}}{\alpha_2(M_Z)}\right) \sim 10^{-162N_{CS}} . \quad (34)$$

Therefore, as advertised, the baryon number violation rate is totally negligible in the SM at zero temperature, but as argued in a seminal paper by Kuzmin, Rubakov and Shaposhnikov [93], at finite temperatures the situation is completely different.

- *Thermal tunneling*

The sphaleron is a field configuration sitting at the top of the potential barrier between two vacua with different Chern-Simons numbers and can be reached simply because of thermal fluctuations [93]. Neglecting $U(1)_Y$, the zero temperature sphaleron solution was first found by Manton and Klinkhamer [73, 83]. At finite temperature the energy obeys an approximate scaling law [85, 86] $E_{sph}(T) = E_{sph}(0) \langle \Phi(T) \rangle / \langle \Phi(0) \rangle$:

$$E_{sph}(T) = \frac{2m_W(T)}{\alpha_2} B(\lambda/g_2) , \quad (35)$$

where $m_W(T) = (1/2)g_2 \langle \Phi(T) \rangle$ is the mass of the W-boson and the function B has a weak dependence on λ/g_2 , where λ is the quartic self coupling of the

Higgs. Below the critical temperature of the electroweak phase transition, the sphaleron rate is exponentially suppressed [94]:

$$\Gamma \sim 2.8 \times 10^5 \kappa T^4 \left(\frac{\alpha_2}{4\pi} \right)^4 \left(\frac{E_{sph}(T)}{B(\lambda^2/g)} \right)^7 e^{-E_{sph}/T}. \quad (36)$$

where κ is the functional determinant which can take the values $10^{-4} \leq \kappa \leq 10^{-1}$ [95]. Above the critical temperature the rate is however unsuppressed. Requiring that the Chern-Simons number changes at most by $\Delta N_{CS} \sim 1$, one can estimate from Eq. (32) that $\Delta N_{CS} \sim g_2^2 l_{sph}^2 W_i^2 \sim 1 \rightarrow W_i \sim \frac{1}{g_2 l_{sph}}$. Therefore a typical energy of the sphaleron configuration is given by

$$E_{sph} \sim l_{sph}^3 (\partial W_i)^2 \sim \frac{1}{g_2^2 l_{sph}}. \quad (37)$$

At temperatures greater than the critical temperature there is no Boltzmann suppression, so that the thermal energy $\propto T \geq E_{sph}$. This determines the size of the sphaleron as

$$l_{sph} \geq \frac{1}{g_2^2 T}. \quad (38)$$

This is exactly the infrared cut-off generated by the magnetic mass of order $\sim g_2^2 T$. Therefore, based on this coherence length scale one can estimate the baryon number violation per volume $\sim l_{sph}^3$, and per unit time $\sim l_{sph}$. On dimensional grounds the transition probability would then be given by

$$\Gamma_{sph} \sim \frac{1}{l_{sph}^3 t} \sim \kappa (\alpha_2 T)^4. \quad (39)$$

where κ is a constant which incorporates various uncertainties. However, the process is inherently non-perturbative, and it has been argued that damping of the magnetic field in a plasma suppresses the sphaleron rate by an extra power of α_2 [96], with the consequence that $\Gamma_{sph} \sim \alpha_2^5 T^4$. Lattice simulations with hard thermal loops also give $\Gamma_{sph} \sim \mathcal{O}(10) \alpha_2^5 T^4$ [97]

2.3.5 Washing out $B + L$

In the early Universe the transitions $\Delta N_{CS} = +1$ and $\Delta N_{CS} = -1$ are equally probable. The ratio of rates for the two transitions is given by

$$\frac{\Gamma_{sph+}}{\Gamma_{sph-}} = e^{-\Delta F/T}, \quad (40)$$

where ΔF is the free energy difference between the two vacua. Because of a finite $B + L$ density, there is a net chemical potential μ_{B+L} . Therefore

$$\Delta F \sim \mu_{B+L}^2 T^2 + \mathcal{O}(T^4) \equiv \frac{n_{B+L}^2}{T^2} + \mathcal{O}(T^4). \quad (41)$$

One then obtains [87]

$$\frac{dn_{B+L}}{dt} = \Gamma_{sph} + - \Gamma_{sph} - \sim N_g \frac{\Gamma_{sph}}{T^3} n_{B+L}. \quad (42)$$

It then follows that an exponential depletion of n_{B+L} due to sphaleron transitions remains active as long as

$$\frac{\Gamma_{sph}}{T^3} \geq H \quad \Rightarrow \quad T \leq \alpha_2^4 \frac{M_P}{g_*^{1/2}} \sim 10^{12} \text{ GeV}. \quad (43)$$

This result is important because it suggests that below $T = 10^{12}$ GeV, the sphaleron transitions can wash out any $B + L$ asymmetry being produced earlier in a time scale $\tau \sim (T^3/N_g \Gamma_{sph})$. This seems to wreck GUT baryogenesis based on $B - L$ conserving groups such as the minimal $SU(5)$.

2.4 Alternatives for baryogenesis

There are several scenarios for baryogenesis (for reviews on baryogenesis, see [78, 90, 91]), the main contenders being GUT baryogenesis, electroweak baryogenesis, leptogenesis, and baryogenesis through the decay of a field condensate, or Affleck-Dine baryogenesis. Here we give a brief description of these various alternatives.

2.4.1 GUT-baryogenesis

This was the first concrete attempt of model building on baryogenesis which incorporates out-of-equilibrium decays of heavy GUT gauge bosons $X, Y \rightarrow qq$, and $X, Y \rightarrow \bar{q}\bar{l}$ (see e.g., [78, 79, 88, 98]). The decay rate of the gauge boson goes as $\Gamma_X \sim \alpha_X M_X$, where M_X is the mass of the gauge boson and $\alpha_X^{1/2}$ is the GUT gauge coupling. Assuming that the Universe was in thermal equilibrium at the GUT scale, the decay temperature is given by

$$T_D \approx g_*^{-1/4} \alpha_X^{1/2} (M_X M_P)^{1/2}, \quad (44)$$

which is smaller than the gauge boson mass. Thus, at $T \approx T_D$, one expects $n_X \approx n_{\bar{X}} \approx n_\gamma$, and hence the net baryon density is proportional to the photon number density $n_B = \Delta B n_\gamma$. However below T_D the gauge boson abundances decrease and eventually they go out-of-equilibrium. The net entropy generated due to their decay heats up the Universe with a temperature which we denote here by T_{rh} . Let us naively assume that the energy density of the Universe at T_D is dominated by the X bosons with $\rho_X \approx M_X n_X$, and their decay products lead to radiation with an energy density $\rho = (\pi^2/30)g_*T_{rh}^4$, where $g_* \sim \mathcal{O}(100)$ for $T \geq M_{GUT}$. Equating the expressions for the two energy densities one obtains

$$n_X \approx \frac{\pi^2}{30} g_* \frac{T_{rh}^4}{M_X}. \quad (45)$$

Therefore the net baryon number comes out to be

$$B \equiv \frac{n_B}{s} \approx \frac{\Delta B n_X}{g_* n_\gamma} \approx \frac{3}{4} \frac{T_{rh}}{M_X} \Delta B. \quad (46)$$

T_{rh} is determined from the relation $\Gamma_X^2 \approx H^2(T_D) \sim (\pi^2/90)g_*T_{rh}^4/M_P^2$. Thus,

$$B \approx \left(\frac{g_*^{-1/2} \Gamma_X M_P}{M_X^2} \right)^{1/2} \Delta B \equiv \left(\frac{g_*^{-1/2} \alpha_X M_P}{M_X} \right)^{1/2} \Delta B. \quad (47)$$

Uncertainties in C and CP violation are now hidden in ΔB , but can be tuned to yield total $B \sim 10^{-10}$ in many models.

Above we have tacitly assumed that the Universe is in thermal equilibrium when $T \geq M_X$. This might not be true, since for $2 \leftrightarrow 2$ processes the scattering rate is given by $\Gamma \sim \alpha^2 T$, which becomes smaller than H at sufficiently high temperatures. Elastic $2 \rightarrow 2$ processes maintain thermal contact typically only up to a maximum temperature $\sim 10^{14}$ GeV, while chemical equilibrium is lost already at $T \sim 10^{12}$ GeV [99, 100]. It has been argued that QCD gas, which becomes asymptotically free at high temperatures, never reaches a chemical equilibrium above $\sim 10^{14}$ GeV [101]. In supergravity the maximum temperature of the thermal bath should not exceed 10^{10} GeV [102] (see Sect. 3.6.3).

2.4.2 Electroweak baryogenesis

A popular baryogenesis candidate is based on the electroweak phase transition, during which one can in principle meet all the Sakharov conditions. There is the sphaleron-induced baryon number violation above the critical temperature, various sources of CP violation, and an out-of-equilibrium environment if the phase transition is of the first order. In that case bubbles of broken $SU(2) \times U(1)_Y$ are nucleated into a symmetric background with a Higgs field profile that changes through the bubble wall [93, 103, 24].

There are two possible mechanisms which work in a different regime; local and non-local baryogenesis. In the local case both CP violation and baryon number violation takes place near the bubble wall. This requires the velocity of the bubble wall to be greater than the speed of the sound in the plasma [104, 105], and the electroweak phase transition to be strongly first order with thin bubble walls.

The second alternative, where the bubble wall velocity speed is small compared to the sound speed in the plasma, appears to be more realistic. In this mechanism the fermions, mainly the top quark and the tau-lepton, undergo CP violating interactions with the bubble wall, which results in a difference in the reflection and the transmission probabilities for the left and right chiral fermions. The net outcome is an overall chiral flux into the unbroken phase from the broken phase. The flux is then converted into baryons via sphaleron transitions inside the unbroken phase. The interactions are taking place in a thermal equilibrium except for the sphaleron transitions, the rate of which is slower than the rate at which the bubble sweeps the space.

One great difficulty with the electroweak baryogenesis is the smallness of CP violation in the SM. It has been pointed out that an additional Higgs doublet [106, 107, 105, 108] would provide an extra source for CP violation in the Higgs sector. However, the situation is much improved in the MSSM where there are two Higgs doublets H_u and H_d , and two important sources of CP violation [109]. The Higgses couple to the charginos and neutralinos at one loop level leading to a CP violating contribution. There is also a new source of CP violation in the mass matrix of the top squarks which can give rise to considerable CP violation [110].

Bubble nucleation depends on the thermal tunneling rate, and the expansion rate

of the Universe. The tunneling rate has to overcome the expansion rate in order to have a successful phase transition via bubble nucleation at a given critical temperature $T_c > T_t > T_0$. The actual value of the baryon asymmetry produced at the electroweak baryogenesis is still an open debate [111, 40, 112, 113, 42], but in general it is hard to generate a large baryon asymmetry. For $T_c \sim 100$ GeV, $N = 3$, $\alpha_2 = 0.033$, and $B(\lambda/g_2) \sim 1.87$, one obtains the condition [114, 87, 115, 26]

$$\frac{E_{sph}(T_c)}{T_c} \geq 7 \log \left[\frac{E_{sph}(T_c)}{T_c} \right] + 9 \log(10) + \log(\kappa). \quad (48)$$

which implies [115]

$$\frac{E_{sph}(T_c)}{T_c} \geq 45 \text{ for } \kappa = 10^{-1}, \quad (49)$$

$$\geq 37 \text{ for } \kappa = 10^{-4}. \quad (50)$$

The standard bound is often taken to be that of Eq. (49). In terms of the Higgs field value at T_c , one then obtains from Eq. (35)

$$\frac{\Phi(T_c)}{T_c} = \frac{g_2}{4\pi B(\lambda/g_2)} \frac{E_{sph}(T_c)}{T_c} \sim \frac{1}{36} \frac{E_{sph}(T_c)}{T_c}, \quad (51)$$

for the above values of α_2, B . Then the bounds in Eqs. (49,50) translate to

$$\frac{\Phi(T_c)}{T_c} \geq 1.3 \quad (1), \quad (52)$$

where the number in parenthesis is for Eq. (50). Eq. (52) respectively, implies that the phase transition should be strongly first order in order that sphalerons do not wash away all the produced baryon asymmetry. This result is the main constraint on electroweak baryogenesis.

In order to save the SM electroweak baryogenesis some attempts such as matter induced effects have also been invoked; exciting the SM gauge degrees of freedom in a time varying Higgs background [116, 117], or via dynamical scalar field which couples to the SM fields [118].

Lattice studies suggest that in the SM the phase transition is strongly first order only below Higgs mass $m_H \sim 72$ GeV [27, 119, 120, 121]. Above this scale the transition is just a cross-over. Such a Higgs mass is clearly excluded by the LEP measurements [65], thus excluding electroweak baryogenesis within the SM.

2.4.3 Electroweak baryogenesis in MSSM

In the MSSM the ratio $\Phi(T_c)/T_c$ can increase considerably. The MSSM Higgs sector at finite temperature has been considered in [122, 123, 124, 125], for lattice studies see [126, 127, 128]. In the MSSM the right handed stop $\bar{t}_R$ couples to the Higgs with a large Yukawa coupling. This leads to a strong first order phase transition [123, 124, 125]. The LEP precision tests then indicate that the lightest left handed stop should be heavy heavy with $m_Q \geq 500$ GeV. This implies the for lightest right handed stop mass

$$m_{\tilde{t}}^2 \approx m_U^2 + 0.15M_Z^2 \cos(2\beta) + m_t^2 \left(1 - \frac{\tilde{A}_t^2}{m_Q^2}\right), \quad (53)$$

where $\tilde{A}_t = A_t - \mu/\tan(\beta)$ is the stop mixing parameter, and μ is the soft-SUSY breaking mass parameter for the right-handed stop. The coefficient β of the cubic term βTH^3 in the effective potential reads

$$\beta_{MSSM} \approx \beta_{SM} + \frac{h_t^3 \sin^3(\beta)}{4\sqrt{2}\pi} \left(1 - \frac{\tilde{A}_t^2}{m_Q^2}\right)^{3/2}, \quad (54)$$

and can be at least one order of magnitude larger than β_{SM} . In principle this modification can give rise to a strong enough first order phase transition.

The sphaleron bound implies Higgs and stop masses in the range [41, 42]

$$110 \text{ GeV} \leq m_H \leq 115 \text{ GeV}, \quad \text{and} \quad 105 \text{ GeV} \leq m_{\tilde{t}_R} \leq 165 \text{ GeV}. \quad (55)$$

The present LEP constraint on the Higgs mass is $m_H \geq 115$ GeV [129]. Hence, even an MSSM-based electroweak baryogenesis may be at the verge of being ruled out. The definitive test of the MSSM based electroweak baryogenesis will obviously come from the Higgs and the stop searches at the LHC and the Tevatron [40, 130, 41, 111, 42].

2.4.4 Leptogenesis

Even if $B+L$ is completely erased by the sphaleron transitions, a net baryon asymmetry in the Universe can still be generated from a non-vanishing $B-L$ [131], even if there were no baryon number violating interactions. Lepton number violation alone can produce baryon asymmetry $B \sim -L$ [132], a process which is known as leptogenesis

(for a recent review [133], and references therein). For lepton number violation one however has to go beyond the SM.

A popular example is $SO(10)$ GUT model, which can either be broken into $SU(5)$ and then subsequently to the SM, or into the SM gauge group directly. The most attractive aspect of $SO(10)$ is that it is left-right symmetric (for details, see [134, 135]), and has a natural foundation for the see-saw mechanism [32, 33] as it incorporates a singlet right-handed neutrino N_R with a mass M_R . A lepton number violation appears when the Majorana right handed neutrino decays into the SM lepton doublet and Higgs doublet, and their CP conjugate state through

$$N_R \rightarrow \Phi + l, \quad N_R \rightarrow \bar{\Phi} + \bar{l}, \quad (56)$$

There also exist $\Delta L = 0$, and $\Delta L = 2$ processes mediated by the right handed neutrino through

$$\frac{(l\Phi)(l\Phi)}{M_R}, \quad \frac{l\Phi\Phi}{M_R}, \quad (57)$$

which are dimension 5 operators [136, 137]. (There are other processes involving t-quarks which may also be important [138, 139]). CP asymmetry is generated through the interference between tree level and one-loop diagrams.

The total baryon asymmetry and total lepton asymmetry can be found in terms of the chemical potentials as [140]

$$B = \sum_i (2\mu_{qi} + \mu_{u_Ri} + \mu_{d_Ri}), \quad L = \sum_i (2\mu_{li} + \mu_{e_Ri}), \quad (58)$$

where i denotes three leptonic generations. The Yukawa interactions establish an equilibrium between the different generations ($\mu_{li} = \mu_l$ and $\mu_{qi} = \mu_q$, etc.), and one obtains expressions for B and L in terms of the number of colors $N = 3$, and the number of charged Higgs fields N_H

$$B = -\frac{4N}{3}\mu_l, \quad L = \frac{14N^2 + 9NN_H}{6N + 3N_H}\mu_l, \quad (59)$$

together with a relationship between B and $B - L$ [140]

$$B = \left(\frac{8N + 4N_H}{22N + 13N_H} \right) (B - L). \quad (60)$$

A similar expression has also been found in [141, 88, 133], although there seems to be small of order one differences. The baryon asymmetry based on the decays of right handed neutrinos in a thermal bath has been computed in [138, 142, 143, 144, 145]. In a recent analysis [145], it was pointed out that the baryogenesis scale is tightly constrained together with the heavy right handed neutrino mass $T_B \sim M_{1,R} = \mathcal{O}(10^{10})$ GeV, with an upper bound on the light neutrino masses $\sum_i m_i < \sqrt{3}$ eV. The current bound on the right handed neutrino mass is around $M_R \sim \mathcal{O}(10^{11})$ GeV for light neutrino masses $m_{1\nu} \approx m_{2\nu} \approx m_{3\nu} \sim \mathcal{O}(0.1)$ eV.

High scale leptogenesis is ruled out in a supersymmetric theory because of the gravitino problem (see Sect. 3.7.1). However, if the masses of the right handed neutrinos are such that the mass splitting is comparable to their decay widths, it is possible to obtain an enhancement in the CP phase of order one [144], and possibly a low scale thermal leptogenesis [146]. Otherwise, one could resort to non-thermal leptogenesis [147, 136, 137, 148], or, to the scattering process discussed in [149], or to sneutrino driven leptogenesis [150, 137].

2.4.5 Baryogenesis through field condensate decay

Scalar condensates may have formed in the course of the evolution of the early Universe. In particular, during inflation all scalar fields are subject to fluctuations driven by the non-zero inflaton energy density so that fields with very shallow potentials may easily take non-zero values. An example is the MSSM, where for the squark and slepton fields there are several directions in the field space where the potential vanishes completely [46, 43]. These directions are called (perturbatively) flat. Field fluctuations along such flat directions will soon be smoothed out by inflation, which effectively stretches out any gradients, and only the zero mode, or the scalar condensate, remains. This mechanism is quite general and applicable to any order parameter with flat enough potential. Baryogenesis can then be achieved by the decay of a condensate that carries baryonic charge, as was first pointed out by Affleck and Dine [44]. As we will discuss, the flat direction condensate can get dynamically charged with a large B and/or L by virtue of CP -violating self-couplings.

Baryogenesis from MSSM flat directions has the virtue that it only requires two already quite popular paradigms: supersymmetry and inflation. In the old version [44] baryons were produced by a direct decay of the condensate, to be discussed in Sect. 2.5.2. It was however pointed out first by Kusenko and Shaposhnikov [151] in the case of gauge mediated supersymmetry breaking, and then by Enqvist and McDonald in [48] in the case of gravity mediated supersymmetry breaking, that the MSSM flat direction condensate most often is not stable but fragments and eventually forms non-topological solitons called Q -balls [47]. These issues will be dealt in Sects. 6 and 7.

2.5 Old Affleck-Dine baryogenesis

2.5.1 Classical motion of the order parameter

In the original Affleck-Dine baryogenesis [44] it was assumed that the order parameter along the flat direction is displaced from the origin because of inflationary fluctuations. Because of inflation, only the long wave-length model of the order parameter will survive so that a spatially constant condensate field is formed along the flat direction. This we shall sometimes call the Affleck-Dine (AD) field. In an expanding Universe the coherent AD field ϕ obeys the usual equation of motion,

$$\ddot{\phi} + 3H\dot{\phi} + \frac{\partial V}{\partial \phi} = 0 , \quad (61)$$

where H is the Hubble parameter.

To follow the time evolution of the AD field, let us consider a toy model with the potential

$$V(\phi) = m^2|\phi|^2 + \lambda(\phi^4 + \phi^{*4}) + \frac{|\phi|^6}{M^2} + \dots . \quad (62)$$

Although this potential is unrealistic in that it does not take correctly into account of supersymmetry breaking induced by the non-zero cosmological constant of the inflationary era, it nevertheless captures the main features of the initial cosmological evolution of the AD field.

The theory Eq. (62) has a partially conserved current $j_\mu = i\phi^*\partial_\mu\phi$, with

$$\partial_\mu j^\mu = \partial_\mu(i\phi^*\partial^\mu\phi - i\partial^\mu\phi^*\phi) = i\lambda(\phi^{*4} - \phi^4) . \quad (63)$$

The current is conserved for small ϕ . The role of the higher order term $|\phi|^6$ is just to stabilize the potential. In the toy model Eq. (62), we identify the baryon number density n_B with j_0 . The model also has a CP invariance under which $\phi \leftrightarrow \phi^*$ but which is violated by the initial conditions, which are taken to be

$$\phi = i\phi_0, \quad \dot{\phi} = 0, \quad (64)$$

where ϕ_0 is real. Writing $\phi = \phi_R + i\phi_I$ one finds the coupled equations of motion (see e.g., [152])

$$\begin{aligned} \ddot{\phi}_I + 3H\dot{\phi}_I + \left[m^2 + 12\lambda\phi_R\phi_I + \frac{3|\phi|^4}{M^2} \right] \phi_I &= 4\lambda\phi_R^3 \\ \ddot{\phi}_R + 3H\dot{\phi}_R + \left[m^2 + \frac{3|\phi|^4}{M^2} \right] \phi_R &= 4\lambda(3\phi_I\phi_R^2 - \phi_I^3). \end{aligned} \quad (65)$$

In a matter dominated Universe $H = 2/(3t)$, so that for large times $t \gg m^{-1}$ the motion is damped and Eq. (65) has then oscillatory solutions of the form

$$\phi_k = \frac{A_k}{mt} \sin(mt + \delta_k), \quad k = I, R, \quad (66)$$

where the amplitudes A_k and the phases δ_k depend on the parameters m, λ, M , and the initial conditions Eq. (64). For large times the baryon number is then found to be

$$n_B = 2(\phi_I\dot{\phi}_R - \phi_R\dot{\phi}_I) = \frac{2A_IA_R}{mt^2} \sin(\delta_I - \delta_R). \quad (67)$$

If $\phi_0^2 \ll mM$, as was tacitly assumed by Affleck and Dine [44], one may disregard the higher-order terms. In that case one obtains [45] $A_I = \phi_0$ and $A_R = a_R\lambda\phi_0^3/m^2$, where $a_R = 0.85$ is determined numerically. Likewise, numerically one finds that $\delta_I - \delta_R = 1.54$. Thus,

$$n_B = \frac{1.7\lambda\phi_0^4}{m^3t^2}, \quad (68)$$

and the generated baryon number per particle is

$$R = \frac{mn_B}{\rho_\phi} = \frac{1.7\lambda\phi_0^2}{m^2}. \quad (69)$$

Eq. (69) is true for matter dominated Universe; for radiation dominated Universe one obtains a similar result, but the numerical prefactor 1.7 should be replaced by -1.3 .

If $\phi_0^2 > mM$, one finds $A_I = a_I(mM)^{1/2}$ and $A_R = a_R(M^3/m)^{1/2}$ with $a_I = 0.94$, $a_R = -2.86$, $\delta_I = 0.11$, and $\delta_R = -0.41$. It then follows from Eq. (67), that

$$n_B = -\frac{2.7\lambda M^2}{mt^2} . \quad (70)$$

Thus, the baryon generation mechanism is remarkably robust. The initial conditions do not matter, nor the actual expansion rate of the Universe. The baryon number generated per ϕ -particle is always large and, with $\lambda \sim m^2/\langle\phi\rangle^2$, typically $n_B \gg 1$. Although these conclusions were derived in a toy model, similar results hold true also for the MSSM flat directions.

Thus, to summarize, along a flat direction where squarks and sleptons have non-zero expectation values, evolution of the AD field condensate, starting from a CP violating initial value, will dynamically generate large baryon number density and charge the condensate with B and/or L .

2.5.2 Condensate decay

To provide the Universe with the observed baryon to entropy ratio, $n_B/s \sim 10^{-10}$, the AD condensate must eventually transform itself into ordinary quarks. Originally [44], it was thought that this could happen via the decay of the AD field components (squarks and sleptons) to ordinary quarks and leptons. The AD condensate can be thought of as a coherent state of ϕ -particles where $\phi = \phi_0 e^{imt}$ and $|\phi_0| \gg m$. When supersymmetry breaking is switched on, the AD field starts to oscillate about the old vacuum $\langle\phi\rangle \gg m$. Writing $\phi = \langle\phi\rangle + \phi'$, one observes that all the fields to which the excitations ϕ' couple are heavy with masses $\mathcal{O}(\langle\phi\rangle)$. The field ϕ' itself has a mass of the order of supersymmetry breaking, $\mathcal{O}(m)$. Therefore, ϕ' can decay to light fields only through loop diagrams involving heavy fields, with an effective coupling of the type $(g^2/\langle\phi\rangle)\phi'\psi\partial\psi^\dagger$, where ψ is a light fermion and g some coupling constant. The decay rate is thus, [44]

$$\Gamma \sim g^4 \frac{m^3}{\phi^2} . \quad (71)$$

Because of the oscillations of the AD field, the Universe will eventually become dominated by the energy density in the oscillations, $\rho_\phi \simeq m^2\phi^2$, so that $H \sim \rho_\phi^{1/2}/M_P$.

The AD field will decay when $\Gamma \simeq H$, or $\phi \simeq (m^2 M_P)^{1/3}$. This implies a reheating temperature $T_{rh} \simeq s^{1/3} \simeq \rho_\phi^{1/4}$ while the baryon number density is $n_B = R\rho_\phi/m$, where R is given in Eq. (69). Therefore one finally obtains

$$\frac{n_B}{s} \simeq \frac{\lambda\phi_0^2}{m^2} \left(\frac{M}{m}\right)^{1/6}. \quad (72)$$

Depending on λ , and the size of the initial fluctuation ϕ_0 of the AD condensate, n_B/s can be either small or large. Therefore determining the initial value is of utmost importance [153]. This requires us to consider theories of inflation in more detail, which will be done in the next Section. Following that, we shall discuss the disappearance of the AD condensate by fragmentation into (quasi)stable lumps of condensate matter, whose state of lowest energy is a spherical non-topological soliton, a Q -ball [47].

3 Field fluctuations during inflation

Apart from explaining the initial condition for the hot Big Bang model, the flatness problem, and the horizon problem, cosmological inflation [154, 13, 14] is one of the most favored candidate for the origin of structure in the Universe (for reviews on inflation, see [15, 155]). There are many models of inflation, but by far the simplest is one in which inflation is generated by the large energy density of a scalar field. The scalar field driven inflation not only explains the homogeneity and the flatness problems but also the observed scale invariance of the density perturbations.

Inflation based on a scalar field theory is described by the following Lagrangian:

$$\mathcal{L} = \frac{M_{\text{P}}^2}{2}R + \frac{1}{2}\partial_\mu\phi\partial^\mu\phi - V(\phi), \quad (73)$$

where R is the curvature scalar. The energy-momentum tensor reads

$$T_{\mu\nu} = \partial_\mu\phi\partial_\nu\phi - \frac{1}{2}g_{\mu\nu}\partial_\rho\phi\partial^\rho\phi - g_{\mu\nu}V(\phi) \quad (74)$$

so that the energy density and the pressure are given by

$$\rho \equiv T_{00} = \frac{1}{2}\dot{\phi}^2 + \frac{1}{2a^2(t)}(\nabla\phi)^2 + V(\phi), \quad (75)$$

$$p \equiv \frac{T_{ii}}{a^2(t)} = \frac{1}{2}\dot{\phi}^2 - \frac{1}{6a^2(t)}(\nabla\phi)^2 - V(\phi). \quad (76)$$

One of the initial conditions for inflation is that there must be a homogeneous patch of the Universe which is bigger than the size of the Hubble horizon [156] (also supported by numerical studies, see [157]). However such a stringent condition can be evaded in a chaotic inflation beginning at the Planck scale [158, 159, 15, 169]. More complicated situation can be obtained if there are several fields that participate in inflation; the classic example is assisted inflation [161, 162].

3.1 Fluctuation spectrum in de Sitter space

The plane wave solution of a massive scalar field $\phi(\mathbf{x}, t)$ in a spatially flat Robertson-Walker metric can be decomposed into Fourier modes by

$$\phi = \frac{1}{(2\pi)^{3/2}} \int d^3k \left(\phi_k(t) e^{ik\cdot\mathbf{x}} + \text{h.c.} \right). \quad (77)$$

Solving the Klein-Gordon equation for the scalar field in a conformal metric: $ds^2 = g_{\mu\nu}dx^\mu dx^\nu = a^2(\tau, x)(d\tau^2 - dx^2)$, the mode function can be given by [163, 164, 165, 166]

$$\begin{aligned}\phi_k(\tau) &= \left(\frac{\pi}{4}\right)^{1/2} H|\tau|^{3/2} \left(c_1 H_\nu^{(1)}(k\tau) + c_2 H_\nu^{(2)}(k\tau)\right), \\ \tau &= -H^{-1}e^{-Ht}, \quad \text{and} \quad \nu^2 = \frac{9}{4} - \frac{m^2}{H^2},\end{aligned}\tag{78}$$

where m is the mass of the scalar field, $H_\nu^{(1)}$ and $H_\nu^{(2)}$ are the Hankel functions and c_1, c_2 are constants. The readers might be tempted to take the limit $\tau \ll 0$, in order to match the above solution with the plane wave solution in a Minkowski background. However this leads to a quasi static de Sitter solution [167]. More technically, it has been shown that using a point splitting regularization scheme, it is possible to obtain a Bunch-Davies vacuum for a de Sitter background which actually corresponds to taking $c_1 = 0$, and $c_2 = 1$. A simple but intuitive way has been developed in [168], where it has been argued that during a de Sitter phase, the main contribution to the two point correlation function comes from the long wavelength modes; $k|\tau| \ll 1$ or $k \ll H \exp(Ht)$. Therefore the two point function is defined by an infrared cutoff which is determined by the Hubble expansion [168]

$$\langle \phi^2 \rangle \approx \frac{1}{(2\pi)^3} \int_H^{He^{Ht}} d^3k |\phi_k|^2.\tag{79}$$

The result of the integration yields [166, 165, 167, 168] an indefinite increase in the variance with time

$$\langle \phi^2 \rangle \approx \frac{H^3}{4\pi^2} t.\tag{80}$$

This result can also be obtained by considering the Brownian motion of the scalar field [160].

For a massive field with $m \ll H$, and $\nu \neq 3/2$, one does not obtain an indefinite growth of the variance of the long wavelength fluctuations, but [166, 165, 167, 168]

$$\langle \phi^2 \rangle = \frac{3H^4}{8\pi^2 m^2} \left(1 - e^{-(2m^2/3H^2)t}\right).\tag{81}$$

In the limiting case when $m \rightarrow H$, the variance goes as $\langle \phi^2 \rangle \approx H^2$. In the limit $m \gg H$, the variance goes as $\langle \phi^2 \rangle \approx (H^3/12\pi^2 m)$ [167]. Only in a massless case $\langle \phi^2 \rangle$ can be treated as a homogeneous background field with a long wavelength mode. This

result plays an important role for the rest of this review as it implies that in a de Sitter phase any scalar field, including the AD condensate, are subject to quantum fluctuations.

3.2 Slow roll inflation

A completely flat potential can render inflation eternal, provided the energy density stored in the flat direction dominates. The inflaton direction is however not completely flat but has a potential $V(\phi)$ with some slope. An inflationary phase is obtained while

$$H^2 \approx \frac{1}{3M_{\text{P}}^2} V(\phi), \quad (82)$$

$$3H\dot{\phi} \approx -V'(\phi), \quad (83)$$

where prime denotes derivative with respect to ϕ . In the above the approximations are: $\dot{\phi}^2 < V(\phi)$, and $\ddot{\phi} < V'(\phi)$, which lead to the slow roll conditions (see e.g. [9])

$$\epsilon(\phi) = \frac{M_{\text{P}}^2}{2} \left(\frac{V'}{V} \right)^2 \ll 1, \quad (84)$$

$$|\eta(\phi)| = M_{\text{P}}^2 \frac{V''}{V} \ll 1. \quad (85)$$

Note that ϵ is positive by definition.

These conditions are necessary but not sufficient for inflation. They only constrain the shape of the potential but not the velocity of the field $\dot{\phi}$. Therefore a tacit assumption behind the success of the slow roll conditions is that the inflaton field should not have a large initial velocity.

Inflation comes to an end when the slow roll conditions are violated, $\epsilon \sim 1$, and $\eta \sim 1$. However, there are certain models where this need not be true, for instance in hybrid inflation models [169], where inflation comes to an end via a phase transition, or in oscillatory models of inflation where slow roll conditions are satisfied only on average [170].

One of the salient features of the slow roll inflation is that there exists a late time attractor behavior. This means that during inflation the evolution of a scalar field at a given field value has to be independent of the initial conditions. Therefore slow roll

inflation should provide an attractor behavior which at late times leads to an identical field evolution in the phase space irrespective of the initial conditions [171]. In fact the slow roll solution does not give an exact attractor solution to the full equation of motion but is nevertheless a fairly good approximation [171]. A similar statement has been proven for multi-field exponential potentials without slow roll conditions (i.e., assisted inflation) [161]. The attractor behavior of the inflaton leads to powerful predictions which can be distinguished from other candidates of galaxy formation [19].

The standard definition of the number of e-foldings is given by

$$N \equiv \ln \frac{a(t_{end})}{a(t)} = \int_t^{t_{end}} H dt \approx \frac{1}{M_{\text{P}}^2} \int_{\phi_{end}}^{\phi} \frac{V}{V'} d\phi, \quad (86)$$

where ϕ_{end} is defined by $\epsilon(\phi_{end}) \sim 1$, provided inflation comes to an end via a violation of the slow roll conditions. The number of e-foldings can be related to the Hubble crossing mode $k = a_k H_k$ by comparing with the present Hubble length $a_0 H_0$. The final result is [9]

$$N(k) = 62 - \ln \frac{k}{a_0 H_0} - \ln \frac{10^{16} \text{ GeV}}{V_k^{1/4}} + \ln \frac{V_k^{1/4}}{V_{end}^{1/4}} - \frac{1}{3} \ln \frac{V_{end}^{1/4}}{\rho_{rh}^{1/4}} \quad (87)$$

where the subscripts *end* (*rh*) refer to the end of inflation (onset of reheating). The details of the thermal history of the Universe determine the precise number of e-foldings, but for most practical purposes it is sufficient to assume that $N(k) \approx 50$, keeping all the uncertainties such as the scale of inflation and the end of inflation within a margin of 10 e-foldings. A significant modification can take place only if there is an epoch of late inflation such as thermal inflation [172], or in theories with a low quantum gravity scale [173].

3.3 Primordial density perturbations

Initially, the theory of cosmological perturbations has been developed in the context of FRW cosmology [175], and for models of inflation in [176, 177, 178, 179, 180]. For a complete review on this topic, see [16]. For a real single scalar field there arise only adiabatic density perturbations. In case of several fluctuating fields there will in general also be isocurvature perturbations. We briefly describe the two perturbations and their observational differences.

3.3.1 Adiabatic perturbations and the Sachs-Wolfe effect

Let us consider small inhomogeneities $\phi(\mathbf{x}, t) = \phi(t) + \delta\phi(\mathbf{x}, t)$ such that $\delta\phi \ll \phi$. Perturbations in matter densities automatically induce perturbations in the background metric, but the separation between the background metric and a perturbed one is not unique. One needs to choose a gauge. A simple choice would be to fix the observer to the unperturbed matter particles, where the observer will detect a velocity of matter field falling under gravity; this is known as the Newtonian or the longitudinal gauge because the observer in the Newtonian gravity limit measures the gravitational potential well where matter is falling in and clumping. The induced metric can be written as

$$ds^2 = a^2(\tau) \left[(1 + 2\Phi)d\tau^2 - (1 - 2\Psi)\delta_{ik}dx^i dx^k \right], \quad (88)$$

where Φ has a complete analogue of Newtonian gravitational potential. In the case when the spatial part of the energy momentum tensor is diagonal, i.e. $\delta T_j^i = \delta_j^i$, it follows that $\Phi = \Psi$ [16]. Right at the time of horizon crossing one finds a solution for $\delta\phi$ as

$$\langle |\delta\phi_k|^2 \rangle = \frac{H(t_*)^2}{2k^3}, \quad (89)$$

where t_* denotes the instance of horizon crossing. Correspondingly, we can also define a power spectrum

$$\mathcal{P}_\phi(k) = \frac{k^3}{2\pi^2} \langle |\delta\phi_k|^2 \rangle = \left[\frac{H(t_*)}{2\pi} \right]^2 \equiv \left[\frac{H}{2\pi} \right]^2 \Big|_{k=aH}. \quad (90)$$

Note that the phase of $\delta\phi_k$ can be arbitrary, and therefore, inflation has generated a Gaussian perturbation.

In the limit $k \rightarrow 0$, one can find an exact solution for the long wavelength inhomogeneities $k \ll aH$ [181, 16], which reads

$$\Phi_k \approx c_1 \left(\frac{1}{a} \int_0^t a dt' \right) + c_2 \frac{H}{a}, \quad (91)$$

$$\frac{\delta\phi_k}{\phi} = \frac{1}{a} \left(c_1 \int_0^t a dt' - c_2 \right), \quad (92)$$

where the dot denotes derivative with respect to physical time. The growing solutions are proportional to c_1 , the decaying proportional to c_2 . Concentrating upon the growing solution, it is possible to obtain a leading order term in an expansion with the help

of the slow roll conditions:

$$\Phi_k \approx -c_1 \frac{\dot{H}}{H^2}, \quad (93)$$

$$\frac{\delta\phi_k}{\dot{\phi}} \approx \frac{c_1}{H}. \quad (94)$$

Note that at the end of inflation, which is indicated by $\ddot{a} = 0$, or equivalently by $\dot{H} = -H^2$, one obtains a constant Newtonian potential $\Phi_k \approx c_1$. This is perhaps the most significant result for a single field perturbation.

In a long wavelength limit one obtains a constant of motion ζ [180, 182, 16] defined as

$$\zeta = \frac{2}{3} \frac{H^{-1} \dot{\Phi}_k + \Phi_k}{1 + w} + \Phi_k, \quad w = \frac{p}{\rho}. \quad (95)$$

If the equation of state for matter remains constant there is a simple relationship which connects the metric perturbations at two different times [180, 182, 16]

$$\Phi_k(t_f) = \frac{1 + \frac{2}{3}(1 + w(t_f))^{-1}}{1 + \frac{2}{3}(1 + w(t_i))^{-1}} \Phi_k(t_i). \quad (96)$$

The comoving curvature perturbation [183] reads in the longitudinal gauge [16] for the slow roll inflation as

$$\mathcal{R}_k = \Phi_k - \frac{H^2}{\dot{H}} \left(H^{-1} \dot{\Phi}_k + \Phi_k \right). \quad (97)$$

For CMB and structure formation we need to know the metric perturbation during the matter dominated era when the metric perturbation is $\Phi(t_f) \approx (3/5)c_1$. Substituting the value of c_1 from Eq. (94), we obtain

$$\Phi_k(t_f) \approx \frac{3}{5} H \frac{\delta\phi_k}{\dot{\phi}} \Big|_{k=aH}. \quad (98)$$

In a similar way it is also possible to show that the comoving curvature perturbations is given by

$$\mathcal{R}_k \approx \frac{H}{\dot{\phi}} \delta\phi \Big|_{k=aH}, \quad (99)$$

where $\delta\phi$ denotes the field perturbation on a spatially flat hypersurfaces, because on a comoving hypersurface $\delta\phi = 0$, by definition. Therefore, on flat hypersurfaces

$$\delta\phi_k = \dot{\phi} \delta t, \quad (100)$$

where δt is the time displacement going from flat to comoving hypersurfaces [19, 9]. As a result

$$\mathcal{R}_k \equiv H\delta t. \quad (101)$$

Note that during matter dominated era the curvature perturbation and the metric perturbations are related to each other

$$\Phi_k = -\frac{3}{5}\mathcal{R}_k. \quad (102)$$

In the matter dominated era the photon sees this potential well created by the primordial fluctuation and the redshift in the emitted photon is given by

$$\frac{\Delta T_k}{T} = -\Phi_k. \quad (103)$$

At the same time, the proper time scale inside the fluctuation becomes slower by an amount $\delta t/t = \Phi_k$. Therefore, for the scale factor $a \propto t^{2/3}$, decoupling occurs earlier with

$$\frac{\delta a}{a} = \frac{2}{3} \frac{\delta t}{t} = \frac{2}{3} \Phi_k. \quad (104)$$

By virtue of $T \propto a^{-1}$ this results in a temperature which is hotter by

$$\frac{\Delta T_k}{T} = -\Phi_k + \frac{2}{3}\Phi_k = -\frac{\Phi_k}{3}. \quad (105)$$

This is the celebrated Sachs-Wolfe effect [184], which we shall revisit when discussing isocurvature fluctuations.

3.3.2 Spectrum of adiabatic perturbations

Now, one can immediately calculate the spectrum of the metric perturbations. For a critical density Universe

$$\delta_k \equiv \left. \frac{\delta \rho}{\rho} \right|_k = -\frac{2}{3} \left(\frac{k}{aH} \right)^2 \Phi_k, \quad (106)$$

where $\nabla^2 \rightarrow -k^2$, in the Fourier domain. Therefore, with the help of Eqs. (90,98), one obtains

$$\delta_k^2 \equiv \frac{4}{9} \mathcal{P}_\Phi(k) = \frac{4}{9} \frac{9}{25} \left(\frac{H}{\dot{\phi}} \right)^2 \left(\frac{H}{2\pi} \right)^2, \quad (107)$$

where the right hand side can be evaluated at the time of horizon exit $k = aH$. In fact the above expression can also be expressed in terms of curvature perturbations [19, 9]

$$\delta_k = \frac{2}{5} \left(\frac{k}{aH} \right)^2 \mathcal{R}_k, \quad (108)$$

and following Eq. (97), we obtain $\delta_k^2 = 4/25 \mathcal{P}_{\mathcal{R}}(k) = (4/25)(H/\dot{\phi})^2(H/2\pi)^2$, exactly the same expression as in Eq. (107). With the help of the slow roll equation $3H\dot{\phi} = -V'$, and the critical density formula $3H^2 M_{\text{P}} = V$, one obtains

$$\delta_k^2 \approx \frac{1}{75\pi^2 M_{\text{P}}^6} \frac{V^3}{V'^2} = \frac{1}{150\pi^2 M_{\text{P}}^4} \frac{V}{\epsilon}, \quad (109)$$

where we have used the slow roll parameter $\epsilon \equiv (M_{\text{P}}^2/2)(V'/V)^2$. The COBE satellite measured the CMB anisotropy and fixes the normalization of $\delta_{\Phi}(k)$ on a very large scale. For a critical density Universe, if we assume that the primordial spectrum can be approximated by a power law and ignoring gravitational waves:

$$\delta_{\Phi}(k) = 1.91 \times 10^{-5} \left(\frac{k}{k_{\text{pivot}}} \right)^{(n-1)/2}, \quad (110)$$

where n is the spectral index and $k_{\text{pivot}} = 7.5a_0H_0$ is the scale at which the normalization is independent of the spectral index.

The spectral index $n(k)$ is defined as

$$n(k) - 1 \equiv \frac{d \ln \mathcal{P}_{\Phi}}{d \ln k}. \quad (111)$$

This definition is equivalent to the power law behavior if $n(k)$ is fairly a constant quantity over a range of k of interest. The power spectrum can then be written as

$$\mathcal{P}_{\Phi}(k) \propto k^{n-1}. \quad (112)$$

If $n = 1$, the spectrum is flat and known as Harrison-Zeldovich spectrum [185]. For $n \neq 1$, the spectrum is tilted and $n > 1$ is known as blue spectrum. In terms of the slow roll parameters, one can write [171]

$$\frac{d\epsilon}{d \ln k} = 2\epsilon\eta - 4\epsilon^2, \quad \frac{d\eta}{d \ln k} = -2\epsilon\eta + \xi^2, \quad \frac{d\xi^2}{d \ln k} = -2\epsilon\xi^2 + \eta\xi^2 + \sigma^3, \quad (113)$$

where

$$\xi^2 \equiv M_{\text{P}}^4 \frac{V'(d^3V/d\phi^3)}{V^2}, \quad \sigma^3 \equiv M_{\text{P}}^6 \frac{V'^2(d^4V/d\phi^4)}{V^3}. \quad (114)$$

Thus one finds [186]

$$n - 1 = -6\epsilon + 2\eta. \quad (115)$$

Slow roll inflation requires that $\epsilon \ll 1$, $|\eta| \ll 1$, and therefore naturally predicts small variation in the spectral index within $\Delta \ln k \approx 1$. The recent Boomerang data suggest [10]

$$|n - 1| \leq 0.1. \quad (116)$$

The rate of change in η is also very small, and can be estimated in a similar way [187]

$$\frac{dn}{d \ln k} = -16\epsilon\eta + 24\epsilon^2 + 2\xi^2. \quad (117)$$

It is possible to extend the calculation of metric perturbation beyond the slow roll approximation basing on a formalism similar to that developed in [188, 189, 190, 191].

3.3.3 Gravitational waves

Gravitational waves are linearized tensor perturbations of the metric and do not couple to the energy momentum tensor. Therefore, they do not give rise a gravitational instability, but carry the underlying geometric structure of the space-time. The first calculation of the gravitational wave production was made in [192], and the topic has been considered by many authors [193]. For reviews on gravitational waves, see [16, 194].

The gravitational wave perturbations are described by a line element $ds^2 + \delta ds^2$, where

$$ds^2 = a^2(\tau)(d\tau^2 - dx^i dx_i), \quad \delta ds^2 = -a^2(\tau)h_{ij}dx^i dx^j. \quad (118)$$

The gauge invariant and conformally invariant 3-tensor h_{ij} is symmetric, traceless $\delta^{ij}h_{ij} = 0$, and divergenceless $\nabla_i h_{ij} = 0$ (∇_i is a covariant derivative). Massless spin 2 gravitons have two degrees of freedom and as a result are also transverse. This means that in a Fourier domain the gravitational wave has a form

$$h_{ij} = h_+ e_{ij}^+ + h_\times e_{ij}^\times. \quad (119)$$

For the Einstein gravity, the gravitational wave equation of motion follows that of a massless Klein Gordon equation [16]. Especially, for a flat Universe

$$\ddot{h}_j^i + 3H\dot{h}_j^i + \left(\frac{k^2}{a^2}\right)h_j^i = 0, \quad (120)$$

As any massless field, the gravitational waves also feel the quantum fluctuations in an expanding background. The spectrum mimics that of Eq. (90)

$$\mathcal{P}_{grav}(k) = \frac{2}{M_{\text{P}}^2} \left(\frac{H}{2\pi}\right)^2 \Big|_{k=aH}. \quad (121)$$

Note that the spectrum has a Planck mass suppression, which suggests that the amplitude of the gravitational waves is smaller compared to that of the adiabatic perturbations. Therefore it is usually assumed that their contribution to the CMB anisotropy is small. The corresponding spectral index can be calculated as [186]

$$n_{grav} = \frac{d \ln \mathcal{P}_{grav}(k)}{d \ln k} = -2\epsilon. \quad (122)$$

Note that the spectral index is negative.

3.4 Multi-field perturbations

In multi-field inflation models contributions to the density perturbations come from all the fields. However unlike in a single scalar case, in the multi-field case there might not be a unique late time trajectory corresponding to all the fields. This is true in particular for those fields that are effectively massless during inflation, such as the MSSM flat direction fields. Therefore, in these cases scalar perturbations will depend on the field trajectories and thus on the choice of initial conditions, with an ensuing loss of predictivity. In a very few cases it is possible to obtain a late time attractor behavior of all the fields; an example is assisted inflation [161]. Let us here nevertheless assume that there is an underlying unique late time trajectory resulting in a simple expression for the amplitude of the density perturbations and the spectral index [195, 155].

3.4.1 Adiabatic and isocurvature conditions

There are only two kinds of perturbations that can be generated. The first one is the adiabatic perturbation discussed previously; it is a perturbation along the late time classical trajectories of the scalar fields during inflation. When the primordial perturbations enter our horizon they perturb the matter density with a generic *adiabatic condition*, which is satisfied when the density contrast of the individual species is related to the total density contrast δ_k

$$\frac{1}{3}\delta_{kb} = \frac{1}{3}\delta_{kc} = \frac{1}{4}\delta_{k\nu} = \frac{1}{4}\delta_{k\gamma} = \frac{1}{4}\delta_k, \quad (123)$$

where b stands for baryons, c for cold dark matter, γ for photons and ν for neutrinos.

The other type is the isocurvature perturbation. During inflation this can be viewed as a perturbation orthogonal to the unique late time classical trajectory. Therefore, if there were N fluctuating scalar fields during inflation, there would be $N - 1$ degrees of freedom which would contribute to the isocurvature perturbation.

The *isocurvature condition* is known as $\delta\rho = 0$: the sum total of all the energy contrasts must be zero. The most general density perturbations is then given by a linear combination of an adiabatic and an isocurvature density perturbations.

3.4.2 Adiabatic perturbations due to multi-field

In a comoving gauge Eq. (97) with $\mathcal{R} = -H\delta\phi/\dot{\phi}$ holds good even for multi-field inflation models, provided we identify each field component of ϕ along the slow roll direction. There also exists a relationship between the comoving curvature perturbations and the number of e-foldings N [181, 196, 195, 155]

$$\mathcal{R} = \delta N = \frac{\partial N}{\partial \phi_a} \delta \phi_a, \quad (124)$$

where N is measured by a comoving observer while passing from flat hypersurface (which defines $\delta\phi$) to the comoving hypersurface (which determines $\mathcal{R}$) [195, 197]. The repeated indices are summed over and the subscript a denotes a component of the inflaton. A more intuitive derivation has been given in [155, 9].

If again one assumes that the perturbations in $\delta\phi_a$ have random phases with an amplitude $(H/2\pi)^2$, one obtains

$$\delta_k^2 = \frac{V}{75\pi^2 M_{\text{P}}^2} \frac{\partial N}{\partial \phi_a} \frac{\partial N}{\partial \phi_a}. \quad (125)$$

For a single component $\partial N/\partial \phi \equiv (M_{\text{P}}^{-2}V/V')$, and then Eq. (125) reduces to Eq. (109). By using slow roll equations we can again define the spectral index

$$n - 1 = -\frac{M_{\text{P}}^2 V_{,a} V_{,a}}{V^2} - \frac{2}{M_{\text{P}}^2 N_{,a} N_{,a}} + 2 \frac{M_{\text{P}}^2 N_{,a} N_{,b} V_{,ab}}{V N_{,c} N_{,c}}, \quad (126)$$

where $V_{,a} \equiv \partial V/\partial \phi_a$, and similarly $N_{,a} \equiv \partial N/\partial \phi_a$. For a single component we recover Eq. (115) from Eq. (126). These results prove useful in constraining the AD potential by cosmological density perturbations, as will be discussed in Sect. 5.3.

3.4.3 Isocurvature perturbations and CMB

One may of course simply assume a purely isocurvature initial condition. For any species the entropy perturbation is defined by

$$S_i = \frac{\delta n_i}{n_i} - \frac{\delta n_\gamma}{n_\gamma}, \quad (127)$$

Thus, if initially there is a radiation bath with a common radiation density contrast δ_r , a baryon-density contrast $\delta_b = 3\delta_r/4$, and a CDM density contrast δ_c , then

$$S = \delta_c - \frac{3}{4}\delta_r = \frac{\rho_r \delta \rho_c - (3/4)\rho_c \delta \rho_r}{\rho_r \rho_c} = \frac{\rho_r + (3/4)\rho_c}{\rho_r \rho_c} \delta \rho_c \approx \delta_c, \quad (128)$$

where we have used the isocurvature condition $\delta \rho_r + \delta \rho_c = 0$, and the last equality holds in a radiation dominated Universe.

However a pure isocurvature perturbation gives five times larger contribution to the Sachs-Wolfe effect compared to the adiabatic case [198, 19, 9]. This result can be derived very easily in a matter dominated era with an isocurvature condition $\delta \rho_c = -\delta \rho_r$, which gives a contribution $\mathcal{R}_k = (1/3)S_k$. Therefore from Eqs. (102,105), we obtain $\Delta T_k/T = -S_k/15$. There is an additional contribution from radiation because we are in a matter dominated era, see Eq. (128), $S \approx \delta_c \equiv -(3/4)\delta_r$. The sum total isocurvature perturbation $\Delta T_k/T = -S/15 - S/3 = -6S/15$, where S is measured on

the last scattering surface. The Sachs-Wolfe effect for isocurvature perturbations fixes the *slope* of the perturbations, rather than the amplitude [199, 200]. Present CMB data rules out pure isocurvature perturbation spectrum [201, 202], although a mixture of adiabatic and isocurvature perturbations remains a possibility [201, 203, 204, 205]. In the latter case it has been argued that the adiabatic and isocurvature perturbations might naturally turn out to be correlated [206, 207]. The most general power spectrum is not a single function but a 5×5 matrix, which contains all possible adiabatic and isocurvature perturbations together with their cross-correlations. As discussed in [207], resolving the perturbation spectrum in all its generality would be an observational challenge that probably would have to wait for the determination of the polarization spectrum by the Planck Surveyor Mission.

It is sometimes useful to consider the ratio α of the total power spectra, defined as $\mathcal{P}_{tot} = \mathcal{P}_{ad} + \mathcal{P}_{iso}$ [208, 204], where α is defined as [204]

$$\alpha = \frac{16}{25} \frac{\mathcal{P}_{iso}}{\mathcal{P}_{ad}} \Big|_{k=aH} = \left| \frac{\delta_{\gamma}^i}{\delta_{\gamma}^a} \right|. \quad (129)$$

where δ_{γ}^i is the perturbation in the photon energy density due to isocurvature perturbations and δ_{γ}^a is the perturbation due to adiabatic perturbations.

3.5 Inflation models

A detailed account on inflation model building can be found in many reviews [15, 98, 155, 9]. Here we briefly recall some of the popular models with a particular emphasis on supersymmetric inflation. First we recapitulate some aspects of non-supersymmetric models.

3.5.1 Non-supersymmetric inflation

The very first attempt to build an inflation model was made in [154], where one loop quantum correction to the energy momentum tensor due to the space-time curvature were taken into account, resulting in terms of higher order in curvature invariants. Such corrections to the Einstein equation admit a de Sitter solution [209], which was presented in [154, 210]. Inflation in Einstein gravity with an additional R^2 term was

considered in [211] (for a discussion of inflation in pure R^2 gravity, see [212]). Such a theory is conformally equivalent to a theory with a canonical gravity [213] with a scalar field having a potential term. A similar situation arises in theories with a variable Planck mass, i.e., in scalar tensor theories [66]. Inflation in these models has been studied extensively [214].

The simplest single field inflation model is arguably chaotic inflation [158, 215] with a generic potential

$$V = \frac{\lambda}{M_{\text{P}}^{\alpha-4}} \phi^\alpha, \quad (130)$$

where α is a positive even integer. In chaotic inflation slow roll takes place for $\phi \gg M_{\text{P}}$, and the two slow roll parameters are given by [9]

$$\epsilon \equiv \frac{\alpha^2}{2} \frac{M_{\text{P}}^2}{\phi^2}, \quad \eta = \alpha(\alpha - 1) \frac{M_{\text{P}}^2}{\phi^2}. \quad (131)$$

Inflation ends when $\epsilon \equiv 1$, or, $\phi \approx \alpha M_{\text{P}}$. The cosmological scales leave the horizon when $\phi = \sqrt{2N\alpha} M_{\text{P}}$, and the spectral indices for scalar and tensor perturbations turn out to be [9]

$$n = 1 - \frac{2 + \alpha}{2N}, \quad r = \frac{3.1\alpha}{N}. \quad (132)$$

The amplitude of the density perturbations, if normalized at the COBE scale, yields the constraint $\lambda \simeq 4 \times 10^{-14}$.

An exponential potential, such as might arise in string theories and theories with extra dimensions,

$$V(\phi) = V_0 \exp\left(-\sqrt{\frac{2}{p}} \frac{\phi}{M_{\text{P}}}\right). \quad (133)$$

would give rise to a power law $a(t) \propto t^p$ for the scale factor, so that inflation occurs when $p > 1$. Multiple exponentials with differing slopes give rise to what has been dubbed as assisted inflation [161].

3.5.2 F-term inflation

In four dimensions the $N = 1$ supersymmetric potential receives two contributions: one from the F-term, which is related to the chiral supermultiplets, and the second from the D-term, which contains the gauge interactions. For a detailed discussion of

supersymmetric inflation we refer to the review by Lyth and Riotto [155]. Here we give a brief resume of the two types of inflation.

Historically, supersymmetric inflation was first introduced to cure some of the problems associated with the fine tuning of new inflation [216], but since then utilizing supersymmetry as a tool for inflation has gained in popularity (we describe supersymmetry in Sect. 4.2, and for supergravity, see Sect. 4.5.2.). The F-term potential can be derived from the superpotential W

$$V(\phi, \phi^*) = F^{*i} F_i, \quad F_i = - \left(\frac{\partial W}{\partial \phi_i} \right)^*, \quad (134)$$

where for renormalizable interactions W has a mass dimension three.

Supersymmetry is broken whenever $|F|^2 \neq 0$. A simple working example is to consider the superpotential

$$W = \lambda S(\phi^2 - \phi_0^2) \quad (135)$$

which is invariant under a global R symmetry with the superfields S and N carrying respectively the R charges 1 and 0. The scalar components of these superfields can be written in the form

$$S = \frac{\sigma}{\sqrt{2}}, \quad \phi = \frac{\phi_1 + i\phi_2}{\sqrt{2}}, \quad (136)$$

where we have used R -transformation in order to make S real. The potential follows from Eq. (134):

$$V = \lambda^2 \phi_0^4 - \lambda^2 \phi_0^2 (\phi_1^2 - \phi_2^2) + \frac{\lambda^2}{4} (\phi_1^2 + \phi_2^2)^2 + \lambda^2 \sigma^2 (\phi_1^2 + \phi_2^2). \quad (137)$$

The supersymmetric vacuum is located at $\sigma = 0$, $\phi_1 = \phi_0$, and $\phi_2 = 0$. Note that the potential has a flat direction along σ -axis when $\sigma > \sigma_{inst} = \phi_0$. When $\sigma < \sigma_{inst}$, the mass squared of ϕ_1 becomes negative and suggests a phase transition along the ϕ_1 direction. When this happens σ , ϕ_1 , and ϕ_2 begin to oscillate around their supersymmetry preserving vacua. If $\phi_1 = \phi_2 = 0$, the height of the potential is given by $V = \lambda^2 \phi_0^4$, and as a consequence one obtains a period of inflation. This is the simplest example of a flat direction giving rise to an inflation potential, and it is known as the hybrid model, first described in a non-supersymmetric context in [215, 160] and in a supersymmetric context in [217].

In order to have a graceful exit from inflation one requires a slope for the flat direction such that σ can roll down and approach σ_{inst} . The flatness of the potential can be lifted in two ways: by radiative corrections [218], or by the low energy soft supersymmetry breaking effects.

Due to supersymmetry breaking the fermions obtain a mass of the order $(\partial^2 W / \partial \phi^2) = \lambda S$, while the two complex scalars receive a mass squared $\lambda^2 S^2 \pm \lambda^2 \phi_0^2$. The one-loop radiative correction to the potential is given by [219]

$$\delta V = \frac{1}{64\pi^2} \sum_i (-)^{f_i} M_i^4 \ln \frac{M_i^2}{M^2}, \quad (138)$$

where f_i denotes the number of fermions, M_i^2 is the fermion mass squared, and M the cut-off or the renormalization scale. The summation should be taken over all helicity states i . In the present example the effective potential along the flat direction is given by

$$V = \lambda^2 \phi_0^4 \left(1 + \frac{C\lambda^2}{8\pi^2} \ln \frac{\sigma}{\sqrt{2}M} \right), \quad (139)$$

where C is a constant essentially counting the states running in the loops. If the loop correction dominates over the tree level potential, there is a period of inflation which typically ends when

$$\sigma = \lambda \sqrt{\frac{CN}{4\pi^2}} M_P, \quad (140)$$

The COBE normalization sets the scale of inflation to

$$V^{1/4} \sim \left(\frac{50}{N} \right)^{1/4} C^{1/4} \lambda \times 10^{15} \text{ GeV} \quad (141)$$

while the spectral index is given by

$$n = 1 - \frac{1}{N} \left(1 + \frac{3C\lambda^2}{16\pi^2} \right). \quad (142)$$

Therefore, depending on the coupling constant λ and the number of e-foldings N , it is possible to have a wide range of inflation energy scales which all provide a spectral index $n \sim 0.96 - 0.98$.

Soft supersymmetry breaking contributions induce $m_\sigma \sim \mathcal{O}(\text{TeV})$. One could also imagine that the mass of σ appears dynamically if σ has couplings to bosons and fermions; these may induce a typical running mass $\propto \sigma^2 \ln(\sigma/M)$ [220, 155, 221].

3.5.3 D-term inflation

In the above discussion we have neglected the gauge contribution. The D-term $D^a = -g_a(\phi^* T^a \phi)$ gives rise to a scalar potential (see [37, 34])

$$V(\phi, \phi^*) = \frac{1}{2} \sum D^a D_a \quad (143)$$

where $(T^a)_i{}^j$ satisfy $[T^a, T^b] = i f^{abc} T^c$ (f^{abc} is the structure constant).

The simplest realization of D-term inflation reproduces the hybrid potential with three chiral superfields, S , ϕ_+ , and ϕ_- with (non-anomalous) $U(1)$ charges 0, +1, -1 [222]. The superpotential can be written as

$$W = \lambda S \phi_+ \phi_- . \quad (144)$$

The scalar potential then reads [222]

$$V = \lambda^2 |S|^2 (|\phi_+|^2 + |\phi_-|^2) + \lambda^2 |\phi_+ \phi_-|^2 + \frac{g^2}{2} (|\phi_+|^2 - |\phi_-|^2 + \xi^2)^2 , \quad (145)$$

where g is the gauge coupling and ξ is the Fayet-Iliopoulos D-term. Note that the potential allows unique supersymmetry preserving vacua with a broken gauge symmetry $S = \phi_+ = 0$, and $\phi_- = \xi$. By virtue of the coupling, when $|S| > S_{inst} = g\xi/\lambda$, the fields $\phi_+, \phi_- \rightarrow 0$, and therefore inflation occurs because of the Fayet-Iliopoulos D-term $V = g^2 \xi^4 / 2$. The slope along the inflaton direction S can be generated by the one-loop contribution and reads

$$V = \frac{g^2 \xi^4}{2} \left(1 + \frac{g^2}{16\pi^2} \ln \frac{\lambda^2 |S|^2}{M_{\text{P}}^2} \right) . \quad (146)$$

Inflation ends when slow roll condition breaks down for $S \sim (g/2\pi\sqrt{2})M_{\text{P}}$, and the predictions for the inflationary parameters are similar to the previous discussion. D-term inflation based on an anomalous $U(1)$ symmetry (which could appear in string theory [223]) is no different.

Hybrid inflation is successful but has also problems that are related to the initial conditions. In [224], it was pointed out that hybrid inflation requires an extremely homogeneous field configuration for the fields orthogonal to the inflaton. In our example the orthogonal fields to the inflaton must be set to zero with a high accuracy

over a region much larger than the initial size of the horizon. It is possible to solve this impasse by having a pre-inflationary matter dominated phase when the field orthogonal to the inflaton direction oscillates and decays into lighter degrees of freedom, gradually settling down to the bottom of its potential [224].

3.5.4 Supergravity corrections

When the field values are close to the Planck scale, supergravity (SUGRA) effects become important and may ruin the flatness of the inflaton potential. The soft breaking mass of the scalar fields are typically [225, 217, 226, 45, 46]

$$m_{soft}^2 \sim \frac{V}{3M_{\text{P}}^2} \sim \mathcal{O}(1)H^2. \quad (147)$$

Once the inflaton gains a mass $\sim H$, the field simply rolls down to the minimum of the potential and inflation stops. Indeed, in SUGRA the slow roll parameter

$$|\eta| \equiv M_{\text{P}}^2 \frac{V''}{V} \sim \frac{m_{SUGRA}^2}{H^2} \sim \mathcal{O}(1), \quad (148)$$

where $m_{SUGRA}^2 \approx m_{SUSY}^2 + (V_{SUSY}/3M_{\text{P}}^2) \sim m_{SUSY}^2 + \mathcal{O}(1)H^2$. Note that the latter contribution dominates in an expanding Universe and violates the slow roll condition. For field values smaller than Planck scale it is always possible to obtain $\epsilon \ll 1$, but in supergravity η can never be made less than one for a single chiral field with a minimal kinetic term. This is known as the η problem in SUGRA models of inflation [217].

When there are more than one chiral superfields, it might be possible to cancel the dominant $\mathcal{O}(1)H$ correction to the inflaton mass by choosing an appropriate Kähler term [226, 217] (see also discussion in [155]). In hybrid inflation models derived from an F-term the dominant $\mathcal{O}(1)H$ correction in the mass term can be canceled if $|N| = 0$ exactly, which however seems to lead to an initial condition problem, as discussed above. The fact that the superpotential is linear in S in Eqs. (135,144) guarantees the cancellation of the dominant contribution in the mass term for a minimal Kähler term $\sim |S|^2$. For non-minimal Kähler potential such as $K = |S|^2 + \beta|S|^4/M_{\text{P}}^2 + \dots$, one obtains $(\partial^2 K / \partial S \partial S^*)^{-1} \sim 1 - 4\beta|S|^2/M_{\text{P}}^2$. These contributions again lead to a problematic $\beta \times \mathcal{O}(1)H$ contribution to the inflaton mass unless the value of the unknown constant β is suppressed.

In [222], it was shown that the η problem does not appear for D-term inflation even for the minimal Kähler potential because the main contribution to the inflation potential does not come from the vev of the inflaton field alone, but from the Fayet-Iliopoulos term which belongs to the D-sector of the potential. Based on this fact many D-term inflation models have been written down [227, 228]. Therefore, hybrid inflation, whether realized as an effective potential coming from F-sector or from D-sector, appears to be among the most promising models for supersymmetric inflation.

3.6 Reheating of the Universe

3.6.1 Perturbative inflaton decay

Traditionally reheating has been assumed to be a consequence of the perturbative decay of the inflaton [229, 198, 22]. After the end of inflation, when $H \leq m_\phi$, the inflaton field oscillates about the minimum of the potential. Averaging over one oscillation results in [230] pressureless equation of state where $\langle p \rangle = \langle \dot{\phi}^2/2 - V(\phi) \rangle$ vanishes², so that the energy density redshifts as during matter domination with $\rho_\phi = \rho_i(a_i/a)^3$ (subscript i denotes the quantities right after the end of inflation). If Γ_ϕ represents the decay width of the inflaton to a pair of fermions, then the inflaton decays when $H(a) = \sqrt{(1/3M_{\text{P}}^2)\rho_i(a_i/a)^{3/2}} \approx \Gamma_\phi$. When the inflaton decays, it releases its energy into the thermal bath of relativistic particles whose energy density is determined by the reheat temperature T_{rh} , given by

$$T_{rh} = \left(\frac{90}{\pi^2 g_*} \right)^{1/4} \sqrt{\Gamma_\phi M_{\text{P}}} = 0.3 \left(\frac{200}{g_*} \right)^{1/4} \sqrt{\Gamma_\phi M_{\text{P}}}. \quad (149)$$

However the inflaton might not decay instantaneously. In such a case there might already exist a thermal plasma of some relativistic species at a temperature higher than the reheat temperature already before the end of reheating [22]. If the inflaton decays with a rate Γ_ϕ , then the instantaneous plasma temperature is found to be [22]

$$T_{inst} \sim \left(g_*^{-1/2} H \Gamma_\phi M_{\text{P}}^2 \right)^{1/4}, \quad (150)$$

²This will be discussed in a more detail in Sect. 5.8.

where g_* denotes the effective relativistic degrees of freedom in the plasma. The temperature reaches its maximum T_{max} soon after the inflaton field starts oscillating around the minimum. Once the maximum temperature is reached, then $\rho_\psi \sim a^{-3/2}$, and $T \sim a^{-3/8}$ until reheating and thermalization is completely over [231, 232, 233, 234, 235].

The process of thermalization has two aspects; achieving kinetic equilibrium, and achieving chemical equilibrium. Kinetic equilibrium can be reached by $2 \rightarrow 2$ scattering and annihilation. For chemical equilibrium one requires particle number changing interactions such as $2 \rightarrow 3$ processes. In [232], soft processes which allow for small momentum transfer with a larger cross-section have been advocated for chemical equilibration, while in [231], hard processes have been invoked. Therefore, depending on the interactions, thermalization time scale could be short, such as in the case of soft scattering processes, or, it could be long compared to the Hubble time if only hard processes are operative. Recently it has been argued [233, 234, 235], that thermalization time scale can be as long as the time it takes for the inflaton decay products with typical energies $\mathcal{O}(m_\phi)$ to lose the energy $\sim (m_\phi - T_{rh})$. The main conclusion is that inelastic scattering interactions $2 \rightarrow 3$ can thermalize the Universe faster compared to elastic interaction $2 \rightarrow 2$. Inelastic interactions can achieve the kinetic and chemical equilibrium both, and therefore, Γ_{inel}^{-1} could be considered as the true thermalization time scale. In [235], the authors have studied thermalization in QCD based approach with emphasis upon late thermalization and hadronization.

3.6.2 Non-perturbative inflaton decay

Much effort has lately been devoted to non-perturbative effects which are essentially non-thermal. These may lead to a rapid transfer of the inflaton energy to other degrees of freedom by the process known as preheating. The requirement is that the inflaton quanta couple to other (essentially massless) fields χ through e.g. terms like $\phi^2 \chi^2$. The quantum modes of χ may then be excited during the inflaton oscillations via a parametric resonance. Preheating has been treated both analytically [236, 237, 238, 239, 240, 241, 242, 243], and on lattice [244].

Like bosons, fermions can also be excited through preheating [245, 246, 136, 247]. In fact it has been argued that fermionic preheating is perhaps more effective than bosonic preheating [246, 136, 248]. However note that supersymmetry is effectively broken during the inflaton oscillations [247, 249, 250]. As a consequence, one naturally expects corrections to the inflaton potential during the oscillations [251]. Therefore in most supersymmetric models of inflation preheating might not turn out to be very relevant.

3.6.3 Gravitino and inflatino problems

The reheat temperature should certainly be above the BBN temperature $T \geq \mathcal{O}(1)$ MeV, but there also exists an upper bound from gravitino overproduction. In supergravity the superpartner of the graviton is a spin-3/2 gravitino, which gets a mass from the super-Higgs mechanism [252] when supersymmetry is spontaneously broken. Typically supergravity is broken in a hidden sector by some non-perturbative dynamics. Supersymmetry breaking is then mediated via gravitational (or possibly other) couplings to the observable sector in such a way that sfermions and gauginos get masses of order electroweak scale [253, 34]. In addition, the gravitino also gets a mass which in the simplest gravity mediated models is of order 1 TeV [254] (see Sect. 4.4).

If the gravitino is not the lightest supersymmetric particle (LSP), it will decay. Gravitino has two helicity states $\pm 3/2$ and $\pm 1/2$. The latter one is mainly the goldstino mode which is eaten by the super-Higgs mechanism. The goldstino coupling strength is inversely proportional to the momentum, so that at low energies the gravitino coupling is mainly dictated by the goldstino mode [255]. At temperatures much above the sparticle masses, it is the massless $\pm 3/2$ mode that governs the gravitino interactions. The helicity $\pm 3/2$ mode can decay into gauge bosons and gauginos through a dimension 5-operator with a lifetime

$$\tau_{3/2 \rightarrow A_{\mu\lambda}} \approx \frac{4M_{\text{P}}^2}{m_{3/2}^3}. \quad (151)$$

Typically $\tau \sim 10^2 - 10^5$ s for a gravitino mass in the range $10 \text{ TeV} \leq m_{3/2} \leq 100 \text{ GeV}$.

Although the gravitino interactions with matter are suppressed by the Planck mass, they can be generated in great abundances very close to the Planck scale [256]. In-

flation would dilute their number density [257], but during reheating they would be regenerated through scattering of gauge and gaugino quanta, with adverse consequences [102, 258, 259, 260, 261, 262, 263, 264]. The resulting gravitino abundance has been estimated to be [102]

$$\frac{n_{3/2}}{s} \approx 2.4 \times 10^{-13} \left(\frac{T_{rh}}{10^9 \text{ GeV}} \right) \left[1 - 0.018 \ln \left(\frac{T_{rh}}{10^9 \text{ GeV}} \right) \right], \quad (152)$$

where s defines the entropy density and T_{rh} denotes the reheating temperature of the Universe. The abundance Eq. (152) could well be increased by an order of magnitude if gravitino interactions with other chiral multiplets are included [260]. In [261] it was argued that at finite temperatures gravitino overproduction could be enhanced, but the calculation was criticized in [262, 263]; for a recent discussion on this topic, see [264].

Since the gravitino is a late decaying particle, BBN yields a restriction on the reheat temperature [258, 2]. For instance, gravitino decay products can enhance the abundance of $D + {}^3\text{He}$ due to photo fission of ${}^4\text{He}$ which implies [2]

$$\begin{aligned} \frac{n_{3/2}}{s} &\leq (10^{-14} - 10^{-11}) \Rightarrow \\ T_{rh} &\leq (10^7 - 10^{10}) \text{ GeV}, \quad 100 \text{ GeV} \leq m_{3/2} \leq 10 \text{ TeV}. \end{aligned} \quad (153)$$

The constraint on the reheating temperature is [2]

$$T_{rh} \leq 2.5 \times 10^8 \left(\frac{m_{3/2}}{100 \text{ GeV}} \right)^{-1} \text{ GeV}, \quad (154)$$

for $m_{3/2} \leq 1.6 \text{ TeV}$.

In gauge mediated supersymmetry breaking scenarios, to be discussed in Sect. 4.4., the gravitino can have a very light mass $\sim 10^{-6} \text{ GeV}$ [265] and can be a hot dark matter candidate [266]. Small (or large) gravitino masses can also be obtained in SUGRA models with non-minimal Kähler terms, such as the no-scale model [267]. In anomaly mediation the gravitino mass is large with $m_{3/2} \sim m_{soft}/\alpha \gg m_{soft}$ [268]. In general, if the gravitino is not LSP and heavier than 10 TeV, it decays before nucleosynthesis and thus does not cause any cosmological problems [269].

Gravitinos could also be produced by non-perturbative processes, as was first described in [247], where the formalism for exciting the helicity $\pm 3/2$ component of the

gravitino was developed. Later the production of the helicity $\pm 1/2$ state, which for a single chiral multiplet is the superpartner of the inflaton known as inflatino, has been studied by several authors [249, 250, 248, 270, 271]. The decay channels of the inflatino have been first discussed in [272, 273]. It has been suggested [272] and also explicitly shown [271] that in realistic models with several chiral multiplets, the helicity $\pm 1/2$ gravitino production is not a problem for nucleosynthesis as long as the inflationary scale is sufficiently higher than the scale of supersymmetry breaking in the hidden sector and the two sectors are gravitationally coupled. A very late decay of inflatino could however be possible, as argued in [273, 274]. In [274], it was argued that if the inflatino and gravitino were not LSP, then late off-shell inflatino and gravitino mediated decays of heavy relics could be significant.

4 Flat directions

4.1 Degenerate vacua

At the level of renormalizable terms, supersymmetric field theories generically have infinitely degenerate vacua. This is a consequence of the supersymmetry and the gauge symmetries (and discrete symmetries such as R -parity) of the Lagrangian, which allow for certain types of interaction terms only. Therefore, in general there are a number of directions in the space of scalar fields, collectively called the moduli space, where the scalar potential is identically zero. In low energy supersymmetric theories such classical degeneracy is accidental and is protected from perturbative quantum corrections by a non-renormalization theorem [275]. In principle the degeneracies could be lifted by non-perturbative effects. However such effects are likely to be suppressed exponentially and thus unimportant because all the couplings of low energy theories are typically weak even at relatively large vevs. Therefore in the supersymmetric limit when $M_p \rightarrow \infty$, the potential for the flat direction always vanishes.

In the MSSM the moduli fields are quark, lepton and Higgs chiral fields. In string theories there are often additional moduli fields associated with the conformal field theory degrees of freedom and world sheet discrete R -symmetries [276]. The moduli space of string theory can also be lifted by a soft supersymmetry breaking masses of the order of the gravitino mass $m_{3/2}$. Since the moduli interactions with others fields are usually Planck mass suppressed, the string moduli are also a cause for worry because they may decay after nucleosynthesis. This problem has been dubbed as the moduli problem [225, 277]. However, the MSSM flat directions are made up of condensates of squarks, Higgses, and sleptons, and can evaporate much before nucleosynthesis.

However there is an effective potential for the flat direction condensate fields which arises as a result of supersymmetry breaking terms and higher dimensional operators in the superpotential. In this sense the MSSM flat directions are only approximately flat at vevs larger than the supersymmetry breaking scale.

4.2 MSSM and its potential

Let us remind the reader that the matter fields of MSSM are chiral superfields $\Phi = \phi + \sqrt{2}\theta\bar{\psi} + \theta\bar{\theta}F$, which describe a scalar ϕ , a fermion ψ and a scalar auxiliary field F . In addition to the usual quark and lepton superfields, MSSM has two Higgs fields, H_u and H_d . Two Higgses are needed because $H^\dagger$, which in the Standard Model gives masses to the u -quarks, is forbidden in the superpotential.

The superpotential for the MSSM is given by [34]

$$W_{MSSM} = \lambda_u Q H_u \bar{u} + \lambda_d Q H_d \bar{d} + \lambda_e L H_d \bar{e} + \mu H_u H_d, \quad (155)$$

where $H_u, H_d, Q, L, \bar{u}, \bar{d}, \bar{e}$ in Eq. (155) are chiral superfields, and the dimensionless Yukawa couplings $\lambda_u, \lambda_d, \lambda_e$ are 3×3 matrices in the family space. We have suppressed the gauge and family indices. Unbarred fields are $SU(2)$ doublets, barred fields $SU(2)$ singlets. The last term is the μ term, which is a supersymmetric version of the SM Higgs boson mass. Terms proportional to $H_u^* H_u$ or $H_d^* H_d$ are forbidden in the superpotential, since W_{MSSM} must be analytic in the chiral fields. H_u and H_d are required not only because they give masses to all the quarks and leptons, but also for the cancellation of gauge anomalies. The Yukawa matrices determine the masses and CKM mixing angles of the ordinary quarks and leptons through the neutral components of $H_u = (H_u^+, H_u^0)$ and $H_d = (H_d^0, H_d^-)$. Since the top quark, bottom quark and tau lepton are the heaviest fermions in the SM, we assume that only the $(3, 3)$ element of the matrices $\lambda_u, \lambda_d, \lambda_e$ are important. In this limit only the third family and the Higgs fields contribute to the MSSM superpotential.

The SUSY scalar potential V is the sum of the F- and D-terms and reads

$$V = \sum_i |F_i|^2 + \frac{1}{2} \sum_a g_a^2 D^a D^a \quad (156)$$

where

$$F_i \equiv \frac{\partial W_{MSSM}}{\partial \phi_i}, \quad D^a = \phi^\dagger T^a \phi. \quad (157)$$

Here we have assumed that ϕ_i transforms under a gauge group G with the generators of the Lie algebra given by T^a .

The μ term provides masses to the Higgsinos

$$\mathcal{L} \supset -\mu(\tilde{H}_u^+ \tilde{H}_d^- - \tilde{H}_u^0 \tilde{H}_d^0) + c.c., \quad (158)$$

and contributes to the Higgs $(mass)^2$ terms in the scalar potential through

$$-\mathcal{L} \supset V \supset |\mu|^2(|H_u^0|^2 + |H_u^+|^2 + |H_d^0|^2 + |H_d^-|^2). \quad (159)$$

Note that Eq. (159) is positive definite. Therefore, it cannot lead to electroweak symmetry breaking without including supersymmetry breaking $(mass)^2$ soft terms for the Higgs fields, which can be negative. Hence, $|\mu|^2$ should almost cancel the negative soft $(mass)^2$ term in order to allow for a Higgs vev of order ~ 174 GeV. That the two different sources of masses should be precisely of same order is a puzzle for which many solutions has been suggested [278, 279, 280, 281].

Note also that Eq. (155) is the minimal superpotential because we have not included terms which are gauge invariant and analytic in the chiral superfields but which violate either baryon number B or lepton number L . The most general gauge invariant and renormalizable superpotential would not only include Eq. (155), but also the terms

$$W_{\Delta L=1} = \frac{1}{2}\lambda^{ijk}L_iL_j\bar{e}_k + \lambda'^{ijk}L_iQ_j\bar{d}_k + \mu'^iL_iH_\mu, \quad (160)$$

$$W_{\Delta B=1} = \frac{1}{2}\lambda''^{ijk}\bar{u}_i\bar{d}_j\bar{d}_k, \quad (161)$$

where $i = 1, 2, 3$ represents the family indices. The chiral supermultiplets carry baryon number assignments $B = +1/3$ for Q_i , $B = -1/3$ for $\bar{u}_i, \bar{d}_i$, and $B = 0$ for all others. The total lepton number assignments are $L = +1$ for L_i , $L = -1$ for $\bar{e}_i$, and $L = 0$ for all the others. The terms in Eq. (160) violate lepton number by one unit, while those in Eq. (161) violate baryon number by one unit.

Unless λ' and λ'' terms are very much suppressed, one would obtain rapid proton decay which violates both B and L by one unit. Many other processes also give rise to violation in baryon and lepton number (for a review, see [74]). Therefore, there must be a symmetry forbidding the terms in Eqs. (160,161), while allowing for the terms in Eq. (155). The symmetry is known as R -parity [255], which is a discrete parity defined for each particle as

$$P_R = (-1)^{3(B-L)+2s} \quad (162)$$

with $P_R = +1$ for the SM particles and the Higgs bosons, while $P_R = -1$ for all the sleptons, squarks, gauginos, and Higgsinos. Here s is spin of the particle. Without the product $(-1)^{2s}$, the expression is known as matter parity [282], and denoted by $P_M = (-1)^{3(B-L)}$. The quantity $(-1)^{2s}$ is equal to 1 whenever conservation of angular momentum holds at a given vertex. In this case matter parity and R -parity are equivalent. If R -parity is conserved then there will be no mixing between the sparticles and the ones which have $P_R = +1$. This completely forbids potentially dangerous terms in Eqs. (160,161).

Matter parity is actually a discrete subgroup of the continuous $U(1)_{B-L}$ group. Therefore, if a gauged $U(1)_{B-L}$ is broken by scalar vevs which carry even integer values of $3(B-L)$, then P_M survives as an exactly conserved discrete remnant [283]. Besides forbidding B and L violation from the renormalizable interactions, R -parity has interesting phenomenological and cosmological consequences. The lightest sparticle with $P_R = -1$, the LSP, must be absolutely stable. If electrically neutral, the LSP is a natural candidate for non-baryonic dark matter [284, 285]. It may be possible to produce LSPs in a next generation collider experiments.

4.2.1 F-and D-renormalizable flat directions of MSSM

For a general supersymmetric model with N chiral superfields X_i , it is possible to find out the directions where the potential Eq. (156) vanishes identically by solving simultaneously

$$D^a \equiv X^\dagger T^a X = 0, \quad F_{X_i} \equiv \frac{\partial W}{\partial X_i} = 0. \quad (163)$$

Field configurations obeying Eq. (163) are called respectively D-flat and F-flat.

D-flat directions are parameterized by gauge invariant monomials of the chiral superfields. A powerful tool for finding the flat directions has been developed in [286, 287, 288, 46, 289, 43], where the correspondence between gauge invariance and flat directions has been employed. The configuration space of the scalar fields of the MSSM contains 49 complex dimensions (18 for Q_i , 9 each for $\bar{u}_i$ and $\bar{d}_i$, 6 for L_i , 3 for $\bar{e}_i$, and 2 each for H_u and H_d), out of which there are 12 real D-term constraints (8 for $SU(3)_C$, 3 for $SU(2)_L$, and 1 for $U(1)_Y$), which leaves a total of 37 complex dimensions

[46, 43]. The trick is to construct gauge invariant monomials forming $SU(3)_C$ singlets and then using them as building blocks to generate $SU(3)_C \times SU(2)_L$, and subsequently the whole $SU(3)_C \times SU(2)_L \times U(1)_Y$ invariant polynomials [46, 43]. However these invariant monomials give only the D-flat directions. For F-flat directions, one must solve explicitly the constraint equations $F_{X_i} = 0$.

A single flat direction necessarily carries a global $U(1)$ quantum number, which corresponds to an invariance of the effective Lagrangian for the order parameter ϕ under phase rotation $\phi \rightarrow e^{i\theta}\phi$. In the MSSM the global $U(1)$ symmetry is $B - L$. For example, the LH_u -direction (see below) has $B - L = -1$.

A flat direction can be represented by a composite gauge invariant operator, X_m , formed from the product of k chiral superfields Φ_i making up the flat direction: $X_m = \Phi_1 \Phi_2 \cdots \Phi_m$. The scalar component of the superfield X_m is related to the order parameter ϕ through $X_m = c\phi^m$.

4.2.2 An example of F-and D-flat direction

The flat directions in the MSSM are tabulated in Table 1. An example of a D-and F-flat direction is provided by

$$H_u = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ \phi \end{pmatrix}, \quad L = \frac{1}{\sqrt{2}} \begin{pmatrix} \phi \\ 0 \end{pmatrix}, \quad (164)$$

where ϕ is a complex field parameterizing the flat direction, or the order parameter, or the AD field. All the other fields are set to zero. In terms of the composite gauge invariant operators, we would write $X_m = LH_u$ ($m = 2$).

From Eq. (164) one clearly obtains $F_{H_u}^* = \lambda_u Q \bar{u} + \mu H_d = F_L^* = \lambda_d H_d \bar{e} \equiv 0$ for all ϕ . However there exists a non-zero F-component given by $F_{H_d}^* = \mu H_u$. Since μ can not be much larger than the electroweak scale $M_W \sim \mathcal{O}(1)$ TeV, this contribution is of the same order as the soft supersymmetry breaking masses, which are going to lift the degeneracy. Therefore, following [46], one may nevertheless consider LH_u to correspond to a F-flat direction.

The relevant D-terms read

$$D_{SU(2)}^a = H_u^\dagger \tau_3 H_u + L^\dagger \tau_3 L = \frac{1}{2} |\phi|^2 - \frac{1}{2} |\phi|^2 \equiv 0. \quad (165)$$

	$B - L$		$B - L$
$H_u H_d$	0	$L H_u$	-1
$\bar{u} \bar{d} \bar{d}$	-1	$Q L \bar{d}$	-1
$LL\bar{e}$	-1	$QQ\bar{u}\bar{d}$	0
$QQQL$	0	$QL\bar{u}\bar{e}$	0
$\bar{u}\bar{u}\bar{d}\bar{e}$	0	$QQQQ\bar{u}$	1
$QQ\bar{u}\bar{u}\bar{e}$	1	$LL\bar{d}\bar{d}\bar{d}$	-3
$\bar{u}\bar{u}\bar{u}\bar{e}\bar{e}$	1	$QLQL\bar{d}\bar{d}$	-2
$QQLL\bar{d}\bar{d}$	-2	$\bar{u}\bar{u}\bar{d}\bar{d}\bar{d}\bar{d}$	-2
$QQQQ\bar{d}LL$	-1	$QLQLQL\bar{e}$	-1
$QL\bar{u}QQ\bar{d}\bar{d}$	-1	$\bar{u}\bar{u}\bar{u}\bar{d}\bar{d}\bar{d}\bar{e}$	-1

Table 1: **Renormalizable F and D flat directions in the MSSM**

Therefore the LH_u direction is also D-flat.

The only other direction involving the Higgs fields and thus soft terms of the order of μ is $H_u H_d$. The rest are purely leptonic, such as $LL\bar{e}$, or baryonic, such as $\bar{u}\bar{d}\bar{d}$, or mixtures of leptons and baryons, such as $QL\bar{d}$. These combinations give rise to several independent flat directions that can be obtained by permuting the flavor indices. For instance, $LL\bar{e}$ contains the directions $L_1 L_2 \bar{e}_3$, $L_2 L_3 \bar{e}_1$, and $L_1 L_3 \bar{e}_2$.

Along a flat direction gauge symmetries get broken, with the gauge supermultiplets gaining mass by super-Higgs mechanism with $m_g = g\langle\phi\rangle$. Several chiral supermultiplets typically become massive by virtue of Yukawa couplings in the superpotential; for example, in the LH_u direction one finds the mass terms $W_{\text{mass}} = \lambda_u \langle\phi\rangle Q\bar{u} + \lambda_e \langle\phi\rangle H_d \bar{e}$.

Of course, there may simultaneously exist several flat directions. For the purpose of AD mechanism it is the lowest dimensional operator which determines the baryonic charge of the eventual condensate. In what follows we will therefore mostly consider a single flat direction.

4.3 Lifting the flat direction

Vacuum degeneracy along a flat direction can be broken in two ways: by supersymmetry breaking, or by higher order non-renormalizable operators appearing in the effective low energy theory. Let us first consider the latter option. Supersymmetry breaking will then be discussed in more detail in Sects. 4.4 and 4.5.

4.3.1 Lifting by non-renormalizable operators

Non-renormalizable superpotential terms in the MSSM can be viewed as effective terms that arise after one integrates out fields with very large mass scales appearing in a more fundamental (say, string) theory. Here we do not concern ourselves with the possible restrictions on the effective terms due to discrete symmetries present in the fundamental theory, but assume that all operators consistent with symmetries may arise. Thus in terms of the invariant operators X_m , one can have terms of the type [45, 46]

$$W = \frac{h}{dM^{d-3}} X_m^k = \frac{h}{dM^{d-3}} \phi^d, \quad (166)$$

where the dimensionality of the effective scalar operator $d = mk$, and h is a coupling constant which could be complex with $|h| \sim \mathcal{O}(1)$. Here M is some large mass, typically of the order of the Planck mass or the string scale (in the heterotic case $M \sim M_{GUT}$). The lowest value of k is 1 or 2, depending on whether the flat direction is even or odd under R -parity.

A second type of term lifting the flat direction would be of the form [45, 46]

$$W = \frac{h'}{M^{d-3}} \psi \phi^{d-1}, \quad (167)$$

where ψ is not contained in X_m . The superpotential term Eq. (167) spoils F-flatness through $F_\psi \neq 0$. An example is provided by the direction $\bar{u}_1 \bar{u}_2 \bar{u}_3 \bar{e}_1 \bar{e}_2$, which is lifted by the non-renormalizable term $W = (h'/M) \bar{u}_1 \bar{u}_2 \bar{d}_2 \bar{e}_1$. This superpotential term gives a non-zero contribution $F_{\bar{d}_2}^* = (h'/M) \bar{u}_1 \bar{u}_2 \bar{e}_1 \sim (h'/M) \phi^3$ along the flat direction.

Assuming minimal kinetic terms, both types discussed above in Eqs. (166,167) yield a generic non-renormalizable potential contribution that can be written as

$$V(\phi) = \frac{|\lambda|^2}{M^{2d-6}} (\phi^* \phi)^{d-1}, \quad (168)$$

where we have defined the coupling $|\lambda|^2 \equiv |h|^2 + |h'|^2$. By virtue of an accidental R -symmetry under which ϕ has a charge $R = 2/d$, the potential Eq. (168) conserves the $U(1)$ symmetry carried by the flat direction, in spite of the fact that at the superpotential level it is violated, see Eqs. (166,167). The symmetry can be violated if there are multiple flat directions, or by higher order operator contributions. However it turns out [46] that the $B - L$ violating terms are always subdominant. This is of importance for baryogenesis considerations, where the necessary $B - L$ violation should therefore arise from other sources.

The process of finding all the possible non-renormalizable superpotential contributions lifting a particular flat direction is similar to finding the D-flat directions discussed in Sect. 4.2.1. All the non-renormalizable operators can be generated from SM gauge monomials with R -parity constraint which allows only even number of odd matter parity fields $(Q, L, \bar{u}, \bar{d}, \bar{e})$ to be present in each superpotential term. At each dimension d , the various $F = 0$ constraints are separately imposed in order to construct the basis for monomials.

As an example, consider flat directions involving the Higgs fields such as $H_u H_d$ and $L H_u$ directions. Even though they are already lifted by the μ term, since μ is of the order of supersymmetry breaking scale, for cosmological purposes they can be considered flat, as was discussed in Sect. 4.2.2. At the $d = 4$ level the superpotential reads

$$W_4 \supset \frac{\lambda}{M} (H_u H_d)^2 + \frac{\lambda_{ij}}{M} (L_i H_u) (L_j H_u). \quad (169)$$

Let us assume $\lambda, \lambda_{ij} \neq 0$. Note that $F_{H_d} = 0$ constraint implies $\lambda H_u^\alpha (H_u H_d) = 0$, which acts as a basis for the monomials. An additional constraint can be obtained by contracting $F_{H_d} = 0$ by $\epsilon_{\alpha\beta} H_d^\beta$, which forms the polynomial $H_u H_d = 0$ in the same monomial basis. Similarly the constraint $F_{H_u} = 0$, along with the contraction yields $\lambda^{ij} (L_i H_u) (L_j H_u) = 0$. This implies that $L_i H_u = 0$ for all i . Therefore the two monomials $L H_u$ and $H_u H_d$ can be lifted by $d = 4$ terms in the superpotential Eq. (169).

The other renormalizable flat directions are $LLE, \bar{u}\bar{u}\bar{d}, Q\bar{d}L, QQQQL, Q\bar{u}Q\bar{d}, \bar{u}\bar{u}\bar{d}\bar{e}$ and $Q\bar{u}L\bar{e}, \bar{d}\bar{d}\bar{d}LL, \bar{u}\bar{u}\bar{u}\bar{e}\bar{e}, Q\bar{u}Q\bar{u}\bar{e}, QQQQ\bar{u}, \bar{u}\bar{u}\bar{d}QdQ\bar{d}$, and $(QQQ)_4 LLL\bar{e}$. These are

lifted primarily by the superpotential terms which involve either H_u or H_d if d is odd, or those which contain neither H_u nor H_d . The complete list of superpotential terms which lift the flat directions can be found in [43]. It was shown that all the MSSM flat directions are lifted by $d = 4, 5, 6, 7, 9$ terms in the superpotential. The unique flat directions involving (Q, u, e) is lifted by $d = 9$, (L, d) by $d = 7$, and (L, d, e) by $d = 5$. The flat directions involving (L, e) , (u, d) and (L, d, e) are all lifted by $d = 6$ terms in the superpotential, while the rest of the flat directions are lifted already by $d = 4$ superpotential terms.

4.3.2 Lifting by soft supersymmetry breaking

Vacuum degeneracy will also be lifted by supersymmetry breaking, as will be discussed in more detail in Sects. 4.4 and 4.5. It is induced by the soft terms, which in the simplest case read

$$V(\phi) = m_0^2 |\phi|^2 + \left[\frac{\lambda A \phi^d}{d M^{d-3}} + \text{h.c.} \right], \quad (170)$$

where the supersymmetry breaking mass m_0 and A are typically of the order of the gravitino mass $m_{3/2}$. An additional soft source for supersymmetry breaking are the gaugino masses m_g . The A -term in Eq. (170) violates the $U(1)$ carried by the flat direction and thus provides the necessary source for $B-L$ violation in AD baryogenesis. In general, the coupling λ is complex and has an associated phase θ_λ . Writing $\phi = |\phi| \exp(i\theta)$, one obtains a potential proportional to $\cos(\theta_\lambda + n\theta)$ in the angular direction. This has n discrete minima for the phase of ϕ , at each of which $U(1)$ is broken.

4.4 Supersymmetry breaking in the MSSM

In the MSSM there are several proposals for supersymmetry breaking, which we shall discuss below. However most of the time it is not important to know the exact mechanism of low energy supersymmetry breaking. This ignorance of the origin of supersymmetry breaking can always be hidden by simply writing down explicitly the soft breaking terms with arbitrary couplings.

4.4.1 Soft supersymmetry breaking Lagrangian

The most general soft supersymmetry breaking terms in the MSSM Lagrangian can be written as (see e.g. [35])

$$\mathcal{L}_{soft} = -\frac{1}{2}(M_\lambda \lambda^a \lambda^a + \text{c.c.}) - (m^2)_j^i \phi_j^* \phi_i - \left(\frac{1}{2} b_{ij} \phi_i \phi_j + \frac{1}{6} a^{ijk} \phi_i \phi_j \phi_k + \text{c.c.} \right), \quad (171)$$

where M_λ is the common gaugino mass $(m^2)_i^j$ are 3×3 matrices determining the masses for squarks and sleptons, denoted as $m_Q^2, m_u^2, m_d^2, m_L^2, m_e^2$; b_{ij} is the mass term for the combination $H_u H_d$; and finally, a^{ijk} are complex 3×3 matrices in the family space which yield the A -terms a_u, a_d, a_e . There are a total of 105 new entries in the MSSM Lagrangian which have no counterpart in the SM. However the arbitrariness in the parameters can be partly removed by the experimental constraints on flavor changing neutral currents (FCNC) and CP violation [290]. In order to avoid FCNC and excessive CP violation, the squark and slepton (mass)² matrices are often taken to be flavor blind, so that the squark and slepton mixing angles can be rotated away. Similarly, one may assume that the ϕ^3 couplings are proportional to the Yukawa coupling matrix, so that $a_u = A_{u0} \lambda_u$, $a_d = A_{d0} \lambda_d$, and $a_e = A_{e0} \lambda_e$. Large CP violating effects can be avoided if the soft parameters do not involve new CP phases in addition to the SM CKM phases. One can also fix μ parameter and b to be real by an appropriate phase rotation of H_u and H_d .

There are a number of possibilities for the origin of supersymmetry breaking. Fayet-Iliopoulos mechanism [291] provides supersymmetry breaking by virtue of a non-zero D-term but requires a $U(1)$ symmetry. However, this mechanism does not work in the MSSM because some of the squarks and sleptons will get non-zero vevs which may break color, electromagnetism, and/or lepton number without breaking supersymmetry. Therefore the contribution from the Fayet-Iliopoulos term should be negligible at low scales.

There are models of supersymmetry breaking by F-terms, known as O’Raifeartaigh models [292], where the idea is to pick a set of chiral supermultiplets $\Phi_i \supset (\phi_i, \psi_i F_i)$ and a superpotential W in such a way that $F_i = -\delta W / \delta \phi_i^* = 0$ have no simultaneous solution. The model requires a linear gauge singlet superfield in the superpotential.

Such singlet chiral supermultiplet is not present in the MSSM. The scale of supersymmetry breaking has to be set by hand.

The only mechanism of supersymmetry breaking where the breaking scale is not introduced either at the level of superpotential or in the gauge sector is through dynamical supersymmetry breaking [293, 294]. In these models a small supersymmetry breaking scale arises by dimensional transmutation. It is customary to treat the supersymmetry breaking sector as a hidden sector which has no direct couplings to the visible sector represented by the chiral supermultiplets of the MSSM. The only allowed interactions are those which mediate the supersymmetry breaking in the hidden sector to the visible sector.

The main contenders are gravity mediated supersymmetry breaking, which is associated with new physics which includes gravity at the string scale or at the Planck scale [253, 34], and gauge mediated supersymmetry breaking, which is transmitted to the visible sector by the ordinary electroweak and QCD gauge interactions [295, 265, 296, 297]. There are other variants of supersymmetry breaking based upon ideas on gravity and gauge mediation with some extensions, such as dynamical supersymmetry breaking (see [298], and references therein), and anomaly mediation (see [268, 299], and references therein), which we do not consider here.

4.4.2 Gravity mediated supersymmetry breaking

Let us assume that supersymmetry is broken by the vev $\langle F \rangle \neq 0$ and is communicated to the MSSM by gravity. On dimensional grounds, the soft terms in the visible sector should then be of the order [34]

$$m_{soft} \sim \frac{\langle F \rangle}{M_P}. \quad (172)$$

Note that $m_{soft} \rightarrow 0$ as $M_P \rightarrow \infty$. In order to obtain a phenomenologically acceptable soft supersymmetry mass $m_{soft} \sim \mathcal{O}(100)$ GeV, one therefore requires the scale of supersymmetry breaking in the hidden sector to be $\sqrt{\langle F \rangle} \sim 10^{10} - 10^{11}$ GeV.

Another possibility is that the supersymmetry is broken via gaugino condensate $\langle 0 | \lambda^a \lambda^b | 0 \rangle = \delta^{ab} \Lambda^3 \neq 0$, where Λ is the condensation scale [300, 34]. If the composite

field $\lambda^a \lambda^b$ belongs to the $\langle F \rangle \sim \Lambda^3/M_{\text{P}}$ -term, then again on dimensional grounds one would expect the soft supersymmetry mass contribution to be [34]

$$m_{\text{soft}} \sim \frac{\Lambda^3}{M_{\text{P}}^2}. \quad (173)$$

In this case the nature of supersymmetry breaking is dynamical and the scale is given by $\Lambda \sim 10^{13}$ GeV.

The supergravity Lagrangian must contain the non-renormalizable terms which communicate between the hidden and the observable sectors. For the cases where the kinetic terms for the chiral and gauge fields are minimal, one obtains the following soft terms [34]

$$m_{1/2} \sim \frac{\langle F \rangle}{M_{\text{P}}}, \quad m_0^2 \sim \frac{|\langle F \rangle|^2}{M_{\text{P}}^2}, \quad A_0 \sim \frac{\langle F \rangle}{M_{\text{P}}}, \quad B_0 \sim \frac{\langle F \rangle}{M_{\text{P}}}. \quad (174)$$

The gauginos get a common mass $M_1 = M_2 = M_3 = m_{1/2}$, the squark and slepton masses are $m_Q^2 = m_{\bar{Q}}^2 = m_d^2 = m_L^2 = m_{\bar{e}}^2 = m_0^2$, and for the Higgses $m_{H_u}^2 = m_{H_d}^2 = m_0^2$. The A -terms are proportional to the Yukawa couplings while $b = B_0 \mu$.

Some particular models of gravity mediated supersymmetry breaking give more detailed estimates of the soft supersymmetry terms. They include: *Dilaton dominated* models [301], which arise in a particular limit of superstring theories, which have $m_0^2 = m_{3/2}^2$, and $m_{1/2} = -A_0 = \sqrt{3}m_{3/2}$; *Polonyi* models [302], where $m_0^2 = m_{3/2}^2$, $A_0 = (3 - \sqrt{3})m_{3/2}$, and $m_{1/2} = \mathcal{O}(m_{3/2})$; and *No-scale* models [303], which also arise in the low energy limit of superstrings and in which the gravitino mass is undetermined at the tree level while the at the string scale $m_{1/2} \gg m_0, A_0, m_{3/2}$.

The predictions for the mass spectrum and other observable can be found renormalization group (RG) equations; these will be described in connection with the dynamical evolution of the AD field. Therefore, a generic flat direction in gravity mediated supersymmetry breaking has two important components: the soft supersymmetry breaking terms, and the RG induced logarithmic dependence of the vev.

4.4.3 Gauge mediated supersymmetry breaking

In gauge mediated supersymmetry breaking one employs a heavy messenger sector which couples directly to the supersymmetry breaking sector but indirectly to the

observable sector via standard model gauge interactions only [295, 304]. As a result the soft terms in the MSSM arise through ordinary gauge interactions. There will still be gravitational communication, but it is a weak effect.

The simplest example is a messenger sector with a pair of $SU(2)$ doublet chiral fields $l, \bar{l}$ and a pair of $SU(3)$ triplet fields $q, \bar{q}$, which couple to a singlet field z with Yukawa couplings λ_2, λ_3 , respectively. The superpotential is given by

$$W_{mess} = \lambda_2 z l \bar{l} + \lambda_3 z q \bar{q}. \quad (175)$$

The singlet acquires a non-zero vev and a non-zero F-term $\langle F_z \rangle$. This can be accomplished either substituting z into an O’Raifeartaigh type model [295, 304], or by a dynamical mechanism [265, 296, 297]. One may parameterize supersymmetry breaking in a superpotential W_{break} by $\langle \partial W_{break} / \partial z \rangle = -\langle F_z^* \rangle$. As a consequence, the messenger fermions acquire masses

$$\mathcal{L} = -(\lambda_2 \langle z \rangle \psi_l \psi_{\bar{l}} + \lambda_3 \langle z \rangle \psi_q \psi_{\bar{q}} + \text{c.c.}), \quad (176)$$

while the scalar messenger partners have a scalar potential given by

$$\begin{aligned} V = & |\lambda_2 \langle z \rangle|^2 (|l|^2 + |\bar{l}|^2) + |\lambda_3 \langle z \rangle|^2 (|q|^2 + |\bar{q}|^2) - (\lambda_2 \langle F_z \rangle l \bar{l} + \lambda_3 \langle F_z \rangle q \bar{q} + \text{c.c.}) \\ & + \text{quartic terms}, \end{aligned} \quad (177)$$

where we have used $\langle \partial W_{mess} / \partial z \rangle = 0$, and we have replaced z and F_z by their vevs. It is easy to read off the eigenvalues of the squared scalar masses and the fermionic and bosonic spectrum of the messenger sector; for $(l, \bar{l})$, $m_{fermions}^2 = |\lambda_2 \langle z \rangle|^2$, and $m_{scalars}^2 = |\lambda_2 \langle z \rangle|^2 \pm |\lambda_2 \langle F_z \rangle|$; for $(q, \bar{q})$, $m_{fermions}^2 = |\lambda_3 \langle z \rangle|^2$, and $m_{scalars}^2 = |\lambda_3 \langle z \rangle|^2 \pm |\lambda_3 \langle F_z \rangle|$.

Supersymmetry breaking is then mediated to the observable fields by one-loop corrections, which generate masses for the MSSM gauginos [265]. The $q, \bar{q}$ messenger loop diagrams provide masses to the gluino and the bino, while $l, \bar{l}$ messenger loop diagrams provide masses to the wino and the bino, i.e., $M_{a=1,2,3} = (\alpha_a / 4\pi) \Lambda$, where $\Lambda = \langle F_z \rangle / \langle z \rangle$.

For squarks and sleptons the leading term comes from two-loop diagrams, e.g. $m_\phi^2 \propto \alpha^2$. The A -terms get negligible contribution at two-loop order compared to

the gaugino masses, they come with an extra suppression of $\alpha/4\pi$ compared with the gaugino mass, therefore $a_u = a_d = a_e = 0$ is a good approximation. The Yukawa couplings at the electroweak scale are generated by evolving the RG equations.

One can estimate [265] the soft supersymmetry breaking masses to be of order

$$m_{soft} \sim \frac{\alpha_a}{4\pi} \frac{\langle F \rangle}{M_s}. \quad (178)$$

If $M_s \sim \langle z \rangle$ and $\sqrt{\langle F \rangle}$ are comparable mass scales, then the supersymmetry breaking can take place at about $\sqrt{\langle F \rangle} \sim 10^4 - 10^6$ GeV.

4.5 Supersymmetry breaking in the early Universe

Non-zero inflationary potential gives rise to supersymmetry breaking, the scale of which is given by the Hubble parameter. At early times this breaking is dominant over breaking from the hidden sector. After the end of inflation, in most models the inflaton oscillates and its finite energy density still dominates and breaks supersymmetry in the visible sector. Supersymmetry is broken also by quantum mechanical effects but these are negligible compared to the classical supersymmetry breaking from the non-zero energy density of the Universe.

4.5.1 Inflaton-induced terms

The early Universe supersymmetry breaking can be transmitted to the MSSM flat directions either by renormalizable or non-renormalizable interactions [46]. However at least for a single flat direction, renormalizable interactions do not lift the MSSM flat directions. In contrast, the effective potential generated by non-renormalizable interactions can induce a mass for the flat direction which is independent of the field values as long as they are below the Planck scale.

At tree level $N = 1$ SUGRA potential in four dimensions is given by the sum of F and D-terms [37]

$$V = e^{K/M_{\text{P}}^2} \left[\left(K^{-1} \right)_i^j F_i F^j - 3 \frac{|W|^2}{M_{\text{P}}^2} \right] + \frac{g^2}{2} \text{Re} f_{ab}^{-1} \hat{D}^a \hat{D}^b, \quad (179)$$

where

$$F^i = W^i + K^i \frac{W}{M_{\text{P}}^2}, \quad \hat{D}^a = -K^i (T^a)_i^j \phi_j + \xi^a. \quad (180)$$

where we have added the Fayet-Iliopoulos contribution ξ^a to the D-term. Here K is the Kähler potential, which is a function of the fields ϕ_i , and $K^i \equiv \partial K / \partial \phi_i$, and $\text{Re} f_{ab}^{-1}$ is the inverse of the real part of the gauge kinetic function matrix.

A particular class of non-renormalizable interaction terms induced by the inflaton arise if the Kähler potential has a form [305, 46, 306]

$$K = \int d^4\theta \frac{1}{M_{\text{P}}^2} (I^\dagger I) (\phi^\dagger \phi), \quad (181)$$

where I is the inflaton whose energy density $\rho \approx \langle \int d^4\theta I^\dagger I \rangle$ dominates during inflation, and ϕ is the flat direction. The interaction Eq. (181) will generate an effective mass term in the Lagrangian in the global supersymmetric limit, given by

$$\mathcal{L} = \frac{\rho_I}{M_{\text{P}}^2} \phi^\dagger \phi = 3H_I^2 \phi^\dagger \phi, \quad (182)$$

where H_I is the Hubble parameter during inflation.

4.5.2 Supergravity corrections

In addition, there are also inflaton-induced supergravity corrections to the flat direction. By inspecting the supergravity potential, one finds the following terms

$$\begin{aligned} & \left(e^{K(\phi^\dagger \phi)/M_{\text{P}}^2} V(I) \right), \quad \left(K_\phi K^{\phi\bar{\phi}} K_{\bar{\phi}} \frac{|W(I)|^2}{M_{\text{P}}^4} \right), \\ & \text{and} \quad \left(K_\phi K^{\phi\bar{I}} D_I \frac{W^*(I)W(I)}{M_{\text{P}}^2} + \text{h.c.} \right). \end{aligned} \quad (183)$$

Above $D_I \equiv \partial/\partial I + K_I W/M_{\text{P}}^2$. All these terms provide a general contribution to the flat direction potential which is of the form[46]

$$V(\phi) = H^2 M_{\text{P}}^2 f\left(\frac{\phi}{M_{\text{P}}}\right), \quad (184)$$

where f is some function. Note that this contribution exists also when the flat direction is lifted by non-renormalizable superpotential terms.

For a minimal choice of flat direction Kähler potential $K(\phi^\dagger, \phi) = \phi^\dagger \phi$, during inflation the effective mass for the flat direction is found to be [46]

$$m_\phi^2 = \left(2 + \frac{F_I^* F_I}{V(I)}\right) H^2. \quad (185)$$

Here it has been assumed that the main contribution to the inflaton potential comes from the F-term. If there were D-term contributions $V_D(I)$ to the inflationary potential, then a correction of order $V_F(I)/(V_F(I) + V_D(I))$ must be taken into account. In purely D-term inflation there is no Hubble induced mass correction to the flat direction during inflation because $F_I = 0$. However, when D-term inflation ends, the energy density stored in the D-term is converted to an F-term and to kinetic energy of the inflaton. Thus again a mass term $m_\phi^2 = \pm \mathcal{O}(1)H^2$ appears naturally, however the overall sign is undetermined [228].

There are additional inflationary contributions to the potential if the flat direction is lifted by the non-renormalizable operators discussed earlier in this Section. These new terms come explicitly from the superpotential part of the flat direction

$$\begin{aligned} & \left(W_\phi K^{\phi\bar{\phi}} K_{\bar{\phi}} \frac{W^*(I)}{M_{\text{P}}^2} + \text{h.c.} \right), \quad \left(W_\phi K^{\phi\bar{I}} D_{\bar{I}} W^*(I) + \text{h.c.} \right), \\ & \text{and} \quad \left(\frac{1}{M_{\text{P}}^2} K_I K^{I\bar{I}} K_{\bar{I}} - 3 \right) \left(\frac{W(\phi)^* W(I)}{M_{\text{P}}^2} + \text{h.c.} \right). \end{aligned} \quad (186)$$

The first one comes from the cross term between the derivative of the flat direction superpotential and the inflaton superpotential, the second is due to the Kähler potential coupling between the flat direction and the inflaton, and the third term is a cross term between the two superpotentials. All these terms give a generalized contribution equivalent to an A -term of the MSSM:

$$V(\phi) = H M_{\text{P}}^3 f \left(\frac{\phi^d}{M_{\text{P}}^d} \right), \quad (187)$$

where d is the dimensionality at which the flat direction is lifted. The induced A -term has an important role to play during the evolution of the flat direction. A possible A -term can also be generated from the expansion of the Kähler potential for field values $I, \phi < M_{\text{P}}$, which is of the form [46]

$$\frac{1}{M_{\text{P}}} \int d^4\theta I \phi^\dagger \phi + \text{h.c.} \sim \frac{F_I^\dagger F_\phi \phi}{M_{\text{P}}} + \text{h.c.},$$

$$\frac{1}{M_{\text{P}}} \int d^2\theta I W_i + \text{h.c.} \sim \frac{F_I}{M_{\text{P}}} W_i + \text{h.c.} \quad (188)$$

Note that the A -terms arise only from terms with a linear coupling of the inflaton superfield to a gauge invariant-operator ϕ_i . If I were a composite field rather than a singlet, then such a term will not arise and an A -term will not be generated. Also, in the case of D-term inflation, the inflaton cannot induce an A -term because $F_I = 0$. More generally, if there is a symmetry preventing a linear coupling of the inflaton, then order H A -terms can be eliminated also in F-term inflation. As long as the thermal bath of the inflaton decay products dominates over the low energy supersymmetry breaking scale, we should have Hubble induced corrections to $m_0^2, m_{1/2}, A$.

If there is a non-minimal dependence of the gauge superfield kinetic terms on the inflaton field, a Hubble-induced gaugino mass can also be produced. Generally the gauge superfield kinetic terms must depend on the field(s) of the hidden sector in order to obtain gaugino masses of roughly the same order as (or larger than) scalar masses, as required by phenomenology. Having $m_{1/2} \sim H$ thus appears to be quite natural unless an R -symmetry forbids terms which are linear in the inflaton superfield [46].

Since the μ -term does not break supersymmetry, there is a priori no reason to assume that a μ -term of order H will be created. (For a discussion, see [279]). In what follows we will treat μ as a free parameter.

So far we have not discussed the sign of the Hubble induced mass correction. In fact with a general Kähler term either sign is possible. Depending on the sign, the dynamical behavior of the AD field is completely different and therefore the predictions depend crucially upon the sign. There are however certain cases where the Hubble-induced terms might not occur at all. An R -symmetry [45, 46] or a special choice of the Kähler potential could forbid the AD field getting the Hubble-induced mass correction [226, 305].

4.6 The potential for flat direction

4.6.1 F-term inflation

Let us collect all the terms which contribute to the flat direction potential, which in the case of F-term inflation can be written as [45, 46]

$$V(\phi) = -C_I H_I^2 |\phi|^2 + \left(a \lambda_d H \frac{\phi^d}{d M^{d-3}} + \text{h.c.} \right) + m_\phi^2 |\phi|^2 + \left(A_\phi \lambda_d \frac{\phi^d}{d M^{d-3}} + \text{h.c.} \right), \\ + |\lambda|^2 \frac{|\phi|^{2d-2}}{M^{2d-6}}. \quad (189)$$

The first and the third terms are the Hubble-induced and low-energy soft mass terms, respectively, while the second and the fourth terms are the Hubble-induced and low-energy A terms. The last term is the contribution from the non-renormalizable superpotential. The coefficients $|C_I|$, a , $\lambda_d \sim \mathcal{O}(1)$, and the coupling $\lambda \approx 1/(d-1)!$. Note that low-energy A_ϕ term is dimensionful.

Note here the importance of the relative sign of the coefficient C_I . At large field values the first term dictates the dynamics of the AD field. If $C_I < 0$, the absolute minimum of the potential is $\phi = 0$ and during inflation the AD field will settle down to the bottom of the potential roughly in one Hubble time. In such case the AD field will not have any interesting classical dynamics. Its presence would nevertheless be felt because of quantum fluctuations. These would be chi-squared in nature since then the classical energy density of the AD field would be due to its own fluctuations.

If $C_I \ll 1$, the AD field takes some time to reach the bottom of the potential, and if it has a non-zero amplitude after the end of inflation, its dynamics is non-trivial.

The most interesting scenario occurs when $C_I > 0$. In this case the absolute value of the AD field settles during inflation to the minimum given by

$$|\phi| \simeq \left(\frac{C_I}{(d-1)\lambda_d} H_I M^{d-3} \right)^{1/d-2}. \quad (190)$$

Here we have ignored the potential term $\propto a$; if $C_I > 0$, the a -term will not change the vev qualitatively. On the other hand, even for $C_I < 0$ the potential Eq. (189) will have a minimum with a non vanishing vev if $|a|^2 > 4(d-1)C_I$. However the origin will also be a minimum in this case. The dynamics then depends on which minimum the AD field will choose during inflation.

The a -term in Eq. (189) violates the global $U(1)$ symmetry carried by ϕ . If $|a|$ is $\mathcal{O}(1)$, the phase θ of $\langle\phi\rangle$ is related to the phase of a through $n\theta + \theta_a = \pi$; otherwise θ will take some random value, which will generally be of $\mathcal{O}(1)$. This is the initial CP -violation which is required for baryogenesis/leptogenesis. In practice, the superpotential term lifting the flat direction is also the B and CP violating operator responsible for AD baryogenesis, inducing a baryon asymmetry in the coherently oscillating ϕ condensate.

4.6.2 D-term inflation

In D-term inflation one does not get the Hubble induced mass correction to the flat direction so that $C_I = 0$. Also the Hubble induced a -term is absent. However the Hubble induced mass correction eventually dominates once D-term induced inflation comes to an end. The potential for a generic flat direction during D-term inflation is given by

$$V(\phi) = m_\phi^2 |\phi|^2 + \left(A_\phi \lambda_d \frac{\phi^d}{d M^{d-3}} + \text{h.c.} \right) + |\lambda|^2 \frac{|\phi|^{2d-2}}{M^{2d-6}}, \quad (191)$$

and after the end of inflation the flat direction potential is given by [228]

$$V(\phi) = (m_\phi^2 - C H^2) |\phi|^2 + \left(A_\phi \lambda_d \frac{\phi^d}{d M^{d-3}} + \text{h.c.} \right) + |\lambda|^2 \frac{|\phi|^{2d-2}}{M^{2d-6}}, \quad (192)$$

where $C \sim \mathcal{O}(1)$. For C positive, the flat direction settles down to one of its minima given by Eq. (190) provided $\phi \geq \sqrt{m_\phi M/\lambda}$, otherwise

$$|\phi| \simeq \left(\frac{2C}{\lambda_d A_\lambda (d-1)} H(t)^2 M^{d-3} \right)^{1/d-2}, \quad (193)$$

Note that in this case that the A -term is also responsible for B and/or L , and CP violation. Another generic point to remember is that in R -parity conserving models the B and/or L violating operators must have even dimensions, so that $d = 4$ yields the minimal operator for AD baryogenesis.

5 Dynamics of flat directions

After the end of inflation $\langle\phi\rangle$ continues first to track the instantaneous local minimum of the scalar potential, obtained by replacing H_I with $H(t)$ in Eq.(190) or by following Eq. (193) in the D-term inflation case. Once $H \simeq m_0 \sim m_{3/2}$, the low-energy soft terms take over. Then m_ϕ^2 becomes positive and $\langle\phi\rangle$ starts to move in a non-adiabatic way (the phase of $\langle\phi\rangle$ differs from the phase of A -term during inflation). As a result $\langle\phi\rangle$ begins a spiral motion in a complex plane, which charges up the flat direction condensate, and eventually leads to generation of a net baryon and/or lepton asymmetry [46].

For baryogenesis purposes it is essential that the AD condensate obtains a non-zero vev during the inflationary epoch. In Sect. 4, we pointed out that a non-zero vev of the flat direction condensate is acquired only when the negative $(\text{mass})^2$ contribution dominates the potential. The MSSM flat directions which are made up of squarks and sleptons have Yukawa and gauge interactions. The couplings render the evolution of a particular flat direction non-trivially, especially when the flat direction has a time varying mass due to the Hubble expansion [305, 46, 307, 308]. Moreover, if thermalization is not instantaneous, thermal effects from reheating can be substantial and might trigger the motion of the flat direction at an earlier time, thereby changing the evolution of the flat direction condensate in a significant way [309, 310].

5.1 Running of the couplings

5.1.1 Running of gravitational coupling

Any flat direction has two kinds of interactions: renormalizable gauge or Yukawa interactions, and a non-trivial coupling to the curvature. Both types of interactions contribute to the logarithmic running of $(\text{mass})^2$ of the flat direction condensate. The coupling to the curvature is generic because in principle any scalar field in an expanding background receives a contribution from the curvature by virtue of the Lagrangian term $\xi R\phi^2$, where ξ is a coupling constant. Note that $R \propto +H^2$ in an expanding background. Any scalar field always gets an additional positive Hubble induced mass

correction, provided ξ is positive. The fundamental theory might have a conformal invariance, in which case the coupling strength $\xi = 1/6$ [311], but it is known that conformal invariance is not protected by any symmetry, and that quantum corrections always break conformal invariance. Especially for the flat direction condensate, spontaneously broken supersymmetry induces soft supersymmetry breaking terms which break conformal invariance, and the value of ξ remains undetermined.

It is of course possible to simply set $\xi = 0$. If initially $\xi = 0$ at some high scale, renormalization effects due to scalar field self-interaction will nevertheless generate a non-zero ξ at lower scales. In an expanding Universe the value of ξ also changes under the influence of a varying curvature (see [312], and references therein). In the simplest case of a single scalar field with a quartic self-interaction strength λ leads (at one-loop level) to a logarithmically running ξ [313]

$$\xi_{eff} = \xi + \left(\xi - \frac{1}{6}\right) \frac{12}{4\pi^2} \lambda \ln \left(\left| \frac{m^2 + \left(\xi - \frac{1}{6}\right) R}{m^2} \right| \right). \quad (194)$$

It is obvious that $\xi = 1/6$ is a fixed point of the RG equation. If the theory has fermions and gauge fields, then obviously the coefficient in front of the logarithmic term in Eq. (194) will be modified [312].

As we have seen in Sect. 4.2, when supersymmetry is promoted to a local theory, a supergravity correction is induced to the flat direction which is proportional to the curvature, and supergravity theories also allow for $\xi R\phi^2$ (e.g. superconformal supergravity [270]).

In the context of MSSM flat directions we have implicitly assumed $\xi = 0$. This is justified from the very definition of F- and D-flat directions. The only leading order self coupling term in the flat direction potential is the Hubble induced A-term in Eq. (189). The overall self coupling constant is relatively large when the flat direction is lifted at $d = 4$, i.e. the suppression is proportional to $\mathcal{O}(1)(H/M_P)$, where we have replaced M by M_P in Eq. (189). In any inflation model the ratio $H_I/M_P \ll 1$, which in conjunction with Eq. (194), suggests that the effect of running on ξ is minimal. For a running ξ the curvature term in Eq. (194) dominates over the mass term. This might not be the case with the flat direction condensate because the condensate also

receives a field dependent mass while it is evolving. As long as the vev dependent mass is larger than the curvature induced mass, the running of any parameter in the theory will be dictated mainly by the renormalizable quantum effects. We therefore conclude that the running of ξ can be neglected.

When the field dependent mass of the flat direction field becomes of order $m^2 \sim \mathcal{O}(H^2)$, it might be prudent to start worrying about the curvature induced term, especially during inflation. A simple inspection of Eq. (194) suggests that ξ is always of order $\ln(\mathcal{O}(1))$, with virtually no alteration in ξ_{eff} . From now onwards we fix the non-minimal coupling to be $\xi = 0$.

5.1.2 Renormalization group equations in the MSSM

Let us consider the running of the flat direction (mass)² below M_{GUT} by assuming that it is the scale where supersymmetry breaking is transmitted to the visible sector, in order to avoid uncertainties about physics between M_{GUT} and M_{P} . The running of low-energy soft breaking masses has been studied in great detail in the context of MSSM phenomenology [314], in particular in connection with radiative electroweak symmetry breaking [315].

Let us recall some of the salient features of the MSSM one-loop RG equations. The ones relevant to flat directions involve the Higgs doublet H_u which couples to the top quark, the right-handed stop $\tilde{u}_3$, the left-handed doublet of third generation squarks $\tilde{Q}_3$ and the A -parameter A_t associated with the top Yukawa interaction. The RG equations read [34]

$$\begin{aligned}
\frac{d}{dq} m_{H_u}^2 &= \frac{3h_t^2}{8\pi^2} \left(m_{H_u}^2 + m_{\tilde{Q}_3}^2 + m_{\tilde{u}_3}^2 + |A_t|^2 \right) - \frac{1}{2\pi^2} \left(\frac{1}{4} g_1^2 |m_1|^2 + \frac{3}{4} g_2^2 |m_2|^2 \right), \\
\frac{d}{dq} m_{\tilde{u}_3}^2 &= \frac{2h_t^2}{8\pi^2} \left(m_{H_u}^2 + m_{\tilde{Q}_3}^2 + m_{\tilde{u}_3}^2 + |A_t|^2 \right) - \frac{1}{2\pi^2} \left(\frac{4}{9} g_1^2 |m_1|^2 + \frac{4}{3} g_3^2 |m_3|^2 \right), \\
\frac{d}{dq} m_{\tilde{Q}_3}^2 &= \frac{h_t^2}{8\pi^2} \left(m_{H_u}^2 + m_{\tilde{Q}_3}^2 + m_{\tilde{u}_3}^2 + |A_t|^2 \right) - \frac{1}{2\pi^2} \left(\frac{1}{36} g_1^2 |m_1|^2 + \frac{3}{4} g_2^2 |m_2|^2 + \frac{4}{3} g_3^2 |m_3|^2 \right), \\
\frac{d}{dq} A_t &= \frac{3h_t^2}{8\pi^2} A_t - \frac{1}{2\pi^2} \left(\frac{13}{36} g_1^2 m_1 + \frac{3}{4} g_2^2 m_2 + \frac{4}{3} g_3^2 m_3 \right).
\end{aligned} \tag{195}$$

Here q denotes the logarithmic scale; this could be an external energy or momentum scale, but in the case at hand the relevant scale is set by the vev(s) of the fields

themselves. h_t is the top Yukawa coupling, while g_i and m_i are respectively the gauge couplings and soft breaking gaugino masses of $U(1)_Y \times SU(2) \times SU(3)$. If h_t is the only large Yukawa coupling (i.e. as long as $\tan\beta$ is not very large), the beta functions for $(\text{mass})^2$ of squarks of the first and second generations and sleptons only receive significant contributions from gauge/gaugino loops. A review of these effects can be found in [314]. Here we only mention the main results for the case of universal boundary conditions, where at M_{GUT} all the scalar masses are m_0^2 and the gauginos have a common soft breaking mass $m_{1/2}$. For a low value of $\tan\beta = 1.65$ ³,

$$m_{H_u}^2 \simeq -\frac{1}{2}m_0^2 - 2m_{1/2}^2 \quad (196)$$

at the weak scale, while $m_{u_3}^2$ and $m_{\tilde{Q}_3}^2$ remain positive. The soft breaking $(\text{mass})^2$ of the first and second generations of squarks is $\simeq m_0^2 + (5 - 7)m_{1/2}^2$, while for the right-handed and left-handed sleptons one gets $\simeq m_0^2 + 0.1m_{1/2}^2$ and $\simeq m_0^2 + 0.5m_{1/2}^2$, respectively. The important point is that the sum $m_{H_u}^2 + m_L^2$, which describes the mass along the $H_u L$ flat direction, is driven to negative values at the weak scale only for $m_{1/2} \gtrsim m_0$. This is intuitively understandable, since Eqs.(195) have a fixed point solution [317] $m_{H_u}^2 + m_{u_3}^2 + m_{\tilde{Q}_3}^2 = A_t = 0$ when $m_{1/2} = 0$.

5.2 Hubble induced radiative corrections

Here we describe radiative corrections in a cosmological set-up relevant for the AD mechanism [308]. When the Hubble induced supersymmetry breaking is dominant, i.e. for $H > \mathcal{O}(\text{TeV})$, the evolution of the soft terms is different from the vacuum RG equations given in Eq. (195). For the low-energy supersymmetry breaking case, constraints from the weak scale (e.g. realization of electroweak symmetry breaking, and experimental limits on the sparticle masses) give information about the soft breaking parameters m_0^2 and $m_{1/2}$. Together with fine tuning arguments, these constraints

³This value corresponds to the case of maximal top Yukawa coupling, so called fixed point scenario [316, 317]. Such a low value of $\tan\beta$ is excluded by Higgs searches at LEP [318], unless one allows stop masses well above 1 TeV. We nevertheless include this scenario in our discussion since it represents an extreme case.

imply that $m_0^2 > 0$ and that $m_0, m_{1/2}$ are $\mathcal{O}(\text{TeV})$. In the Hubble induced supersymmetry breaking case m_0^2 and $m_{1/2}$ are determined by the scale of inflation (and the form of the Kähler potential). At low scales the Hubble induced terms are completely negligible because at temperature $T \sim M_W$, $H \sim \mathcal{O}(1)$ eV; at present the Hubble parameter is tiny, $H_0 \sim \mathcal{O}(10^{-33})$ eV.

There exists an even more fundamental difference between the Hubble induced and ordinary radiative corrections. In Minkowski space the loop contributions to beta functions freeze at a scale of the order of the mass of the particles in the loop. In an expanding Universe the horizon radius $\propto H^{-1}$ defines an additional natural infrared cut-off for the theory. The masses of particles coupled to the flat direction receive contributions from two sources. There is a supersymmetry preserving part proportional to the vev $\langle\phi\rangle$, and the Hubble induced supersymmetry breaking part. The loop contributions to beta functions should thus be frozen at a scale given by the largest of $|\langle\phi\rangle|$ and H (recall that h_t and gauge couplings are close to one). In particular, if the squared mass of the flat direction condensate is positive at very large scales but turns negative at some intermediate scale Q_c , the origin of the flat direction potential will cease to be a minimum, provided the Hubble parameter is less than Q_c . On the other hand, if $m_\phi^2 < 0$ at the GUT scale, its running should already be terminated at the scale $|\langle\phi\rangle|$ determined by Eq.(190).⁴

In the following two subsections we discuss separately the cases of positive and negative GUT-scale (mass)² for the flat direction condensate.

5.2.1 The case with $C_I \approx -1$

In this case all scalar fields roll towards the origin very rapidly and settle there during inflation, provided radiative corrections to their masses are negligible. A typical flat

⁴Here we note that the Hubble cut-off usually plays no role in loop corrections to the inflaton potential. In most inflation models the masses of the fields which may run in the loop are larger than the Hubble expansion rate due to the presence of a finite coupling to the inflaton. This will happen if the (time varying) inflaton vev is large and the couplings are not very small. In those cases, which are somewhat similar to our case with $C_I > 0$, one can trust the usual loop calculation evaluated in a flat space time background [218].

A_t/H	$m_{1/2}/H$	$Q_c(h_t = 2)$	$Q_c(h_t = 0.5)$
+1/3 (-1/3)	1/3	$\times$	$\times$
+1/3 (-1/3)	1	$10^6 - 10^7$	10^3
+1/3 (-1/3)	3	10^{11}	$10^6 - 10^7$
+1 (-1)	1/3	$\times$	$\times$
+1 (-1)	1	$10^6 - 10^7$	10^5 ($\times$)
+1 (-1)	3	10^{11}	10^8 (10^6)
+3 (-3)	1/3	$\times$	10^7
+3 (-3)	1	10^{14} (10^7)	10^9 (10^3)
+3 (-3)	3	10^{15} (10^{11})	10^{10} (10^6)

Table 2: The scale Q_c (in GeV) where the squared mass of the $H_u L$ flat direction changes sign, shown for $C_I = -1$ and several values for the ratios A_t/H and $m_{1/2}/H$ as well as the top Yukawa coupling h_t , all taken at scale $M_{\text{GUT}} = 2 \cdot 10^{16}$ GeV, from [308]

direction condensate ϕ is a linear combination $\phi = \sum_{i=1}^N a_i \varphi_i$ of the MSSM scalars φ_i , implying that $m_\phi^2 = \sum_{i=1}^N |a_i|^2 m_\varphi^2$. In [308], it was noticed that with small values for the μ parameter, the running of m_ϕ^2 crucially depends on $m_{1/2}$.

Let us consider sample cases with gaugino masses $m_{1/2} = (H; 3H; H/3)$, the A-term⁵ $A_t(M_{\text{GUT}}) = (\pm H; \pm 3H; \pm H/3)$, top Yukawa $h_t(M_{\text{GUT}}) = (2, 0.5)$ and couplings $g_1(M_{\text{GUT}}) = g_2(M_{\text{GUT}}) = g_3(M_{\text{GUT}}) = 0.71$, and follow the running of scalar soft masses from M_{GUT} down to 10^3 GeV, where low-energy supersymmetry breaking becomes dominant. The main result is that only the LH_u flat direction can acquire a negative (mass)² at low scales. In this case $m_\phi^2 = (m_{H_u}^2 + m_L^2 + \mu^2)/2$, where the last term is from the Hubble induced μ term. The results are summarized in Table 2, where it has been assumed that $\mu(M_{\text{GUT}}) \lesssim H/4$ so that the μ -term contribution to m_ϕ^2 is negligible. In general m_ϕ^2 changes sign at a higher scale for $h_t(M_{\text{GUT}}) = 2$.

⁵The RG equations (195) for A_t show that the relative sign between A_t and $m_{1/2}$ matters, since it affects the running of $|A_t|$, and subsequently, scalar soft masses. Without loss of generality we take the common gaugino mass $m_{1/2}$ to be positive.

This is expected since a large Yukawa coupling naturally maximizes the running of $m_{H_u}^2$. Furthermore, the difference between $A_t/m_{1/2} < 0$ and $A_t/m_{1/2} > 0$ becomes more apparent as $|A_t/m_{1/2}|$ increases and h_t decreases. The quasi fixed-point value of $A_t/m_{1/2}$ is positive [316, 317]. Positive input values of A_t will thus lead to positive A_t at all scales, but a negative $A_t(M_{\text{GUT}})$ implies that $A_t \simeq 0$ for some range of scales, which diminishes its effect in the RG equations, see Eq. (195). The sign of $A_t(M_{\text{GUT}})$ is more important for smaller h_t , since then $A_t/m_{1/2}$ will evolve less rapidly.

It was noticed in [308] that the squared mass of the $H_u L$ flat direction does not change sign when $m_{1/2} = H/3$, except for⁶ $A_t = \pm 3H$ and $h_t = 0.5$. This can be explained by the fact that for small $m_{1/2}$ and small or moderate $|A_t|$ we are generally close to the fixed point solution

$$m_{H_u}^2 \simeq -\frac{1}{2}H^2; \quad m_{u_3}^2 \simeq 0; \quad m_{Q_3}^2 \simeq \frac{1}{2}H^2. \quad (197)$$

Nevertheless, even for $m_{1/2} \ll H$ the squared mass of the LH_u flat direction as well as $m_{u_3}^2$ are $< 0.2H^2$ above 1 TeV, because of the fixed point behavior. This implies that the LH_u flat direction can still be viable for baryogenesis, as pointed out by McDonald [324]. Flat directions built out of $\tilde{u}_3$ will be marginal at best, since the decrease in $m_{u_3}^2$ will be counteracted by other contributions to m_ϕ^2 ; e.g. for the $\bar{u}_3 \bar{d}_1 \bar{d}_2$ flat direction we find $m_\phi^2 > 2H^2/3$ at all scales.

The AD mechanism for baryogenesis should always work if $Q_c > H_I$, since in that case the global minimum of the potential during inflation is located at $|\langle \phi \rangle| \neq 0$. Note that in this case the vev $|\langle \phi \rangle|$ is usually determined by Q_c rather than by Eq. (190).

For scales close to Q_c the mass term in the scalar potential Eq. (189) can be written as $\beta_\phi H^2 |\phi|^2 \log(|\phi|/Q_c)$, where the coefficient β_ϕ can be obtained from the RG equations. If $\beta_\phi > 0$, which is true for the $H_u L$ flat direction for $C_I < 0$, this term will reach a minimum at $\log(|\phi|/Q_c) = -1$. If $Q_c < (H_I M_{\text{GUT}}^{d-3})^{1/d-2}$ the non-renormalizable contributions to the scalar potential are negligible for $|\phi| \sim Q_c$, so that the minimum of the quadratic term essentially coincides with the minimum of the complete potential given by Eq. (189). In models of high scale inflation (e.g. chaotic inflation models),

⁶For this choice of parameters, A_t runs initially very slowly. It will therefore remain large for some time and helps $m_{H_u}^2$ to decrease quickly towards lower scales.

the Hubble constant during inflation H_I can be as large as 10^{13} GeV. This implies that m_ϕ^2 for the $H_u L$ flat direction can only become negative during inflation if $m_{1/2}^2 \gg H^2$, which includes the “no-scale” scenario studied in [305].

The region of the parameter space safely allowing AD leptogenesis is much larger in models of intermediate and low scale inflation (e.g. some new inflation models [14]) where H_I is substantially smaller. In such models one can easily have $H_I < Q_c$ at least for the $H_u L$ flat direction, unless $m_{1/2}^2 \ll H^2$ or $\mu^2 \gtrsim m_{1/2}^2$ [308].

If $Q_c < H_I$, the condensate ϕ settles at the origin during inflation and its post-inflationary dynamics will depend on the process of thermalization. If the inflaton decay products thermalize very slowly, m_ϕ^2 is only subject to zero-temperature radiative corrections and $\langle \phi \rangle$ can move away from the origin once $H \lesssim Q_c$; a necessary condition for this scenario is that inflatons do not directly decay into fields that are charged under $SU(3) \times SU(2) \times U(1)_Y$. If $Q_c \gg 1$ TeV, ϕ will readily settle at the new minimum and AD leptogenesis can work.

The situation will be completely different if inflatons directly decay into some matter fields. In such a case the plasma of inflaton decay products has a temperature $T \sim (\Gamma_d H M_{\text{Planck}}^2)^{1/4}$ [22] (Γ_d is the inflaton decay rate). Fields which contribute to the running of m_ϕ^2 are in thermal equilibrium (recall that the flat direction field is stuck at $\phi = 0$) and their back reaction results in thermal corrections of order $+T^2$ to m_ϕ^2 . For generic models of inflation $T > H$, implying that thermal effects exceed radiative corrections. Therefore $\langle \phi \rangle$ remains at the origin at all times and AD leptogenesis will not work.

5.2.2 The case with $C_I \approx +1$

In this case all flat directions are viable for baryogenesis purposes provided the running of m_ϕ^2 is negligible. Radiative corrections may change the sign (in this case to positive) at small vev(s), resulting in the entrapment of ϕ at the origin.

A quantitative study of the sample cases discussed above can be summarized as follows [308]. The squared mass of the LH_u flat direction is always negative at small scales, unless $\mu^2 \gtrsim H^2/2$. For $m_{1/2} = 3H$, m_ϕ^2 changes sign twice; it is positive for

scales Q between roughly 10^{14} and 10^6 GeV, the precise values depending on h_t and A_t . Slepton masses only receive positive contributions from electroweak gauge/gaugino loops. As a result, the squared mass of the $LL\bar{e}$ flat direction remains negative down to 1 TeV, unless $m_{1/2} > 2H$; for $m_{1/2} \gtrsim 3H$, $Q_c \gtrsim 10^9$ GeV even for this flat direction. The squared masses of all squarks (except $\tilde{u}_3$) change sign at $Q_c > 1$ TeV unless $m_{1/2} \lesssim H/3$; we find $Q_c \simeq 10^{10}$ (10^{15}) GeV for $m_{1/2}/H = 1$ (3). This is due to the large positive contribution $\propto m_3^2$ to the squared squark masses at scales below M_{GUT} . The corresponding values for the $\bar{u}_3\bar{d}_i\bar{d}_j$ and $LQ\bar{d}$ flat directions are usually somewhat smaller, due to the Yukawa terms in the β -function and the slower running of the slepton masses, respectively; however, the values of Q_c listed in Table 2 are still a fair approximation.

The positive contribution to the scalar potential from the non-renormalizable superpotential term now dominates $-H^2$ (see Eq. (190)). If $Q_c > (H_I M_{\text{GUT}}^{d-3})^{1/d-2}$, m_ϕ^2 is positive for all vev(s) and hence the flat direction will settle at the origin during inflation and remain there. In such a case the flat direction is not suitable for AD baryogenesis. This can easily happen for flat directions involving squarks in models with low scale inflation, but is not likely for high scale inflation models (unless $m_{1/2} \gtrsim 3H$). For $H_I < Q_c < (H_I M_{\text{GUT}}^{d-3})^{1/d-2}$, feasible for some flat directions in both intermediate/high scale and low scale models. During inflation the potential has two minima, at $\langle\phi\rangle = 0$ and at $|\langle\phi\rangle| \sim (H_I M_{\text{GUT}}^{d-3})^{1/d-2}$. Depending on the initial conditions, ϕ can roll down towards either of them and settle there but only the latter one will be useful for AD baryogenesis.

If $Q_c < H_I$, the flat direction condensate will settle at the value determined by Eq. (190) (the only minimum during inflation) and remain there. The appearance of another minimum at the origin after inflation, which is possible once $H < Q_c$, does not change the situation since these minima are separated by a barrier. In this case radiative corrections will not change the picture qualitatively; however, they will still modify the quantitative analysis, since C_I in Eq.(190) will become scale-dependent.

In brief, the main conclusion is that among the flat directions LH_u is the only robust one in the sense that it gives rise to AD leptogenesis independently of the sign

of C_I .

5.2.3 Running of the flat direction field in no-scale supergravity

So far we have dealt with a minimal choice of the Kähler potential. An alternative is non-flat Kähler potential; an example of this is provided by e.g. no-scale models, for which $K \sim \ln(z + z^* + \phi_i^\dagger \phi)$, where z belongs to supersymmetry breaking sector, and ϕ_i belongs to the matter sector, and both are measured in terms of reduced Planck mass (for a review, see [303]). In no-scale models there exists an enhanced symmetry known as the Heisenberg symmetry [325], which is defined on the chiral fields as $\delta z = \epsilon^* \phi^i$, $\delta \phi^i = \epsilon^i$, and $\delta y^i = 0$, where y^i are the hidden sector fields, such that the combinations $\eta = z + z^* - \phi_i^* \phi^i$, and $y_i = 0$ are invariant. For a especial choice

$$K = f(\eta) + \ln[W(\phi)/M_{\text{P}}^3]^2 + g(y), \quad (198)$$

The $N = 1$ supergravity potential reads [305, 307]

$$V = e^{f(\eta)+g(y)} \left[\left(\frac{f'^2}{f''} - 3 \right) \frac{|W|^2}{M_{\text{P}}^2} - \frac{1}{f'^2} \frac{|W_i|^2}{M_{\text{P}}^2} + g_a (g^{-1})^a_b g^b \frac{|W|^2}{M_{\text{P}}^2} \right]. \quad (199)$$

Note that there is no cross term in the potential such as $|\phi_i^* W|^2$. As a consequence any tree level flat direction remains flat even during inflation [305] (in fact it is the Heisenberg symmetry which protects the flat directions from obtaining Hubble induced masses [325]). The symmetry is broken by gauge interactions or by coupling in the renormalizable part of the Kähler potential. Then the mass of the flat direction condensate arises from the running of the gauge couplings.

For $f(\eta) = -3 \ln \eta$, the one-loop corrected supergravity induced mass term has been calculated in [326, 327, 305], which gives an effective mass for the flat direction field in the presence of finite energy density stored in the inflaton sector. The typical mass of the flat direction has been computed and comes out to be $m_\phi^2 \sim 10^{-2} H^2$ during inflation. The only constraint is that flat direction must not involve stops [305].

5.3 Post-inflationary running of the flat direction

Now we focus on the running of the flat direction after inflation. Here we must take into account the low energy supersymmetry breaking effects. In particular, the running of

the condensate mass will depend on how supersymmetry is transmitted to the visible sector.

5.3.1 Gravity mediated supersymmetry breaking

For gravity-mediated supersymmetry breaking the scalar potential along a flat direction has been evaluated as [48, 49]

$$U(\Phi) \approx m_\phi^2 \left(1 + K \log \left(\frac{|\Phi|^2}{M^2} \right) \right) |\Phi|^2 + \frac{\lambda^2 |\Phi|^{2(d-1)}}{M_P^{2(d-3)}} + \left(\frac{A_\lambda \lambda \Phi^d}{d M_P^{d-3}} + h.c. \right), \quad (200)$$

where m_ϕ is the conventional gravity-mediated soft supersymmetry breaking scalar mass term ($m_\phi \approx 100$ GeV), K is a parameter which depends on the flat direction, and the logarithmic contribution parameterizes the running of the flat direction potential with $M = (M_P^{d-3} m_{3/2} / |\lambda|)^{(1/(d-2))}$. In the gravity mediated case $|A_\lambda| < d m_{3/2}$, for $d = 4, 6$.

K can be computed from the RG equations, which to one loop have the form

$$\frac{\partial m_i^2}{\partial t} = \sum_g a_{ig} m_g^2 + \sum_a h_a^2 \left(\sum_j b_{ij} m_j^2 + A^2 \right), \quad (201)$$

where a_{ig} and b_{ij} are constants, m_g is the gaugino mass, h_a the Yukawa coupling, A is the A -term, and $t = \ln M_X / q$. The full RG equations have been listed in [34]. The potential along the flat direction is then characterized by the amount of stop mixture (where appropriate), the values of gluino mass and A , and in the special case of the $d = 4$ $H_u L$ -direction, on the $H_u H_d$ -mixing mass parameter μ_H .

The mass of the AD scalar ϕ is the sum of the masses of the squark and slepton fields ϕ_i constituting the flat direction, $m_S^2 = \sum_a p_i^2 m_i^2$, where p_i is the projection of ϕ along ϕ_i , and $\sum p_i^2 = 1$. The parameter K is then given simply by

$$K = \frac{1}{q^2} \frac{\partial m_S^2}{\partial t} \Big|_{t=\log q}. \quad (202)$$

To compute K , one has to choose the scale q . The appropriate scale is given by the value of the AD condensate amplitude when it first begins to oscillate at $H \approx m_\phi$ or

$$Q = |\phi_0| = \left[\frac{m_\phi^2 M^{2(d-3)}}{(d-1)\lambda^2} \right]^{\frac{1}{2(d-2)}}, \quad (203)$$




Figure 1: Contours of K for two $d=4$ flat directions in the $(A, \xi \equiv m_g/m(t=0))$ -plane: (a) $K = 0$ (b) $K = -0.01$; (c) $K = -0.05$; (d) $K = -0.1$. The directions are (i) $Q_3 Q_3 Q L$; (ii) $Q Q Q L$, no stop; (iii) $\bar{u}_3 \bar{u} \bar{d} \bar{e}$; (iv) $\bar{u} \bar{u} \bar{d} \bar{e}$ with equal weight for all $\bar{u}$ -squarks, from [329].

The RG running of the flat directions in the case of gravity mediated supersymmetry breaking was studied in [329, 330], where unification at $t = 0$ was assumed and all the other Yukawa couplings except the top Yukawa were neglected.

The contours of K for the $d = 4$ $\bar{u} \bar{u} \bar{d} \bar{e}$ and $Q Q Q L$ directions are shown in Fig. (1) in the (A, ξ) -plane, where $\xi \equiv m_g/m(t=0)$ (for $\tan \beta = 1$ and $\lambda = 1$). These are representative of all the other directions, too, except for $H_u L$. For $\xi \sim \mathcal{O}(1)$, typical value for K is found to be about -0.05 . Similar contours can be obtained for the $d = 6$ $(\bar{u} \bar{d} \bar{d})^2$ and $(Q L \bar{d})^2$ directions, see [329]. For all the squark directions with no stop, as long as h_b and h_u can be neglected, K is always negative, and the contours of equal K do not depend on A . In the presence of stop mixing $K < 0$ is no longer automatic

even in the purely squark directions. The more there is stop, the larger value of ξ is required for $K < 0$. Even for pure stop directions, positive K is typically obtained only for relatively light gaugino masses with $\xi \lesssim 0.5$.

In contrast to the squark directions, K was found [329] to be always positive in the $H_u L$ -direction. This is due to the fact $H_u L$ does not involve strong interactions which in other directions are mainly responsible for the decrease of the running scalar masses. Very roughly, instability is found when $m_{\tilde{g}} \gtrsim m_{\tilde{t}}$, although the exact condition should be checked case by case. In general, the sign of K could be deduced from the observation of SUSY parameters such as $\tan \beta$, the gluino mass and the supersymmetry breaking parameter m_ϕ [329].

5.3.2 Gauge mediated supersymmetry breaking

A similar analysis can be made for the gauge mediated case, where supersymmetry breaking is transmitted to the observable sector below some relatively low messenger sector scale Λ_S , above which the potential is completely flat (see Sect. 4.4.3.). In the gravity mediated scenario the soft masses stay intact, modulo RG running, up to the Planck scale; in gauge mediation the masses simply disappear above Λ_S . For a large condensate vev, one can integrate out the gauge and chiral fields coupled to the flat direction in order to obtain an effective low energy theory. In such a case, as was first pointed out by Kusenkov and Shaposhnikov [151], the potential along the flat direction obtains a logarithmic correction of the form [151, 331]

$$U(\Phi) = m_\phi^4 \log \left(1 + \frac{|\Phi|^2}{m_\phi^2} \right) + \frac{\lambda^2 |\Phi|^{2(d-1)}}{M_{\text{P}}^{2(d-3)}} + \left(\frac{A_\lambda \lambda \Phi^d}{d M_{\text{P}}^{d-3}} + \text{h.c.} \right). \quad (204)$$

where $m_\phi \sim 1 - 100$ TeV. Because of the differences in the potential, the dynamical evolution of the condensate field will be markedly different from the gravity mediated case. Because the messenger sector is not constrained by experiments, one cannot provide a detailed description of the mass parameters. Here one should note that in order to have an AD condensate, the A -term is actually constrained. In the gauge mediated case $|A_\lambda| \leq (10^{-4} - 10^{-7}) m_\phi$, for $d = 4, 6$.

Eqs. (200) and (204) are two book-keeping equations which are useful for the rest of this review.

5.4 Density perturbations from the flat direction condensate

The role of MSSM flat directions is not just limited to generating the lepton and/or baryon asymmetry in the Universe, but they also play an interesting role in the dynamics of density perturbations.

5.4.1 Energetics of flat direction and the inflaton field

Once the flatness of the flat direction potential is lifted by non-renormalizable terms, for large field values the condensate energy density can dominate over the inflaton potential. This could be disastrous: either inflationary expansion would come to a halt, or the flat direction condensate fluctuations might ruin the successful predictions for the angular power spectrum [52, 332, 53, 54].

In [53], the generation of adiabatic density perturbations was studied for both D- and F-term inflation models. Note that in the former case there is no Hubble induced mass correction to the flat direction condensate. The scalar potential for F- and D-flat direction of dimension d is given by (see Eq. (191) in Sect. 4.6.2.)

$$V(\phi) \approx \frac{\lambda^2 |\Phi|^{2(d-1)}}{M^{2(d-3)}}, \quad (205)$$

where only the dominant term from Eq. (191) corresponding to superpotential term of the form $W = \lambda \Phi^n / n M^{d-3}$ has been kept. Throughout this discussion R -parity is conserved and therefore we deal only with even dimensions $d = 4, 6, 8$.

For illustrative purposes let us assume that the D-term inflationary potential is given by (see Eq. (146) in Sect. 3.5.3.)

$$V(S) = \frac{g^2 \xi^4}{2} + \frac{g^4 \xi^4}{32\pi^2} \ln \left(\frac{S^2}{Q^2} \right), \quad (206)$$

where S is the inflaton component; Q is here the renormalization scale. For a large initial vev for ϕ , $S \sim \mathcal{O}(M_P)$, the dynamics is first dominated by $V(\phi)$. For a sufficiently large vev of ϕ the effective condensate mass squared $V''(\phi)$, becomes larger than H^2 . This occurs if $\phi > \phi_H$, where [53]

$$\phi_H = \frac{2^{\frac{(d-1)}{2(d-2)}}}{(6(2d-2)(2d-3))^{\frac{1}{2(d-2)}}} \left(\frac{g}{\lambda} \right)^{\frac{1}{d-2}} \xi^{\frac{2}{d-2}} M_P^{\frac{d-4}{d-2}}. \quad (207)$$

If initially $\phi_i > \phi_H$, then ϕ will first oscillate in its potential with a decreasing amplitude: $\phi(t) \propto a^{-3/d}(t)$ [230]. This period ends before the onset of inflaton domination. The system then enters a regime where both ϕ and S are slowly rolling.

The slow rolling dynamics of the scalar fields is given by the solution of

$$3H\dot{\Psi}_a = -\frac{\partial V(\Psi_a)}{\partial \Psi_a} \quad ; \quad H = \left(\frac{\sum_a V(\Psi_a)}{3M_{\text{P}}^2} \right)^{1/2}, \quad (208)$$

where $\Psi_a \equiv S, \phi$. By taking the ratio of the equations for ϕ and S , one obtains

$$\frac{\partial \phi}{\partial S} = \frac{16\pi^2(d-1)\lambda^2\phi^{(2d-3)}S}{2^{d-2}g^4\xi^4M_{\text{P}}^{2(d-3)}}, \quad (209)$$

which has a general solution of the form

$$\phi = \phi_i \left[1 + \alpha_d \phi_i^{2d-4} (S_i^2 - S^2) \right]^{-1/(2d-4)} \quad ; \quad \alpha_d = \frac{16\pi^2(d-2)(d-1)\lambda^2}{2^{d-2}M_{\text{P}}^{2(d-3)}g^4\xi^4}, \quad (210)$$

where ϕ_i and S_i are the initial values at the onset of inflation. There are two features about this solution. First, since S_i is large compared with the value of S at $N = 50$ e-foldings before inflation, we see that for sufficiently large ϕ_i the value of ϕ at late times is *fixed* by S_i ,

$$\phi \equiv \phi_* \approx \left(\frac{1}{\alpha_d} \right)^{\frac{1}{2d-4}} \frac{1}{S_i^{1/(d-2)}}. \quad (211)$$

This is true if $\phi_i > \phi_*$, otherwise, ϕ simply remains at ϕ_i . Second, we can relate S_i to the total number of e-foldings during the $V(S)$ dominated period of inflation. In general, for sufficiently large ϕ_i , we could have an initial period of $V(\phi)$ dominated inflation. During this period S does not significantly change from S_i . The potential is dominated by $V(\phi)$ once $\phi > \phi_S$, where [53]

$$\phi_S = \frac{\sqrt{2}M_{\text{P}}^{\frac{d-3}{d-1}}}{\lambda^{\frac{1}{d-1}}} \left(\frac{g^2\xi^4}{2} \right)^{\frac{1}{2(d-1)}}. \quad (212)$$

ϕ_S is generally less than ϕ_H (see Eq. (207)), therefore ϕ will be slow rolling during $V(S)$ domination.

From Eq. (210), we find that the condition for S to change significantly from S_i at a given value of ϕ is given by

$$S_i < \frac{1}{\alpha_d^{1/2}} \left(\frac{1}{\phi} \right)^{d-2}. \quad (213)$$

The condition for S to change significantly during $V(\phi)$ dominated inflation is given by Eq. (213) with $\phi = \phi_S$; one finds

$$S_i < S_{i_c} \approx \frac{2^{\frac{d-2}{2(d-1)}} g^{\frac{d}{d-1}} \xi^{\frac{2}{d-1}} M_{\text{P}}^{\frac{d-3}{d-1}}}{4\pi \lambda^{\frac{1}{d-1}}} . \quad (214)$$

Since S_{i_c} is small compared with M_{P} , whereas the value of S required to generate 50 e-foldings of inflation, $S_{50} = g\sqrt{50}M_{\text{P}}/(2\pi)$ is close to M_{P} , it follows that $S_i (> S_{50})$ will generally be larger than S_{i_c} , and so the inflaton will remain at S_i until the Universe becomes inflaton dominated.

In this case the total number of e-foldings during inflaton domination is given by N_S where $S_i = (g/2\pi)N_S^{1/2}M_{\text{P}}$. If $\phi_i > \phi_*$, then ϕ at $N \approx 50$ e-foldings of inflation will be given by [53]

$$\phi_* \approx \left(\frac{1}{\alpha_d}\right)^{\frac{1}{2d-4}} \left(\frac{2\pi}{gM_{\text{P}}N_S^{1/2}}\right)^{\frac{1}{d-2}} . \quad (215)$$

Note that the dependence on N_S is quite weak; for the case of $d = 4$ ($d = 6$) AD baryogenesis, $\phi_* \propto N_S^{-1/4}$ ($N_S^{-1/8}$). If there is no large number of inflationary e-foldings one can essentially fix the value of ϕ_* . In this case one can *predict* the magnitude of the baryonic isocurvature perturbation.

Imposing a chaotic-type initial condition $V(\phi_i) \approx M_{\text{P}}^4$ yields

$$\phi_i \approx \frac{\sqrt{2}M_{\text{P}}}{\lambda^{\frac{1}{d-1}}} . \quad (216)$$

By directly solving the slow roll equations for ϕ and S , we obtain the total number of e-foldings of inflation:

$$N_T = N_\phi + N_S \approx \frac{1}{4(d-1)} \frac{\phi_i^2}{M_{\text{P}}^2} + \frac{4\pi^2 S_i^2}{g^2 M_{\text{P}}^2} , \quad (217)$$

where N_ϕ is the number of e-foldings during $V(\phi)$ domination, provided $\phi_i > \phi_S$. $V(S)$ will dominate the total number of e-foldings only if

$$N_S > \frac{1}{2(d-1)\lambda^{2/(d-1)}} . \quad (218)$$

Since $N_S > 50$, the above condition will be satisfied so long as λ is not very small (for example, if $\lambda \approx 1/(d-1)!$). In this case the value of ϕ at the time when the

CMB perturbations are generated will be determined mainly by the total number of e-foldings of inflation, i.e. $N_T \approx N_S$.

5.4.2 Adiabatic perturbations during D-term inflation

The potential for the flat direction condensate is far from flat, and so if the magnitude of the flat direction condensate is large, it will cause a large deviation from scale-invariance to the adiabatic perturbation. This will impose an upper limit on the amplitude of the flat direction condensate at 50 e-foldings before the end of inflation. If we assume that the flat direction follows a late time attractor trajectory together with the inflaton, then following the analyses in [53], and Kawasaki and Takahashi [54], the flat direction induced adiabatic density perturbation can be estimated from Eq. (125).

For a potential of the form $V = V(S) + V(\phi)$, one obtains (with the help of Eqs. (84,85,115, 126)) [53]

$$\eta = -\frac{M_{\text{P}}^2}{(V'_S + V'_\phi)V} \left[V'_S V''_S + V'_\phi V''_\phi - \frac{2(V'_S + V'_\phi)(V''_S V'^2_S + V''_\phi V'^2_\phi)}{(V'^2_S + V'^2_\phi)} \right] \quad (219)$$

and

$$\epsilon = \frac{M_{\text{P}}^2}{(V'_S + V'_\phi)V} \left[\frac{(V'_S + V'_\phi)(V'^2_S + V'^2_\phi)}{2V} \right]. \quad (220)$$

Particularly, for the case of D-term inflation, if $V'_\phi < V'_S$ and $V''_\phi < V''_S$, we obtain the conventional result, see Eq. (115). Here the isocurvature contribution to the spectral index has been neglected; we will discuss it in the next subsection.

Since the main contribution to the scale-dependence of the perturbations comes from η , let us estimate the deviation from scale-invariance due to the presence of the flat direction condensate. Note that when $V''_\phi > V''_S$, with $V'_\phi \ll V'_S$ and $V_\phi \ll V_S$ still satisfied, we can expand η in order to obtain corrections to the conventional D-term inflation model [53]

$$\eta \approx M_{\text{P}}^2 \frac{V''_S}{V_S} - M_{\text{P}}^2 \frac{V'_\phi V''_\phi}{V_S V'_S}. \quad (221)$$

The condensate scalar induced deviation from the scale invariance in the spectral index is given by

$$\Delta n_\phi \approx -\frac{2V''_\phi V'_\phi M_{\text{P}}^2}{V_S V'_S}. \quad (222)$$

Requiring that $|\Delta n_\phi| < K$, the present CMB observations imply that $n = 1.2 \pm 0.3$; adopting $K < 0.2$ imposes an upper bound on ϕ ,

$$\phi < \phi_c = k_d \left(\frac{K}{\sqrt{N}} \right)^{\frac{1}{4d-7}} g^{\frac{5}{4d-7}} \lambda^{\frac{-4}{4d-7}} \xi^{\frac{8}{4d-7}} M_{\text{P}}^{\frac{4d-15}{4d-7}} , \quad (223)$$

where

$$k_d = \left(\frac{2^{2(d-1)}}{128\pi(d-1)^2(2d-3)} \right)^{\frac{1}{4d-7}} . \quad (224)$$

For the case of minimal $d = 4$ AD baryogenesis, one obtains [53]

$$\phi_c = 0.53 \left(\frac{K}{\sqrt{N}} \right)^{\frac{1}{9}} \left(g^5 \lambda^{-4} \xi^8 M_{\text{P}} \right)^{\frac{1}{9}} \sim 10^{16} \text{ GeV} , \quad (225)$$

while for $d = 6$ AD baryogenesis scenario, one gets

$$\phi_c = 0.77 \left(\frac{K}{\sqrt{N}} \right)^{\frac{1}{17}} \left(g^5 \lambda^{-4} \xi^8 M_{\text{P}}^9 \right)^{\frac{1}{17}} \sim 10^{17} \text{ GeV} . \quad (226)$$

5.4.3 Adiabatic perturbations during F-term inflation

During F-term inflation, the dominant part of the flat direction potential is given by (see Eq. (189), Sect. 4.6.1.)

$$V_{\text{total}}(\phi) \approx \frac{C_I H^2 \phi^2}{2} + V(\phi) , \quad (227)$$

where $V(\phi)$ is the usual part from the non-renormalizable superpotential term. Here we assume that $C_I \approx -\mathcal{O}(1)$. In such a case the local minimum of the flat direction condensate is given by Eq. (190), denoted here by ϕ_m .

Note that if ϕ is close to ϕ_m ($|\delta\phi| \equiv |\phi - \phi_m| \lesssim \phi_m$), then inflation will damp $\delta\phi$ to be close to zero. The equation of motion for perturbations around this local minimum is given by

$$\delta\ddot{\phi} + 3H\delta\dot{\phi} = -kH^2\delta\phi ; k = (2d-4)C_I \gtrsim 1 , \quad (228)$$

which has a solution of the form:

$$\delta\phi = \delta\phi_o e^{\alpha H t} ; \alpha = \frac{1}{2}(-3 + \sqrt{9-4k}) . \quad (229)$$

As long as $Ht \gg 1$, i.e. there are a significant number of e-foldings, the amplitude of the flat direction condensate will be damped to be exponentially close to the minimum of its local minimum.

In general, it is likely that the initial value of ϕ will not be close to ϕ_m . It has been shown that the deviation of the adiabatic perturbation from scale-invariance implies that the value of the potential at $N \approx 50$ cannot be very much larger than ϕ_m [53]. The deviation from scale-invariance due to the flat direction condensate is then

$$\Delta n_\phi = -\frac{2}{\xi} \frac{d\xi}{dN} = -\frac{3V'(\phi)}{V(\phi) + V(S)} \frac{\partial \phi}{\partial N} . \quad (230)$$

For $\phi \gg \phi_m$, the ϕ field will be rapidly oscillating in its potential and the change in the amplitude of ϕ over an e-folding due to damping by expansion will be $\partial \phi / \partial N \sim -\phi$. Requiring that $|\Delta n_\phi| < K$ imposes an upper bound on ϕ

$$\phi \lesssim \left(\frac{Kd}{6(d-1)\lambda^2} \right)^{\frac{1}{2(d-1)}} \sqrt{2} H^{\frac{1}{d-1}} M^{\frac{d-2}{d-1}} . \quad (231)$$

For $d = 4$, one finds [53]

$$\frac{\phi}{\phi_m} \lesssim \frac{0.8}{C_I^{1/4}} \left(\frac{\lambda M_P}{H} \right)^{\frac{1}{6}} , \quad (232)$$

while for $d = 6$

$$\frac{\phi}{\phi_m} \lesssim \frac{0.9}{C_I^{1/8}} \left(\frac{\lambda M_P}{H} \right)^{\frac{1}{20}} , \quad (233)$$

where we have used $K = 0.2$. For typical values of H_I , the scale-invariance of the density perturbations implies that ϕ at $N \approx 50$ e-foldings cannot be much more than an order of magnitude greater than ϕ_m . Since there is no reason for ϕ to be close to this upper limit when $N \approx 50$, it is most likely that ϕ will be close to ϕ_m when the primordial perturbations responsible for large scale structure formation have left the horizon during inflation.

5.4.4 Isocurvature fluctuations in D-term inflation

The isocurvature perturbation in the baryon number arises from the AD scalar if the angular direction is effectively massless, i.e. mass is small compared with H during

and after inflation [52, 53, 54]. The resulting perturbations will be unsuppressed until the baryon number of the Universe is generated. This in turn requires that there are no order H corrections to the supersymmetry-breaking A -terms.

The baryon number from AD baryogenesis is generated at $H \approx m_{susy} \sim 100$ GeV when the A -term can introduce B and CP violation into the coherently oscillating AD scalar. If the phase of the AD scalar relative to the real direction (defined by the A -term) is θ , then the baryon number density given by (see Eq. (253), in Sect. 5.5.)

$$n_B \approx m_{susy} \phi_o^2 \sin 2\theta , \quad (234)$$

where ϕ_o is the amplitude of the coherent oscillations at $H \approx m_{susy}$. One can then obtain fluctuation in the baryon number or an isocurvature perturbation as

$$\frac{\delta n_B}{n_B} = \frac{2\delta\theta}{\tan(2\theta)} = \frac{H}{\pi\phi \tan(2\theta)} , \quad (235)$$

where $\delta\theta \approx (H/2\pi\phi)$ is generated by quantum fluctuations of the AD field at the time when the perturbations cross the horizon. The magnitude of the CMB isocurvature perturbation relative to the adiabatic perturbation can be written as [53, 203, 208], (see Eq. (129), in Sect. 3.4.3.)

$$\alpha = \left| \frac{\delta_\gamma^i}{\delta_\gamma^a} \right| = \frac{\omega}{3} \left(\frac{2M^2 V'(S)}{V(S) \tan(2\theta) \phi} \right) , \quad (236)$$

where $V(S)$ is the inflaton potential (see Sect. 5.4.1). For purely baryonic isocurvature perturbations $\omega = \Omega_B/\Omega_m$, where Ω_B is the baryon density and Ω_m is the total matter density. For the case of D-term inflation one obtains [53]

$$\alpha = \frac{1}{6\pi} \frac{g\omega M}{\phi N^{1/2} \tan(2\theta)} , \quad (237)$$

where $N \approx 50$.

Requiring that the deviations from the spectral index due to the AD scalar are acceptably small, for $d = 4$, one finds

$$\alpha > \alpha_c = \frac{3.3\omega(g\lambda)^{4/9}}{K^{1/9} \tan(2\theta)} , \quad (238)$$

and for $d = 6$

$$\alpha > \alpha_c = \frac{0.18\omega(g^3\lambda)^{4/17}}{K^{1/17}\tan(2\theta)} . \quad (239)$$

The range of Ω_B allowed by nucleosynthesis is $0.006 \lesssim \Omega_B \lesssim 0.036$ (for $0.6 \lesssim h \lesssim 0.87$) [3]. For $\Omega_m = 0.4$, $K = 0.2$, and for $d = 4$, we obtain

$$\alpha_c = (0.06 - 0.36) \frac{(g\lambda)^{4/9}}{\tan(2\theta)} , \quad (240)$$

and for $d = 6$

$$\alpha_c = (3.0 \times 10^{-3} - 0.018) \frac{(g^3\lambda)^{4/17}}{\tan(2\theta)} . \quad (241)$$

(The lower limits should be multiplied by 0.4 for the case $\Omega_m = 1$.) If, for example, $g \sim \lambda \sim 0.1$ and $\tan(2\theta) \lesssim 1$, one would obtain a lower bound $\alpha \gtrsim 10^{-2}$ for $d = 4$ and $\alpha \gtrsim 10^{-3}$ for $d = 6$. Such small isocurvature contamination could be detectable in future CMB experiments.

Present CMB and large-scale structure observations require that $\alpha \lesssim 0.1$ [203, 208]. COBE normalization combined with the value of σ_8 (the rms of the density field on a scale of 8 Mpc) as obtained from X-ray observations of the local cluster together with the shape parameter $\Gamma \approx \Omega_m h = 0.25 \pm 0.05$ [333] from galaxy surveys, which is also consistent with the recent observations of high-redshift supernovae [20, 21]) yields the limit $\alpha \lesssim 0.07$. The limit may however rely too much on COBE normalization, which is just one experimental result among many.

Future CMB observations by MAP will be able to probe down to $\alpha \approx 0.1$, while PLANCK (with CMB polarization measurements) should be able to see isocurvature perturbations as small as 0.04 [204] (see also [205]). For the case of minimal ($d = 4$) AD baryogenesis, there is a good chance that PLANCK will be able to observe isocurvature perturbations at least if inflation is driven by D-term. For higher dimension AD baryogenesis ($d \geq 6$) the situation is less certain.

All this assumes that ϕ can take any value. This is true if $\phi_i < \phi_*$, in which case ϕ remains at its initial value ϕ_i . We have seen that the dynamics of the flat direction during D-term inflation implies that if $\phi_i > \phi_*$ then ϕ will equal ϕ_* at $N \approx 50$. In this case we can fix the magnitude of the isocurvature perturbation. For $d = 4$, $N \approx 50$

and $\Omega_m = 0.4$ it is given by [53]

$$\alpha = \alpha_* \approx (0.17 - 1.03) \left(\frac{N_S}{50} \right)^{1/4} \frac{(g\lambda)^{1/2}}{\tan(2\theta)} . \quad (242)$$

(For $\Omega_m = 1$ this should be multiplied by 0.4.) For $d = 6$ and $\Omega_m = 0.4$,

$$\alpha = \alpha_* \approx (4.4 \times 10^{-3} - 2.6 \times 10^{-2}) \left(\frac{N_S}{50} \right)^{1/8} \frac{g^{3/4} \lambda^{1/4}}{\tan(2\theta)} . \quad (243)$$

If $g, \lambda \gtrsim 0.1$ then for the $d = 4$ case one expects $\alpha_* \approx 0.01 - 0.1$. For the $d = 6$ case the isocurvature perturbation might just be at the observable level.

It is important that one can fix the isocurvature perturbation to be not much larger than the lower bound coming from adiabatic perturbations. This is because there is typically a very small range of values of ϕ over which the isocurvature perturbation is less than the present observational limit, $\alpha \lesssim 0.1$, but larger than the adiabatic perturbation lower bound, $\alpha \gtrsim 0.01$ for $d = 4$.

5.4.5 Isocurvature fluctuations in F-term inflation

If the flat direction condensate is stuck in a local minimum $\phi \approx \phi_m$ given by Eq. (190), the isocurvature perturbation is given by [52, 53]

$$\alpha \approx \frac{2\omega}{3} \frac{H}{\tan(2\theta) \delta_\rho \phi_m} , \quad (244)$$

where $\delta_\rho = 3\delta T/T \approx 3 \times 10^{-5}$ is the density perturbation. Given H , d , and the value of ϕ_m , the magnitude of the isocurvature perturbation is essentially fixed. For $d = 4$ and $\Omega_m = 0.4$, the isocurvature perturbation has been found to be

$$\alpha = (3.1 - 18.6) \times 10^2 \frac{\lambda^{1/2}}{C_I^{1/4} \tan(2\theta)} \left(\frac{H_I}{M_P} \right)^{1/2} \quad (245)$$

while for $d = 6$

$$\alpha = (2.9 - 17.4) \times 10^2 \frac{\lambda^{1/4}}{C_I^{1/8} \tan(2\theta)} \left(\frac{H_I}{M_P} \right)^{3/4} . \quad (246)$$

If we require that $\alpha \lesssim 0.1$ we find the upper bounds $H_I/M_P \lesssim 10^{-7}/\lambda$ (for $d = 4$) and $H_I/M_P \lesssim 10^{-5}/\lambda^{1/3}$ (for $d = 6$). For typical values of H the isocurvature perturbation

in the F-term inflation can be close to the present observational limits. In [54], however, it was pointed out that there would be negligible isocurvature perturbations produced from the MSSM flat directions, and the present observations would not be able place any independent constraint upon the initial amplitude of the flat directions.

5.4.6 MSSM flat directions as a source for curvature perturbations

In a very recent development, a new paradigm has been laid, where adiabatic density perturbations were generated from the decay of the pure isocurvature perturbations. Though it was first suggested in [319], but it was implemented recently in a pre-Big-Bang scenario [320], where the axion field which generates isocurvature perturbations decays late in the Universe.

In order to create pure adiabatic density perturbations it is important that the field, known as curvaton [321], is subdominant during inflation, but becomes dominant during the late phase of the Universe especially when it is decaying. The curvaton field σ generates isocurvature perturbations during inflation, assuming that the perturbations generated from the inflaton field can be negligible, and the curvaton mass $m_\sigma^2 \approx V_{\sigma\sigma} \ll H_{inf}^2$. In this limiting case the power spectrum for the curvaton will be equivalent to a massless scalar field during inflation; $\mathcal{P}_\sigma = H_*^2/4\pi^2$, where $*$ denotes the epoch when perturbations are crossing the horizon $k = a_*H_*$. The curvaton field follows its trajectory after the end of inflation, and when $H \sim m_\sigma$, the curvaton oscillates and eventually decays through its coupling to the SM relativistic degrees of freedom. While oscillating it produces density contrast $\delta_\sigma = 2\delta\sigma/\sigma$, assuming that $H_* < \sigma_*$ [321], and the perturbation spectrum is given by

$$\mathcal{P}_{\delta_\sigma}^{1/2} = 2 \frac{\mathcal{P}_\sigma^{1/2}}{\sigma} = \frac{H_*}{\pi\sigma_*}. \quad (247)$$

When the curvaton decays it converts all its isocurvature perturbations to the adiabatic ones by following that, before decaying the relativistic degrees of freedom due to the inflaton decay products gives rise to density perturbations in the radiation as $\zeta_r = (1/4)\delta_r$, and $\zeta_\sigma = (1/3)\delta_\sigma$. With these results the curvature perturbations is

given by [321]

$$\zeta = \frac{4\rho_r\zeta_r + 3\rho_\sigma\zeta_\sigma}{4\rho_r + 4\rho_\sigma} \approx \frac{1}{3}\delta_\sigma, \quad (248)$$

supposing that before decaying ζ_r is negligible. If the curvaton decay products do not dominate the Universe, then there will adiabatic and isocurvature perturbations both [321, 322].

In [323], the authors have pointed out that the MSSM flat directions which are lifted by the non-renormalizable operators such as $d = 7, 9$ are the best candidate for the curvaton. The flat directions $d = 7$, $LL\bar{d}\bar{d}\bar{d}$ (lifted by $H_u LLLddd$), and $d = 9$, $Q\bar{u}Q\bar{u}Q\bar{u}\bar{e}$ (lifted by $Q\bar{u}Q\bar{u}Q\bar{u}H_d\bar{e}\bar{e}$), which are lifted by superpotential: $W \sim \sigma^{d-1}\psi/M^{d-3}$. Note that ψ is the superfield other than the curvaton, does not produce any A -term in the potential, since $\langle\psi\rangle = 0$, and therefore does not give rise to any $U(1)$ violating terms in the flat direction potential. It was shown in [323], that these flat directions can dominate the energy density of the Universe and while decaying the squarks and sleptons can directly decay in the MSSM relativistic degrees of freedom. Therefore the virtue of this scenario is that the MSSM flat direction is solely responsible for reheating the Universe, barring any need for speculation from the inflaton coupling to the SM fields. Inflation was supposed to happen in the hidden sector of the theory, which does not necessarily couple to the SM sector.

5.5 Baryon number asymmetry

In both D-and F-term inflation the inflaton and other scalar fields begin to oscillate coherently about the minimum of their respective potential after the end of inflation, and the post-inflationary evolution of the flat direction condensate is no exception. If C_I and C are positive in Eqs. (189) and (192), the corresponding vevs are either given by Eqs. (190) or (193). In fact, in D-term inflation models for $|C|$ less than about 0.5, it is possible to have a positive H^2 correction and still generate the observed baryon asymmetry as shown by McDonald [324]. In addition, there has been attempts for AD baryogenesis in F-term inflation, basing on the low energy effective action of the heterotic string theory, where inflation is driven by T -moduli [334]. Flat directions

beyond MSSM involving a triplet Higgs has also been considered [335]. Here we will mainly concentrate on the negative H^2 correction only.

After inflation, $\langle\phi\rangle$ initially continues to track the instantaneous local minimum of the scalar potential, which can be derived by replacing H_I with $H(t)$ in Eq.(190). Once $H \simeq m_0$, the low-energy soft terms take over. The condensate mass squared turns positive, and since the phase of $\langle\phi\rangle$ differs from the phase of A , $\langle\phi\rangle$ starts to change non-adiabatically.

In an absence of H corrections to the A -terms, the initial phase θ of the AD field relative to the real direction is random and so typically ≈ 1 . As a result $\langle\phi\rangle$ starts a spiral motion in the complex plane (see Figs. (2), and forthcoming discussion on the condensate trajectory), which leads to a generation of a net baryon and/or lepton asymmetry [45, 46]⁷.

The baryon number density is related to the AD field as

$$n_{B,L} = \beta i(\dot{\phi}^\dagger \phi - \phi^\dagger \dot{\phi}), \quad (249)$$

where β is corresponding baryon and/or lepton charge of the AD field. The equations of motion for the AD field are given by

$$\ddot{\phi} + 3H\dot{\phi} + \frac{\partial V(\phi)}{\partial \phi^*} = 0. \quad (250)$$

The above two equations lead to

$$\begin{aligned} \dot{n}_{B,L} + 3Hn_{B,L} &= 2\beta \text{Im} \left[\frac{\partial V(\phi)}{\partial \phi^*} \phi \right], \\ &= 2\beta \frac{m_\phi}{dM^{d-3}} \text{Im}(a\phi^d). \end{aligned} \quad (251)$$

By integrating Eq. (251), we obtain the baryon and/or lepton number as

$$a^3(t)n_{B,L}(t) = 2\beta|a|\frac{m_\phi}{M^{d-3}} \int^t a^3(t')|\phi(t')|^d \sin(\theta) dt', \quad (252)$$

Note that a introduces an extra CP phase which we may parameterize as $\sin(\delta)$. After a few expansion times, the amplitude of the oscillations will become damped by the expansion of the Universe and the A -term, which is proportional to a large power of

⁷There have been attempts for AD baryogenesis in local domains, see [336]

ϕ , will become gradually negligible. The net baryon and/or lepton asymmetry is then given by [46]

$$\begin{aligned} n_{B,L}(t_{osc}) &= \beta \frac{2(d-2)}{3(d-3)} m_\phi \phi_0^2 \sin 2\theta \sin \delta, \\ &\approx \beta \frac{2(d-2)}{3(d-3)} m_\phi \left(m_\phi M^{d-3}\right)^{2/(d-2)} \sin 2\theta \sin \delta, \end{aligned} \quad (253)$$

where $\sin \delta \sim \sin 2\theta \approx \mathcal{O}(1)$.

When the inflaton decay products have completely thermalized with a reheat temperature T_{rh} , the baryon and/or lepton asymmetry is given by [337]

$$\begin{aligned} \frac{n_{B,L}}{s} &= \frac{1}{4} \frac{T_{rh}}{M_{\text{P}}^2 H(t_{osc})^2} n_{B,L}(t_{osc}), \\ &= \frac{d-2}{6(d-3)} \beta \frac{T_{rh}}{M_{\text{P}}^2 m_\phi} \left(m_\phi M^{d-3}\right)^{2/(d-2)} \sin 2\theta \sin \delta, \end{aligned} \quad (254)$$

where we have used $H(t_{osc}) \approx m_\phi$, and s is the entropy density of the Universe at the time of reheating. For $d = 4$, the baryon-to-entropy ratio turns out to be [337]

$$\frac{n_{B,L}}{s} \approx 1 \times 10^{-10} \times \beta \left(\frac{m_{3/2}}{m_\phi}\right) \left(\frac{M}{M_{\text{P}}}\right) \left(\frac{T_{rh}}{10^9 \text{ GeV}}\right), \quad (255)$$

and for $d = 6$

$$\frac{n_{B,L}}{s} \approx 5 \times 10^{-10} \times \beta \left(\frac{m_{3/2}}{m_\phi}\right) \left(\frac{1 \text{ TeV}}{m_\phi}\right)^{1/2} \left(\frac{M}{M_{\text{P}}}\right)^{3/2} \left(\frac{T_{rh}}{100 \text{ GeV}}\right), \quad (256)$$

where we have taken the net CP phase to be $\sim \mathcal{O}(1)$. The asymmetry remains frozen unless there is additional entropy production afterwards. Note that for $d = 4$, the required reheat temperature of the Universe is below the gravitino overproduction bound (see Sect. 3.6.2). For higher dimensional non-renormalizable operators, a low reheat temperature is favorable, which is indeed a good news.

In this regard low scale inflation, which guarantees a low reheat temperature, has been given some consideration [338] (see also [339] where AD baryogenesis after a brief period of thermal inflation, required to solving the cosmological moduli problem, has been discussed). Although, in [340], it was pointed out that in gauge mediated supersymmetry breaking it is hard to reconcile AD baryogenesis with a moduli problem.

Among the host of MSSM flat directions which are lifted by non-renormalizable operator and listed in Table 1, the LH_u flat direction carrying the lepton number

is the candidate for producing lepton asymmetry in the Universe (there has been some earlier attempts of direct baryogenesis via $\bar{u}d\bar{d}$ directions, see e.g. [341]). The lepton asymmetry calculated above in Eqs. (255,256) can be transformed into baryon number asymmetry via sphalerons $n_B/s = (8/23)n_L/s$. AD leptogenesis has important implications in neutrino physics also, because in the MSSM, the LH_u direction is lifted by the $d = 4$ non-renormalizable operator which also gives rise to neutrino masses [46]:

$$W = \frac{1}{2M_i} (L_i H_u)^2 = \frac{m_{\nu_i}}{2\langle H_u \rangle^2} (L_i H_u)^2, \quad (257)$$

where we have assumed the see-saw relation $m_{\nu_i} = \langle H_u \rangle^2 / M_i$ with diagonal entries for the neutrinos ν_i , $i = 1, 2, 3$. The final n_B/s can be related to the lightest neutrino mass since the flat direction moves furthest along the eigenvector of $L_i L_j$ which corresponds to the smallest eigenvalue of the neutrino mass matrix [46].

$$\frac{n_L}{s} \approx 1 \times 10^{-10} \times \beta \left(\frac{m_{3/2}}{m_\phi} \right) \left(\frac{T_{rh}}{10^8 \text{ GeV}} \right) \left(\frac{10^{-6} \text{ eV}}{m_{\nu_l}} \right), \quad (258)$$

where m_{ν_l} denotes the lightest neutrino.

5.6 Thermal effects

In our discussion on the baryon/lepton asymmetry we have tacitly assumed that the asymmetry has been generated before the Universe has thermalized and reheated. This might not be the case if there were light degrees of freedom which have thermalized with an instantaneous plasma temperature $T_{inst} \leq (H\Gamma_d M_P)^{1/4}$ before the inflaton has decayed. We remind that the bulk of energy density is still in the form of inflaton oscillations, and only a fraction of the energy density has gone into these light Standard Model degrees of freedom. If the MSSM flat direction couples with this thermal bath, there arises a modification in the flat direction potential [309, 310, 342].

5.6.1 Thermal corrections to the flat direction potential

Besides the D-term couplings of the form $g^2 \phi^2 \alpha^2$, where α is some field with gauge interactions, there are also F-term Yukawa couplings to fields χ which result in a term $h^2 |\phi|^2 |\chi|^2$ in the flat direction potential. χ and α obtain large masses due to the

flat direction vev, and therefore do not feel the effect of temperature if the condensate amplitude ϕ is large. As pointed out by Allahverdi, Campbell and Ellis [309], the back-reaction effect induces a mass-squared term $h^2 T^2$ for the flat direction. If this exceeds the negative Hubble induced mass squared term, the flat direction oscillations starts earlier than otherwise expected. In order for thermal correction to play a significant role the couplings h, g must have intermediate strength. Otherwise, a large coupling would induce a large vev dependent mass for α , which would prevent its thermal excitations, and a very small coupling would not have significant thermal backreaction at all.

For the inflationary scale $H_I \sim 10^{13}$ GeV and $M = M_{GUT}$, it has been found [309] that a generic MSSM flat direction with a Yukawa coupling $h \sim 10^{-2}$ starts oscillating at $H \gg 10^2$ GeV for $4 \leq d \leq 8$. Since thermal effects induce early oscillations, baryon asymmetry is also produced much earlier, which could have interesting consequences.

Another thermal effect has been discussed by Anisimov and Dine [310, 342]. All the flat directions which are lifted at large d give rise to a large mass for α , and consequently one should account for their effect by integrating out the heavy modes. This would result in terms like

$$\frac{A}{16\pi^2} F_{\mu\nu}^2 \ln \frac{|\phi|^2}{M^2}. \quad (259)$$

In particular for the flat direction LH_u the effective potential thus obtained has the form [310]

$$V_{eff} = \alpha_s^2(T) a_g T^4 \ln \frac{|\phi|^2}{M^2}, \quad (260)$$

where $\alpha_s \equiv g_s^2/4\pi$, and $a_g = \frac{3N_g}{288} \left(5\frac{N_f}{4} + 7\frac{N}{2} \right)$ includes leading order contribution of the gluons, gluinos, and quarks to the free energy for a non-abelian group $SU(N)$. The oscillations in the flat direction are induced when $H_{osc}^2 = \partial^2 V_{eff} / \partial |\phi|^2 = \alpha_s^2 a_g T^4 / |\phi|^2$ and one can check that [309] for $d = 4$, the thermal mass correction $\sim h^2 T^2$ wins over the logarithmic counterpart Eq. (260), but for $d = 5$ and/or 6, the logarithmic correction dominates and the oscillations start earlier than otherwise one would have expected.

There could also be a thermal enhancement of the A -term [309], which can arise

from the cross terms

$$W \supset h\phi\alpha\alpha + \lambda_d \frac{\phi^d}{dM^{d-3}}. \quad (261)$$

However a symmetry forbids such enhancement [342], although the situation might change if one adds more terms in the superpotential such as [342].

$$W = \frac{1}{M} \left(aI + b \frac{I^2}{M} \right) \frac{\phi^d}{M^{d-3}}. \quad (262)$$

5.6.2 Thermal evaporation of the flat direction

It has been argued that in general the flat direction condensate decays as a result of scattering with the thermalized decay products of the inflaton [46]. Usually the scattering interactions preserve B and L , and therefore the previously produced baryon and/or lepton asymmetry remains unchanged. In [310] it was assumed that thermalized fermions scatter with the condensate with a rate

$$\Gamma_{scatt} \simeq yg^2T. \quad (263)$$

where yT corresponds to the mass of the condensate. A complete evaporation was found to be avoided only after reheating if $T_{rh} \leq (yg^2)^{2/3} M_{\text{P}}^{5/6} H^{1/6}$, which is usually satisfied for a reasonable range of reheat temperatures and Yukawa couplings.

5.7 Baryosynthesis and neutrino mass

As discussed in Sect. 5.4., the lepton asymmetry via LH_u direction leads to a prediction on the lightest neutrino mass. It is however pertinent to include also the finite temperature effects [343]. At finite T , the flat direction potential for LH_u direction can be written as [343]

$$\begin{aligned} V_{total} = & \left(m_\phi^2 - C_I H^2 + \sum_{f_k |\phi| < T} c_k f_k^2 T^2 \right) |\phi|^2 + \frac{m_{3/2}}{8M} (a_m \phi^4 + \text{h.c.}) + \frac{H}{8M} (a_H \phi^4 + \text{h.c.}) \\ & + a_g \alpha_s^2 T^4 \ln \left(\frac{|\phi|^2}{T^2} \right) + \frac{|\phi|^6}{4M^2}, \end{aligned} \quad (264)$$

where c_k are real positive constants and couplings $f_k = 1 - 10^{-5}$ in MSSM [337]. The mismatch in phases between a_m and a_H leads to the helical motion of the flat direction.

Once the inflaton decay products generate a thermal plasma with a temperature $T = (T_{rh} M_P H_I)^{1/4}$, thermal corrections take over the Hubble induced term

$$H^2 \leq m_\phi^2 + \sum_{f_k |\phi| < T} c_k f_k^2 T^2 + a_g \alpha_s^2(T) \frac{T^4}{|\phi|^2}. \quad (265)$$

The flat direction starts to oscillate when [343]

$$H_{osc} \approx \max \left[m_\phi, H_i, \alpha_s T_{rh} \left(\frac{a_g M_P}{M} \right)^{1/2} \right], \quad (266)$$

where H_i is given by [337, 343]

$$H_i \approx \min \left\{ \frac{1}{f_i^4} \frac{M_P T_{rh}^2}{M^2}, (c_i^2 f_i^4 M_P T_{rh}^2)^{1/3} \right\}. \quad (267)$$

The lepton asymmetry is then given by [343]

$$a^3(t) n_L(t) \approx \frac{m_{3/2}}{2M} \int^t dt' a^3(t') \mathcal{I}m(a_m \phi^4), \quad (268)$$

The right hand side of Eq. (268) initially increases until $H \approx H_{osc}$, after which the integrand is rapidly damped because $a^3 \phi^4 \sim t^{-n}$ for $n > 1$. The final lepton asymmetry is determined approximately by the configuration at the time when the oscillations commence [343]

$$n_L = \frac{m_{3/2}}{2M} \mathcal{I}m(a_m \phi^4) t \Big|_{H=H_{osc}} = \frac{1}{3} m_{3/2} M H_{osc} \delta_{eff}, \quad (269)$$

where $\delta_{eff} = \sin(4\arg(\phi) + \arg(a_m))$ is the net CP phase. The final baryon to entropy ratio turns out to be [343]

$$\frac{n_B}{s} = 10^{-11} \delta_{eff} \times \left(\frac{m_\nu l}{10^{-8} \text{eV}} \right)^{-3/2} \left(\frac{m_{3/2}}{1 \text{ TeV}} \right). \quad (270)$$

Note that the final expression obtained does not depend upon the reheat temperature T_{rh} , mainly due to the fact that the H_{osc} is determined by thermal correction $\sim T^4 \ln(|\phi|^2)$. This is however true only for $10^8 \text{ GeV} \leq T_{rh} \leq 10^{12} \text{ GeV}$ as pointed out by Fujii, Hamaguchi, and Yanagida in [343]. For $10^5 \text{ GeV} \leq T_{rh} \leq 10^8 \text{ GeV}$, the dependence on reheat temperature appears as $n_B/s \propto T_{rh}^{1/3}$, because then the thermal mass term $\sim T^2 |\phi|^2$ dominates.

It is possible [343] to obtain the right amount of baryon asymmetry with the lightest neutrino mass $m_{\nu l} \simeq (0.1 - 3) \times 10^{-9} \text{ eV}$ and with a CP phase $\delta_{eff} \simeq (0.1 - 1)$ for a fairly wide range of reheat temperature $10^5 \text{ GeV} \leq T_{rh} \leq 10^{12} \text{ GeV}$.




Figure 2: Affleck-Dine condensate formation with $x = \phi_1$ and $y = \phi_2$, for (a) gravity mediated case with $d = 4$ (solid) and $d = 6$ (dashed), and (b) gauge mediated case with $d = 4$, $m_\phi = 1$ TeV (solid) and $m_\phi = 10$ TeV (dashed) with the initial condition $\theta_i = -\pi/10$, from [344].

5.8 Trajectory of a flat direction

Let us now turn our attention to the dynamical evolution of the flat direction after the end of inflation. Here we assume that the flat direction is tracking its minimum which is determined by Eq. (190). The trajectory of the flat direction depends upon the potentials Eqs. (200) and (204). Here we sketch the main differences between the gravity and gauge mediated cases.

Jokinen [344] has studied numerically the trajectories of the flat direction condensate in gravity and gauge mediated cases, following Eqs. (200) and (204). The rotation of the condensate depends on the low energy supersymmetry breaking mass terms. The classical motion for the condensate $\phi = (\phi_1 + i\phi_2)/\sqrt{2}$ is illustrated in Fig. (2). In the gravity mediated case, Fig. (2a), we see that the orbit is a spiraling ellipse and in the gauge mediated case, Fig. (2b), a precessing trefoil. From Fig. (2), one can see that there is a twist on the orbit much before the rotation starts properly. This is the time of the phase transition, when the condensate ϕ starts to rotate in the pit of the symmetry breaking minimum. The rotation begins when the symmetry breaking minimum is the vacuum, and ends when it has become a false vacuum, and twists when the false vacuum has completely vanished forming a kink on the orbit. It is possible to produce a condensate through a second order phase transition, but the charge in that case will be small. It should also be noted that in the gravity mediated

case condensate formation starts when $C_I H^2 \sim m_{3/2}^2$ for all values of d , A and a . In the gauge mediated case the condensate formation starts at $C_I H^2 \sim m_\phi^4/|\phi|^2$, so that the formation happens earlier if the condensate mass m_ϕ is increased, as can be seen from the different positions of the kink in Fig. (2b).

5.9 Instability of the coherent condensate

5.9.1 Negative pressure

The effective equation of state of a coherent scalar field oscillating in a potential $U(\phi)$ with a frequency which is large compared with H is obtained by averaging $p/\rho = (|\dot{\phi}|^2/\rho) - 1$ over one oscillation cycle T . The result is [230]

$$p = (\gamma - 1)\rho , \quad (271)$$

where

$$\gamma = \frac{2}{T} \int_0^T \left(1 - \frac{U(\phi)}{\rho} \right) dt . \quad (272)$$

For the case $U \sim m^2 \phi^2$, one finds $\gamma = 1$, so that one effectively obtains the usual case of pressureless, non-relativistic cold matter.

When the motion of the condensate field is not simply oscillatory, such as in the case for the condensate trajectory, one can generalize Eq. (272) by integrating over the orbit c of the AD field. In that case

$$\gamma = \frac{2 \int_c d|\phi| (1 - U(\phi)/\rho)^{\frac{1}{2}}}{\int_c d|\phi| (1 - U(\phi)/\rho)^{-\frac{1}{2}}} . \quad (273)$$

In practice the orbits are nearly elliptical. Then the arc length is given by

$$d|\phi| = \frac{d\phi_1}{\sqrt{2}} \sqrt{1 + \frac{B^2 \phi_1^2}{A^4 (1 - \phi_1^2/A^2)}} , \quad (274)$$

where A and $B \leq A$ are respectively the semi-major and the semi-minor axis of the ellipse, and $\phi_1 = \text{Re}\phi/\sqrt{2}$. For a circular orbit $B = A$, whereas for pure oscillation (no charge in the condensate) $B = 0$.

It is therefore obvious that small corrections to a harmonic potential of a coherent condensate can easily generate a pressure. As we have seen, in the gravity mediated




Figure 3: Pressure-to-energy density ratio, $p/\rho \equiv w = \gamma - 1$, in the gravity mediated case vs. (a) time in logarithmic units for $d = 4$, (b) different initial conditions for $d = 4, 6$; (c) ellipticity $\varepsilon = B/A$ vs. initial conditions for $d = 4$ (thin lines), $d = 6$ (thick lines), D-term (solid), F-term (dashed) with $K = -0.01$ and $t = 100m_{3/2}^{-1}$. In (b) w is shown at $t = 300m_{3/2}^{-1}$ with dotted lines for the $d = 4$ D-term case, from [344].

case quantum corrections typically modify the flat direction mass terms by

$$U(\phi) = \frac{1}{2}m_\phi^2\phi^2 + Km_\phi^2\phi^2 \log\left(\frac{\phi^2}{\mu^2}\right) + \dots, \quad (275)$$

where K is some constant. If one writes

$$U(\phi) = \frac{1}{2}m_\phi^2\phi^2 \left(\frac{\phi^2}{\mu^2}\right)^x \quad (276)$$

one finds that

$$\gamma = \frac{1+x}{1+\frac{x}{2}}, \quad p = \frac{x}{2+x}. \quad (277)$$

In the case of the logarithmic potential $x \simeq 2K$. There arises a negative pressure $p = K\rho$ whenever $K < 0$ or whenever x is small and negative. This is a sign of an instability of the condensate under arbitrarily small perturbations.

This is exactly the situation one finds in the MSSM with flat directions. The effective mass $m_{\text{eff}}^2(\phi) \equiv dU/d\phi^2$ decreases for a range in ϕ , albeit for different reasons, both for gravity mediated and gauge mediated supersymmetry breaking. According to Eq. (273) this results in a negative pressure, which has been computed numerically by Jokinen [344]. The results are shown in Figs. (3) and (4).

The pressure-to-energy density ratio; $w = \gamma - 1$ for the gravity mediated case is plotted in Fig. (3a), which shows that there are time-dependent oscillations in pressure [344]. The average pressure is slightly on the negative side. The average value of w is




Figure 4: Pressure-to-energy density ratio, w , in the gauge mediated D-term case (without Hubble induced A -term) vs. time in logarithmic units for (a) $d = 4$ and (b) $d = 6$; (c) ellipticity of the orbit where $d = 4$ (thin lines) and $d = 6$ (thick lines). The scalar masses $m_\phi = 1, 10, 100$ TeV are denoted respectively with solid, dotted and dashed lines, from [344].

shown in Fig. (3b) at $t \sim 100m_{3/2}^{-1}$ for a few different initial conditions. In Fig. (3c), the ellipticity of the orbit, $\varepsilon = B/A$, is plotted to show that w is more negative if ε is small. It should be noted that w achieves values which are more negative than the absolute lower bound coming from pure oscillation.

A similar analysis has been made for the gauge mediated supersymmetry breaking case. The results have striking similarities [344]. In Figs. (4a) and (4b), the time development of the pressure-to-energy density ratio, w , for $d = 4$ and $d = 6$ has been depicted. One can see that the pressure is always negative. The calculation of average pressure is even more involved than in the gravity mediated case, since the oscillation frequency becomes very large. In Fig. (4c), the ellipticity of the orbit is shown as a function of different initial conditions. Jokinen [344] has pointed it out that quite generically $\varepsilon \lesssim 0.1$.

5.9.2 Growth of perturbations in the AD condensate

As a result of internal negative condensate pressure the quantum fluctuations in the scalar condensate grow according to [9]

$$\ddot{\delta}_{\mathbf{k}} = -K\mathbf{k}^2\delta_{\mathbf{k}}. \quad (278)$$




Figure 5: Energy-to-charge ratio, x , in the gravity mediated case vs. (a) time in logarithmic units for $d = 4, 6$; (b) the D-term case; (c) the F-term (with Hubble induced A -term) case with $d = 4, 5, 6, 7$ (solid, dash-dot, dashed and dotted lines), $K = -0.01$ and $t = 100m_{3/2}^{-1}$, from [344].

If $K < 0$, quantum fluctuations of the condensate field at the scale $\lambda = 2\pi/|\mathbf{k}|$ will grow exponentially in time as

$$\delta\phi_{\mathbf{k}}(t) = \delta\phi(0)\exp(-K\mathbf{k}^2t) . \quad (279)$$

In reality the onset of non-linearity sets the scale at which the spatial coherence of the condensate can no longer be maintained and the condensate fragments. For the AD condensate the initial perturbation originates from inflation. Note that since the AD condensate carries a global charge, due to charge conservation the energy-to-charge ratio changes as the the condensate fragments.

The energy-to-charge ratio has been estimated numerically for both gravity and gauge mediated cases by Jokinen [344]. The time evolution of the energy-to-charge ratio x is shown in Fig. 5, where x is also plotted for various initial phases in F-and D-term inflation models. For the gauge mediation case the plots are quite different from the gravity mediated case, see Fig. (6).

5.9.3 The true ground state

Under the negative pressure the homogeneous AD condensate fragments and forms lumps. The question then is, what is the true ground state? The answer is, a non-topological soliton with a fixed charge, called the Q -ball [345, 346, 47], which in general is made up of a complex scalar field with a global $U(1)$ symmetry, for which the




Figure 6: Energy-to-charge ratio, x , in the gauge mediated case vs. (a) time in logarithmic units of time $d = 4$, (b) $d = 4$ and (c) $d = 6$ with D-term (thin lines, without Hubble induced A-term) and F-term (thick lines, with Hubble induced A-term) and $m_\phi = 1, 10, 100$ TeV (solid, dashed, dotted lines) at $t = 4 \cdot 10^5 m_\phi^{-1}, 10^5 m_\phi^{-1}, 4 \cdot 10^4 m_\phi^{-1}$ ($d = 4$) and $t = 4 \cdot 10^9, 10^9, 4 \cdot 10^8 m_\phi^{-1}$ ($d = 6$), from [344].

Lagrangian is

$$\mathcal{L} = \partial_\mu \phi \partial^\mu \phi^* - U(\phi \phi^*) . \quad (280)$$

When supplemented by the CP violating terms, this is the Lagrangian for the MSSM flat directions.

The conserved current is $j^\mu = i(\phi \partial^\mu \phi^* - \phi^* \partial^\mu \phi)$, and the conserved charge, and energy are given by

$$Q = \int d^3x j^0 , \quad (281)$$

$$E = \int d^3x [|\dot{\phi}|^2 + |\nabla \phi|^2 + U(\phi \phi^*)] \quad (282)$$

If the charge is kept fixed, the state of lowest energy is found by minimizing [47]

$$E_\omega = E - \omega(Q - i \int d^3x j^0) , \quad (283)$$

with respect to variations in ϕ and the Lagrange multiplier ω . From $\delta_\phi E_\omega = 0$, it follows that $\dot{\phi} - i\omega\phi = 0$, so that we may write

$$\phi(t, \mathbf{x}) = e^{i\omega t} \varphi(\mathbf{x}) \quad (284)$$

where φ may be chosen real by virtue of $U(1)$ invariance. The charge and energy of such a configuration read

$$Q = 2\omega \int d^3x \varphi^2 , \quad (285)$$

$$E = \int d^3x [(\nabla\varphi)^2 + U(\varphi^2) + \omega^2\varphi^2] , \quad (286)$$

and one has to minimize $E_\omega = \int d^3x[(\nabla\varphi)^2 + \hat{U}_\omega(\varphi^2)] + \omega Q$ where

$$\hat{U}_\omega(\varphi^2) = U(\varphi^2) - \omega^2\varphi^2 . \quad (287)$$

To find a localized configuration that vanishes at spatial infinity one may make use of the spherical rearrangement theorem, which implies that E_ω is minimized by $\varphi(\mathbf{x})$ which is spherically symmetric and monotonically decreasing. This is equivalent to solving the equation of motion

$$\frac{d^2\varphi}{dr^2} + \frac{2}{r} \frac{d\varphi}{dr} - \varphi \frac{dU_\omega}{d\varphi^2} = 0 . \quad (288)$$

If by convention we set the globally symmetric minimum to $\varphi = 0$ with $U(0) = 0$, one can then show that a non-trivial solution to Eq. (288) is obtained whenever $U(\varphi^2)/\varphi^2$ has a minimum at $\varphi \neq 0$, i.e. $U(\varphi^2)$ grows more slowly than $m_\phi^2\varphi^2$ over some range. We will discuss Q -balls in detail in the next Sects. 6 and 7.

5.10 Numerical studies of fragmentation

Although the homogeneous AD condensate is not the ground state, it is not obvious that the ground state should always be reached within cosmic time scales. It is then essential to study the dynamical evolution of the AD condensate. Since the Q -ball formation is inherently a non-linear phenomenon, analyzing small perturbations is not sufficient to determine the full dynamical evolution of the AD condensate. One can nevertheless gather some information about the gross features of the condensate fragmentation by perturbative considerations alone.

5.10.1 Perturbation theory

Negative pressure is equivalent to an attractive force between the condensate quanta which induces a growing mode in spatial perturbations. A linearized description of the evolution of perturbations has been given by Kusenko and Shaposhnikov in [151], and by Enqvist and McDonald in [48, 347, 49, 328]. For a maximally charged AD

condensate ($B = 0$ in Eq. (274)), the linearized perturbation takes the form $\phi = \phi(t) + \delta\phi(x, t)$, and $\theta = \theta(t) + \delta\theta(x, t)$, where the homogeneous condensate is described by

$$\Phi = \frac{\phi(t)}{\sqrt{2}} e^{i\theta(t)} , \quad (289)$$

with $\phi(t) = (a_o/a)^{3/2} \phi_o$ and $\dot{\theta}(t)^2 \approx m_\phi^2$. Initially $\delta\phi(x, t)$ and $\delta\theta(x, t)$ should satisfy the relationship [151, 48, 49]

$$\delta\theta_i \approx \left(\frac{\delta\phi}{\phi} \right)_i . \quad (290)$$

The solution of the linear perturbation equations then has the form [48]

$$\delta\phi \approx \left(\frac{a_o}{a} \right)^{3/2} \delta\phi_o \exp \left(\int dt \left(\frac{1}{2} \frac{\mathbf{k}^2}{a^2} \frac{|K| m_\phi^2}{\dot{\theta}(t)^2} \right)^{1/2} \right) e^{i\mathbf{k} \cdot \mathbf{x}} \quad (291)$$

and

$$\delta\theta \approx \delta\theta_i \exp \left(\int dt \left(\frac{1}{2} \frac{\mathbf{k}^2}{a^2} \frac{|K| m_\phi^2}{\dot{\theta}(t)^2} \right)^{1/2} \right) e^{i\mathbf{k} \cdot \mathbf{x}} . \quad (292)$$

For the gravity mediation case, the above condition applies if $|\mathbf{k}^2/a^2| \lesssim |2K m_\phi^2|$, and H^2 is small compared with m_ϕ^2 and $|K| \ll 1$. If the first condition is not satisfied, then the gradient energy of the perturbations produces a positive pressure larger than the negative pressure due to the attractive force from the logarithmic term, preventing the growth of the perturbations.

For the case of a matter dominated Universe, the exponential growth factor is then [48, 49]

$$\int dt \left(\frac{1}{2} \frac{\mathbf{k}^2}{a^2} \frac{|K| m_\phi^2}{\dot{\theta}(t)^2} \right)^{1/2} = \frac{2}{H} \left(\frac{|K|}{2} \frac{\mathbf{k}^2}{a^2} \right)^{1/2} . \quad (293)$$

The largest growth factor will correspond to the largest value of $\mathbf{k}^2$ for which growth can occur,

$$\left. \frac{\mathbf{k}^2}{a^2} \right|_{max} \approx 2|K| m_\phi^2 . \quad (294)$$

The value of H at which the first perturbation goes non-linear is [48, 49]

$$H_i \approx \frac{2|K| m_\phi}{\alpha(\lambda)} , \quad (295)$$

with

$$\alpha(\lambda) = -\log\left(\frac{\delta\phi_o(\lambda)}{\phi_o}\right) , \quad (296)$$

where ϕ_o is the value of ϕ when the condensate oscillations begin at $H \approx m_\phi$. A typical value of $\alpha(\lambda)$ (e.g. with $d = 6$) is $\alpha(\lambda) \approx 30$. The initial non-linear region has a radius λ_i at H_i , which is given by [48, 49]

$$\lambda_i \approx \frac{\pi}{|2K|^{1/2}m_\phi} . \quad (297)$$

For the case of a non-maximally charged condensate the situation is slightly different. It is likely that the initial radius and the time at which the spatial perturbations initially go non-linear will roughly be the same [49] as for the maximally charged condensate. In general, the charge density of the initial non-linear lumps will essentially be the same as that of the original homogeneous condensate.

The perturbative evolution of a single condensate lump was considered in [49]. In terms of $\phi = (\phi_1 + i\phi_2)/\sqrt{2}$, the initial lumps are described by

$$\phi_1(r, t) = A \cos(m_\phi t)(1 + \cos(\pi r/r_0)) \quad (298)$$

$$\phi_2(r, t) = B \sin(m_\phi t)(1 + \cos(\pi r/r_0)) , \quad (299)$$

for $r \leq r_0$ and by $\phi_{1,2} = 0$ otherwise. The initial radius of the lump is $2r_0$, where $r_0 = \pi/(\sqrt{2}|K|^{1/2}m_\phi)$. The maximally charged condensate lump corresponds to $A = B$, while the non-maximal lump has $A > B$. The total energy and charge in a fixed volume are given by [328, 329]

$$E = 4\pi \int_V drr^2 \rho \quad , \quad Q = 4\pi \int_V drr^2 q \sim AB, \quad (300)$$

with $Q_{\max} = A^2$.

In [49], the behavior of the solutions was found in a perturbative analysis to depend on K , and to a greater extent on $Q/Q_{\max}$. The condensate lump was found to pulsate while charge is flowing out until the lump reaches a (quasi-)equilibrium pseudo-breather configuration, also called Q -axiton, with the lump pulsating with only a small difference between the maximum and minimum field amplitudes. For the Q -axiton,

in which the attractive force between the scalars is balanced by the gradient pressure of the scalar field, the energy per unit charge is much larger than m_ϕ ; indeed, the Q -axiton exists even if $Q = 0$. Only for a maximally charged Q -axiton are the properties similar to that of the corresponding Q -ball. It is however unclear whether Q -axitons are just an artifact of perturbation expansion.

5.10.2 Lattice simulations

The features of the fragmentation of the AD condensate cannot be fully captured by studying various mean field theory approaches, such as in large N-approximation and Hartree-approximation [239, 238, 243]. The formation of a Q -ball is a non-linear process for which various mode-mode interactions become important. This can be seen by expanding the perturbed ϕ and θ as shown by Kasuya and Kawasaki in [348, 349, 350]

$$\begin{aligned}\delta\ddot{\phi} + 3H\delta\dot{\phi} - 2\dot{\theta}\phi\delta\dot{\phi} - \frac{\nabla^2}{a^2}\delta\phi + U''(\phi)\delta\phi &= 0, \\ \phi\ddot{\theta} + 3H\phi\delta\dot{\theta} + 2(\dot{\phi}\delta\dot{\theta}) - 2\frac{\dot{\phi}}{\phi}\dot{\theta}\delta\phi - \phi\frac{\nabla^2}{a^2}\delta\theta &= 0.\end{aligned}\tag{301}$$

Although the potentials differ in the gauge and gravity mediated cases, it is nevertheless always possible to identify the fastest growing amplified mode. In the gravity mediated case we have already obtained that by inspecting Eq. (293). A similar analysis can be performed for the gauge mediated case by noting that $U''(\phi) \approx -2m_\phi^4/\phi^2$. Taking into account the conservation of charge $\dot{\theta}\phi^2a^3 = \text{const.}$, along with the approximation of a circular orbit, one may simplify Eq. (301) by seeking a solution of the form $\delta\phi = \delta\phi_0 \exp(\alpha t + ikx)$ and $\delta\theta = \delta\theta_0 \exp(\alpha t + ikx)$. In order to further simplify the analysis, one can also assume $a = \text{const.}$ and $\phi = \phi_0 = \text{const.}$, so that the phase velocity $\dot{\theta} = (U'/\phi)^{1/2} \approx \sqrt{2}m_\phi^2/\phi_0$. If α is real and positive, the fluctuations grow exponentially and become non-linear. Solving for $\delta\phi_0, \delta\theta_0$, Kasuya and Kawasaki finds for gauge mediated case [351]

$$\alpha^4 + 2\left(\frac{\mathbf{k}^2}{a^2} + \frac{2m_\phi^4}{\phi_0^2}\right)\alpha + \left(\frac{\mathbf{k}^2}{a^2} - \frac{4m_\phi^4}{\phi_0^2}\right)\mathbf{k}^2 = 0.\tag{302}$$

Note that in order for α to be positive, one must require the expression in the second parenthesis to be negative. This means that the instability band for the fluctuations

is given by

$$0 < \frac{\mathbf{k}}{a} < \frac{2m_\phi^2}{\phi_0}. \quad (303)$$

The most amplified mode appears at $(\mathbf{k}/a)_{max} \approx (3/2)^{1/2} m_\phi^2 / \phi_0$ in the gauge mediated case.

Various groups have studied the fragmentation of the AD condensate and the formation of Q -balls numerically. In [330], condensate fragmentation was simulated numerically on a $2 + 1$ dimensional 100×100 lattice, starting with a uniform AD-condensate with $\phi_0 = 10^9$ GeV and with an arbitrary phase ω , $\phi = \phi_0 e^{i\omega t} + \delta\phi$ with uniformly distributed random noise $\delta\phi \sim \mathcal{O}(10^{-13})|\phi_0|$ added to the amplitude and phase. The parameter values chosen for the simulations were $m_\phi = 10^2$ GeV, $K = -0.1$, and $\lambda = 1/2$. The results indicate that first the charge density of the condensate decreases uniformly due to the expansion of the Universe. As time progresses a growing mode can be seen to develop. White noise is still present but the growing mode soon starts to dominate. This process continues until lumps of positive charge develop. These are dynamically arranged in string-like features but the filament texture is a transient feature which disappears soon, see forthcoming Figs. (9,10).

The further evolution of the flat direction depends on the initial energy-to-charge ratio of the condensate, defined as

$$x \equiv \frac{E}{mQ}, \quad (304)$$

and hence on the value of ω . If $x = 1$, i.e. the energy-to-charge ratio of the condensate is equal to that of a Q -ball, no negatively charged Q -balls, anti- Q -balls, are formed. After the modes grow non-linear, the lumps just evolve into Q -balls and finally freeze due to the expansion of the Universe.

If $\omega < 1$ so that $x \gg 1$, the fragmentation process has a much more complicated history [330]. After the positively charged lumps have formed, expanded linearly and then developed non-linearly, the extra energy stored in them causes the lumps to fragment as they evolve into Q -balls. In this process a large number of negatively charged Q -balls forms. The total charge in the negative and positive Q -balls is approximately equal so that the initial charge in the condensate is in fact negligible compared to the amount of charge and anti-charge created.




Figure 7: Q -ball formation in gravity mediated supersymmetry breaking. From the left, the first plot shows the amplitude of the condensate after Q -balls have formed at $z = 6.3$. The second plot shows around 40 Q -balls with a largest charge $Q \simeq 5.16 \times 10^{16}$, and the third plot shows the power spectra of the condensate fluctuations ($k|\delta\varphi_k|^2$) when the amplitude of the fluctuations has become as large as the homogeneous mode: $\langle\delta\varphi^2\rangle \sim \varphi^2$. The top panels (a) and (c) show the full fluctuations calculated on one dimensional lattices, while the bottom panels (b) and (d) show the linearized fluctuations without mode mixing, from [350]

Full 3 + 1 dimensional simulations have been presented by Kasuya and Kawasaki [348, 349, 350] (for both gravity mediated and gauge mediated cases), see Figs. (7,8), and by Multamäki and Vilja [352] (for a wide range of the energy-to-charge ratio in the gravity mediated case with), see Figs. (9,10).

In [348], the authors simulated Q -ball formation in the gravity mediated scenario on a $(64)^3$ lattice with a lattice spacing $\Delta\zeta = 0.1$. The results are shown in Fig. (7). The initial fluctuations in the real and the imaginary direction were taken to be $\delta\phi_1 = \delta\phi_2 \sim \mathcal{O}(10^{-7})$, together with initial field values $\phi_1(0) = \phi_2(0) \sim \mathcal{O}(10^7)$ GeV. In order to obtain the spectrum the authors relied on a 1d lattice with $N = 1024$ and $\Delta\zeta = 0.1$. The result is depicted in the third plot of Fig. (7) for two different comoving time scales. One can see the marked difference between the linearized perturbations and the lattice simulated ones. In the latter case the spectrum does not fall sharply, which can be attributed to mode-mode interactions or rescattering effects which kick




Figure 8: Configuration of Q -balls on a three dimensional lattice in the gauge mediated case. More than 30 Q -balls have formed, and the largest one has the charge with $Q \simeq 1.96 \times 10^{16}$. The second plot shows the power spectrum of the condensate fluctuations when the amplitude of the fluctuations has become as large as the homogeneous mode. The top panels (a) and (c) show the full fluctuations calculated on one dimensional lattices, while the bottom panels (b) and (d) show the linearized fluctuations without mode mixing, from [349].

the lower momentum modes higher, leading to a broadening and smoothening of the spectrum. Note that in linearized fluctuations the instability band is almost the same as Eq. (294). For example $\mathbf{k}/m_\phi = \sqrt{2}a(\tau)|K|^{1/2} \approx 2$ for $|K| = 0.01$ and $\tau = 5.5 \times 10^3$.

In [349], Kasuya and Kawasaki repeated their simulation for the gauge mediated case, which is shown in Fig. (8). Note that the size of the Q -ball is bigger than in the gravity mediated case.

Multamäki and Vilja [352] had typical lattice sizes of 120^3 . They verified that the 2 dimensional simulations [330] capture all of the essential features of the AD condensate fragmentation: transient filament structures resulting in a large number of Q -balls and anti- Q -balls, which can also be seen in Fig. (10). Note that when the condensate has the exact energy-to-charge ratio of a Q -ball so that $x = 1$, no anti- Q -balls form in Fig. (9), whereas for $x \gg 1$, the number of Q -balls and anti- Q -balls are practically equal.




Figure 9: Formation and evolution of charged lumps on 3D lattice from comoving time $\tau = 875 - 3000$, when $x = 1$. Only positively charged Q -balls have formed, from [352].




Figure 10: Formation and evolution of charged lumps from comoving time $\tau = 1500 - 7500$, when $x = 10^5$. Here positive and negatively charged Q -balls have formed, depicted in different shades, from [352].

5.11 Equilibrium ensembles

Some of the lattice results can be understood by analytical arguments. In particular, the distributions and number densities of Q -balls and anti- Q -balls may be obtained simply by maximizing the entropy. Such approach appears justified, since after fragmentation, the AD lumps are expected to interact vigorously and the field fragments will settle to the state of lowest energy by emitting and exchanging smaller fragments. If the interaction is fast enough compared with the expansion rate of the Universe, i.e. $\Gamma = n_{tot}\sigma v > H = (2/3t)$, where n_{tot} is the total number of Q -balls and anti- Q -balls, $\sigma \approx \pi R_Q^2$ is the geometric cross-section of a Q -ball collision, and v is the average velocity of a Q -ball, then it is naturally expected that the final state should consist of an equilibrium distribution of Q -balls and anti- Q -balls in a state of maximum entropy. Considerations supporting this has been given in [330] for the case of gravity mediated supersymmetry breaking.

The Q -ball (anti- Q -ball) distributions $N_+(Q, p)$ ($N_-(Q, p)$) are subject to the following constraints:

$$\begin{aligned} E_{tot} &= E_+ + E_- , \quad E_{\pm} = \int dQ dp E(Q, p) N_{\pm}(Q, p) \\ Q_{tot} &= Q_+ - Q_- , \quad Q_{\pm} = \int dQ dp Q N_{\pm}(Q, p) , \end{aligned} \quad (305)$$

where $E(Q, p) \approx \sqrt{p^2 + m^2 Q^2}$ is the energy of a single Q -ball, E_+ (E_-) and Q_+ (Q_-) are the energy and charge of Q -balls (anti- Q -balls), and E_{tot} and Q_{tot} are respectively the total energy and charge of Q -balls and anti- Q -balls, which are equal to the energy and charge of the initial AD-condensate (unless significant amounts of energy and/or charge are transformed into radiation). It then follows from Eq. (305) that

$$x \equiv E_{tot} \geq \frac{m (Q_+ + Q_-)}{m Q_{tot}} \geq 1. \quad (306)$$

This condition is independent of the precise form of the Q -ball distributions.

If all the baryon asymmetry resides in Q - and anti- Q -balls, then at times earlier than $10^{-6}s$, $Q_{tot}/Q_+ \sim \Delta B \sim 10^{-8}$. (Since $B - L$ is conserved in the MSSM, this holds also for the purely leptonic flat directions.) From Eq. (306) it follows

$$x \simeq 10^8 . \quad (307)$$

Even if all of the baryon asymmetry were not carried by Q -balls, a natural expectation is $x \gg 1$ so that the number of Q -balls, N_+ , and the number of anti- Q -balls, N_- , are approximately equal and the total number of Q -balls is $N_{tot} \approx 2N_+$. The main bulk of the Q -balls may be expected to be relativistic [330], and the collision rate much larger than the Hubble rate.

Indeed, one can verify that the equilibrium assumption is self-consistent, and it is also supported by numerical studies in 2 dimensions [330] as well as by the 3 dimensional simulations of Multamäki and Vilja [352], who actually observe the Q -ball distribution to relax into equilibrium as shown in Fig. (10).

6 Q -balls

Q -balls with a global charge have many interesting astrophysical and cosmological consequences. They may reheat the Universe [353, 354] and serve as candidates for dark matter candidate [151, 355, 356, 351, 357, 358]. They may provide baryogenesis and leptogenesis [48, 49, 347] and while decaying, produce LSP dark matter [359, 360, 351, 361, 362]. They could be responsible for the generation of cosmic magnetic fields [363]. Q -balls could stabilize neutron stars [364] or even form solitonic q-stars [365, 366, 367]. They could act as a laboratory for physics beyond the electroweak scale [331].

6.1 Q -ball as a non-topological soliton

6.1.1 Proofs of existence

The Q -ball is a generic ground state in a broad class of theories with interacting scalar fields carrying some conserved global charge [345, 346, 368, 369, 47, 365]. The Q -ball is an example of a non-topological soliton whose boundary condition at infinity is the same as that for the vacuum state, unlike in the case of topological solitons such as magnetic monopoles [370, 371] (a detailed review on non-topological solitons can be found in, e.g. [372, 365]).

Q -balls are quite generic solitons in $3 + 1$ dimensions, which can be associated with many scalar fields with various $U(1)$ charges [50], with a non-Abelian symmetries [373, 374, 375], and also with local gauge symmetries [376, 377, 378, 356, 379, 380, 381]. The main difference which distinguishes global Q -balls from a local Q -balls is that in the latter case the charge of the stable Q -ball is bounded from above.

Recall that in order to find a Q -ball solution one must minimize the energy E_ω (see Eqs. (285,286)) for a fixed charge with respect to the variations of ω and $\varphi(x)$ independently. Obviously one could try finding directly a Q -ball solution by solving Eq. (288). A more effective way is to seek a bounce solution $\bar{\varphi}(x)$ for tunneling in three Euclidean dimensions in the potential given by Eq. (287) [382, 384, 158]. Note that the first term in Eq. (286) is then the three dimensional Euclidean action $S_3[\bar{\varphi}_\omega(x)]$ of

this bounce solution which satisfies Eq. (288) in radial coordinates with the boundary conditions $\varphi(0) = \varphi'(\infty) = 0$.

A theorem [47, 365] (see also [382]) states that if there exists a range of φ however small, and the potential $U(\varphi^2)$ contains an attractive interaction however weak, then a non-topological soliton solution exists for

$$\nu^2 \leq \omega^2 < m_\phi^2, \quad (308)$$

where the mass parameter is defined as $U(\varphi^2) \rightarrow m_\phi^2 \varphi^2$ when $\varphi \rightarrow 0$. The value of $\omega = \sqrt{k^2 + m_\phi^2}$ determines the frequency of the φ quanta in the field space. For a plane wave solution it is always true that $\omega^2 > m_\phi^2$ while for a solitonic solution $\omega^2 < m_\phi^2$. This suggests that there exists a parabola $\nu^2 \varphi^2$ tangent to $U(\varphi^2)$ at $\varphi = \pm\varphi_0$, with $\nu^2 < m_\phi^2$. Another useful way of expressing this is through

$$\frac{U(\varphi)}{\varphi^2} = \min, \quad \text{for } \varphi = \varphi_0 > 0. \quad (309)$$

For a sufficiently large Q given by Eq. (285), the energy of a soliton is then given by

$$E = |\nu Q| < m_\phi |Q|, \quad (310)$$

which ensures its stability against decay into plane wave solutions with $\varphi \simeq \varphi_0$ inside and $\varphi \simeq 0$ outside the soliton. Note that the global $U(1)$ symmetry is thus broken inside the soliton by the vev, however, remains unbroken outside the soliton. The most crucial piece is the presence of a global $U(1)$ charge of the Q -ball which actually prevents it from decaying and makes the soliton stable. (For analytical results on Q -ball properties, see [383]). The above discussion can be repeated in the presence of several charges q_i , and an analogue of Eq. (309) has been established by Kusenko in [50]. The main difference between a single and multi-charged Q -ball is that there exists different vevs corresponding to different charges, which modifies the appropriate bounce solution Eq. (309).

6.1.2 Beyond thin wall solution

The above discussion tacitly assumes a thin wall approximation where the edge of a soliton is sharply defined, which means that the gradient energy is small compared

to the volume energy. As φ_0 increases, the thin wall limit breaks down for a fixed charge, and one must seek other methods in order to guarantee the existence of a Q -ball solution as pointed out by Kusenko [382]. For a flat potential, which mimics the large field value situation, the equations of motion Eq. (288) can be solved near the origin $r = 0$, and for large r . The Q -ball profile can be found to be

$$\begin{aligned}\varphi(r) &= \varphi_0 \frac{\sin(\omega r)}{\omega r}, \quad r < R, \\ &= \varphi_1 e^{-m_\phi r}, \quad r \geq R,\end{aligned}\tag{311}$$

where the values of $\varphi_0, \varphi_1, \omega$ and R are such that they minimize E_ω , while $\varphi(r)$ interpolates $r = R$, the size of the Q -ball, continuously. In a thick wall limit one can also write down E_ω in terms of dimensionless variable $\xi = \omega x$ and $\psi = \varphi/\omega$, while neglecting all the terms in $\hat{U}_\omega$ except the constant term and the $\omega^2 \varphi^2$ term [382, 158], one obtains $E_\omega \approx a\omega + b/\omega^3 + \omega Q$, where a, b are constants independent of ω . The size of the Q -ball turns out to be $R \propto 1/\omega$, and the vev of the field in the Q -ball interior is $\varphi_0 \propto Q^{1/4}$ from Eq. (311). Note that there is no classical limit on the charge of a Q -ball. In fact no matter how small Q is there always exists a value ω close to m_ϕ , for which E_ω is minimized. Quantum stability requires that $Q \geq 1$. When $Q \rightarrow 1$, the quantum corrections will indeed become important. For $Q \geq 7$ the Q -balls are quantum mechanically stable configurations [386].

The Q -ball has been shown to be classically stable by Coleman [47] (see also [365, 382]). The semiclassical approach obviously breaks down when quantum fluctuations around the Q -ball are comparable to the energy of the system itself. This happens when $Q \approx 1$. Even though the charge becomes small, the Q -ball size remains large in comparison to the Compton length m_ϕ^{-1} [382, 385, 386]. One can also establish a Virial theorem, which holds for any Q and does not require any approximation [50].

6.2 Varieties of Q -balls

6.2.1 Thin wall Q -balls

There exist thin wall Q -ball solutions, where the boundary is a well defined edge, as well as thick wall Q -balls, where the boundary is not localized in a narrow region and

the soliton is typically described by a Gaussian profile. Both Q -ball types may appear within the same theory.

The thin wall Q -ball is the simplest and arises naturally in any suitable scalar potential that allows for the existence of a Q -ball. As mentioned already, thin wall solution has the profile $\varphi(r) \approx \varphi_0 \delta(r - R)$ in the radial direction, where R is the size of the radius of a spherically symmetric Q -ball. This obviously neglects the surface energy contribution and yields [47]

$$\frac{E}{Q} = \min \sqrt{\frac{U(\varphi^2)}{\varphi^2}} \approx \omega_c. \quad (312)$$

Energy is thus growing linearly with charge. Note that the radius of such a Q -ball can be very large,

$$Q = 2\omega_c \varphi_0^2 V = \frac{8\pi}{3} \pi R^3 \omega_c \varphi_0^2. \quad (313)$$

These are useful relationships for the purposes of this Section.

6.2.2 Thick wall Q -balls in the gauge mediated case

Thick wall Q -balls have been widely considered in the literature within gauge and gravity mediated supersymmetry breaking scenarios. In both cases the mass of the Q -ball grows more slowly compared to a thin wall case, i.e. the scalar potential grows slower than φ^2 . In this case Q -ball never reaches a thin wall regime, even if Q is large. The value of φ inside a Q -ball extends as far as the gradient terms allow, and the mass of a Q -ball is proportional to Q^p , where $p < 1$.

In the context of gauge mediated supersymmetry breaking the AD potential takes the form (without the non-renormalizable contributions) [151, 331] (see Eq. (204), in Sect. 5.3.2.)

$$U(\varphi) \approx m_\phi^4 \log \left(1 + \frac{|\varphi|^2}{m_\phi^2} \right), \quad (314)$$

where $m_\phi \sim \mathcal{O}(\text{TeV})$ represents the supersymmetry breaking scale. In [387], the authors have considered the effective potential of the form

$$U \sim F^2 \left[\log \left(\frac{|\varphi|^2}{m_\phi^2} \right) \right], \quad (315)$$

where $F^{1/2} \gg m_\phi$. Despite the difference between the forms of Eqs. (314) and (315), the dynamics of the flat direction is similar to the one given in Eq. (314), which yields the equation of motion

$$\varphi'' + \frac{2}{r}\varphi' = -\omega^2\varphi. \quad (316)$$

For large r , $\varphi(r) \sim \exp(-m_\phi r)$, where m_ϕ is the mass of φ near the origin. The interpolating solution was already presented in Eq. (311). The energy of such a Q -ball grows as [151]

$$E \approx \frac{4\sqrt{2}}{3}\pi m_\phi Q^{3/4}. \quad (317)$$

The profile of the Q -ball is given by $\varphi(r) \sim \exp(-m_\phi r)$. The radius and the value of the vev inside the Q -ball is roughly given by [151]

$$R \approx \frac{1}{\sqrt{2}m_\phi}Q^{1/4}, \quad (318)$$

$$\varphi_0 \approx \frac{m_\phi}{\sqrt{2}\pi}Q^{1/4}. \quad (319)$$

6.2.3 Thick wall Q -balls in the gravity mediated case

If the potential grows only slightly slower than φ^2 as in the case of gravity mediated supersymmetry breaking scenarios, the potential may be approximately written as [48, 347, 49] (see Eq. (200), in Sect. 5.3.1.)

$$U(\varphi) \approx m_\phi^2 \left(1 + K \log \left[\frac{\varphi^2}{M^2} \right] \right) \varphi^2, \quad (320)$$

where $K < 0$, and M is the largest mass scale. Note that at small vevs we have again neglected the non-renormalizable contributions in the above potential. The mass scale is given by $m_\phi \sim m_{3/2} \sim \mathcal{O}(\text{TeV})$.

The Q -ball equation of motion is written as

$$\varphi'' + \frac{2}{r}\varphi' = -\omega_0^2\varphi + m_\phi^2\varphi K \log \left(\frac{\varphi^2}{M^2} \right), \quad (321)$$

where ω_0 is defined by

$$\omega_0^2 = \omega^2 - m_\phi^2(1 + K). \quad (322)$$

For thin wall Q -balls, the initial value of φ is very close to φ_{c1} , the value of φ for which the right-hand side of Eq. (321) vanishes. In this case φ will remain close to φ_{c1} up to a radius of order $\omega_0^{-1} \log(\varphi_{c1}/\delta\varphi(0))$, where $\delta\varphi(0) = (\varphi_{c1} - \varphi(0))$. It will then decrease to zero over a distance $\delta r \approx \omega_0^{-1}$, corresponding to the width of the wall of a Q -ball. The radius of a thin wall Q -ball can be made arbitrarily large by choosing $\delta\phi(0)$ small enough.

For a thick wall Q -ball, the initial value of φ can be much smaller than φ_{c1} . In this case the non-renormalizable terms may be neglected. In general, the right-hand side of the Q -ball equation of motion vanishes for three values of φ , which correspond to φ_{c1} , φ_{c2} and zero. φ_{c2} corresponds to the point at which, assuming that the non-renormalizable terms can be neglected, the first two terms on the right-hand side of Eq. (321) cancel, and we obtain [49]

$$\begin{aligned}\varphi_{c2}(r) &= Me^{\left(\frac{\omega_0^2}{2Km_\phi^2}\right)}, \\ &= Me^{-(1-\omega^2/m_\phi^2-2K)}e^{Km_\phi^2r^2/2}.\end{aligned}\tag{323}$$

In deriving the last correspondence we have used Eq. (322). Note that φ_{c2} is an attractor, in a sense that if $\varphi(0)$ is close to φ_{c2} it will tend towards φ_{c2} as r increases.

The radius of a thick wall Q -ball in the gravity mediated case is given by [48, 347, 49]

$$R \approx \frac{1}{|K|^{1/2}m_\phi},\tag{324}$$

where R is defined as the radius within which 90% of the Q -ball energy is found, and [48, 347, 49]

$$\omega_0 \approx |K|^{1/2}m_\phi.\tag{325}$$

Since typically $|K|$ is small compared with 1, we find $\omega \approx m_\phi$. In the gravity mediated supersymmetry case the size of a Q -ball does not depend on charge, unlike in the gauge mediated case, see Figs. (7,9,10), where the sizes of the Q -balls are all equal.

One may take a Gaussian ansatz for the profile of a thick wall Q -ball [49]

$$\varphi(r) = \varphi(0)e^{-\frac{r^2}{R^2}},\tag{326}$$

provided one identifies $\omega_0^2 \approx 3|K|m_\phi^2$ and $R^2 \approx 2(|K|m_\phi^2)^{-1}$.

The total charge of the Gaussian thick wall Q -ball is given by [49]

$$Q = \int dr 4\pi r^2 \omega \varphi_0^2 e^{-\frac{2r^2}{R^2}} = \left(\frac{\pi}{2}\right)^{3/2} \omega \varphi_0^2 R^3 \quad (327)$$

while the total energy is given by [49]

$$E \approx \frac{3}{2} \left(\frac{\pi}{2}\right)^{3/2} \varphi_0^2 R + \left(\frac{\pi}{2}\right)^{3/2} m_\phi^2 \varphi_0^2 R^3, \quad (328)$$

where the second term in the above equation is the combined contribution from the potential energy and the charge term, where we have used $\omega^2 \approx m_\phi^2$. Since R is large compared with m_ϕ^{-1} for small $|K|$, the potential plus charge term dominates the energy. The radius within which 90% of the energy lies is then given by $R_c = 1.25R$. The energy per unit charge is given by [49]

$$\frac{E}{Q} = \frac{m^2}{\omega} \approx \left(1 + \frac{3|K|}{2}\right) m_\phi, \quad (329)$$

where we have used the Gaussian result $\omega_0^2 = 3|K|m_\phi^2$. For all practical purposes we can take $E \approx m_\phi Q$. Although the energy per unit charge is larger than m_ϕ , the mass of the scalar at small values of φ will have the form $m_\phi(1 + \alpha|K|)$ (with $\alpha \gtrsim 1$) once the logarithmic correction to the potential is included, so that the binding energy per unit charge will be positive and of order $|K|m_\phi$.

The last two examples Eqs. (314,320) exhibit two extremes of any thick wall type Q -ball. Any thick wall Q -ball should belong somewhere in between, such as in hybrid case.

6.2.4 Hybrid case: gauge and gravity mediated Q -ball

In Sect. 6.2.2., we discussed the Q -ball potential in the gauge mediated case, but it is true that any generic flat direction is also lifted by gravity mediation as well. In the gauge mediated case the full flat direction potential relevant for Q -ball formation should read as [349, 351]

$$U(\varphi) = m_\phi^4 \log \left(1 + \frac{|\varphi|^2}{m_\phi^2}\right) + m_{3/2}^2 |\varphi|^2 \left[1 + |K| \log \left(\frac{|\varphi|^2}{M_{\text{P}}^2}\right)\right], \quad (330)$$

where $m_{3/2}$ takes values between 100 KeV and 1 GeV. The gaugino loops lead to $K < 0$ as discussed before (see Sect. 5.3.1.), but Yukawa couplings give rise to $K > 0$. On the other hand if the Yukawas dominate, the AD condensate can be stabilized and Q -balls can only form for $\varphi \leq \varphi_{eq}$. The second term in the above potential dominates when

$$\varphi \geq \varphi_{eq} \equiv \sqrt{2} \frac{m_\phi^2}{m_{3/2}}. \quad (331)$$

There are two distinct regimes. When $\varphi \geq \varphi_{eq}$, the Q -ball properties resemble the gravity mediated thick wall case. Otherwise, when $\varphi < \varphi_{eq}$, the Q -ball properties are similar to the gauge mediated thick wall case. The energy per unit charge can be written as [349, 351]

$$\frac{E}{Q} \sim \begin{cases} m_\phi Q^{-1/4} & \varphi \leq \varphi_{eq} \\ m_{3/2} & \varphi \geq \varphi_{eq}. \end{cases} \quad (332)$$

Obviously in between there should be a hybrid regime which interpolates smoothly between the gauge and gravity mediated cases.

6.2.5 Effect of gravity on Q -balls

In principle gravity can give a significant contribution to the Q -ball energy, as shown in a study by Multamäki and Vilja [388]. In a thin wall case, the interesting result is that gravity limits the maximal size of a Q -ball. The reason is that besides the rotational motion in the complex field space which generates outward pressure, there exists a gravitational attraction. Since the gravitational contribution to the Q -ball energy is negative, it is possible that gravity can render an otherwise unstable Q -ball stable. The effect of gravity on Q -balls remains small provided the soliton is much larger than the Schwarzschild radius and the charge smaller than the gravitational charge $Q_g \sim (M_P/m_\phi)^4$, which is quite large $Q_g \sim 10^{64}$ for $m_\phi \sim 100$ GeV.

6.2.6 Q -balls and local gauge invariance

So far we have considered Q -ball solutions in theories with a global $U(1)$ symmetry. The symmetry group can however be extended to include global non-Abelian symmetries [373, 374, 375, 380]. The existence of Q -balls in a supersymmetric Wess-Zumino

model has been demonstrated by Axenides, Floratos and Kehagias [375], who showed that Q -balls form domains of manifestly broken supersymmetry.

A gauged Q -ball has some additional interesting properties [376, 378, 356, 379, 381]. Taking a complex scalar field $\phi(\mathbf{r}, t) = f(\mathbf{r}, t) \exp(-i\theta(\mathbf{r}, t))/\sqrt{2}$ coupled to an Abelian gauge field A_μ , the charge of a given field configuration is given by [376]

$$Q_\phi = \int d^3r f^2 (\dot{\theta} - eA_0) , \quad (333)$$

where e is the gauge charge. Since the gauge field inside a Q -ball is broken by the non-zero vev of φ which couples to the gauge field canonically, the gauge field is massive and acts as a $U(1)$ superconductor, provided the Compton wavelength of the gauge field is smaller than the size of a Q -ball. As a result of broken gauge symmetry, there is an extra source of electrostatic self-energy contribution which comes from the electrostatic repulsive force due to the presence of a gauge charge. The gauged charges are also repelled to reside on the boundary of a Q -ball. In a gauged Q -ball, for a fixed charge Q , both radius and energy are relatively large. The remarkable feature is that there is a maximum allowed charge and correspondingly a maximum radius. This is due to the repulsive electrostatic potential A_0 which tends to destabilize the gauged Q -ball. The maximum charge-to-radius ratio is governed by the gauge charge: $Q_{max}/R_{max} = 4\pi(m_\phi - \omega_c)/e^2$.

In [379], it has been argued that the presence of fermions could stabilize the gauged Q -ball. During Q -ball formation, fermions could be trapped inside the Q -ball. Those with charges equal to the charge of the scalar quanta would be repelled from the inside, whereas fermions with opposite charges would remain and render the Q -ball neutral. Large electric fields inside the Q -ball could also lead to a pair production and to a subsequent screening of the charge.

An interesting application of a gauged Q -ball could be the hadronic structure of QCD, as in the Friedberg-Lee model [389], where hadrons are modeled by phenomenological non-topological solitons. Gauged Q -balls in theories with a Chern-Simons terms has also been considered in [377].

6.3 Q -ball decay

6.3.1 Surface evaporation to fermions

In the MSSM, the scalar field forming a Q -ball can interact with fermions. Then a Q -ball can decay into a pair of fermionic quanta. Q -ball decay has been considered in [390, 391], where fermion production was studied in a classical background of a Q -ball. As it was first pointed out in [390], for a large Q the Q -ball evaporates through its physical surface and there exists an upper bound on the evaporation rate per unit area. This behavior has been verified by numerical studies [391].

In order to understand this, let us imagine that a region inside a Q -matter with a vanishing ϕ , which might appear due to fermion pair production, forms a cavity. Suppose L is the linear size of the cavity. The energy of Q -matter formerly inside the cavity is given by [390]

$$E_Q \sim \omega_0^2 \phi_0^2 L^3, \quad (334)$$

while the charge Q within that region is given by [390]

$$\Delta Q \sim \omega_0 \phi_0^2 L^3. \quad (335)$$

N massless fermions inside the cavity will have a free Fermi gas distribution, and therefore, the energy of the fermions is given by [390]

$$E_\psi \sim \hbar \frac{N^{4/3}}{L} \sim \left(\frac{\hbar}{L} \right) \left(\frac{\Delta Q}{\hbar} \right)^{4/3}. \quad (336)$$

The ratio of energies is given by [390]

$$\frac{E_Q}{E_\psi} \sim \left(\frac{\hbar \omega_0^2}{\phi_0^2} \right)^{1/3}. \quad (337)$$

In the semiclassical limit $\hbar \rightarrow 0$, cavitation is energetically forbidden irrespective of the size of a cavity. In a sense, it is the Pauli exclusion principle which keeps the Q -ball stable. Fermions are produced but there exists a Fermi pressure which prevents further production. Inside Q -matter fermions gain mass of order $g\phi_0$ and saturate the Fermi energy so that fermions can be produced only from the surface.

Following [390], let us assume that each fermion carries the energy $\sim \hbar\omega_0/2$. A simple bound on the average fermionic pair production from the surface can then

be obtained by noting that maximum current density can be reached only when the outward moving fermions from the Q -wall are occupied while every level in the inward moving fermions is empty. Then, by assuming massless Weyl fermions with a single helicity state for each $\mathbf{k}$, one obtains the limit on the outward moving current [390]

$$\langle \mathbf{n} \cdot \mathbf{j} \rangle_{\omega \leq \omega_0/2} \leq \frac{1}{(2\pi)^3} \int_0^{\omega_0/2} k^2 dk \int_0^1 \cos(\theta) d(\cos(\theta)) \int_0^{2\pi} d\phi = \frac{\omega_0^3}{192\pi^2}. \quad (338)$$

Integrating this expression over the enclosing surface yields the maximum evaporation rate per unit area in the limit $R \rightarrow \infty$ [390]

$$\frac{dQ}{dt dA} \leq \frac{\omega_0^3}{192\pi^2}. \quad (339)$$

A detailed calculation relies on estimating the reflection and transmission coefficients of the ingoing and outgoing waves outside the Q -ball and matching these with the solutions obtained inside the Q -ball. In order to calculate the transmission coefficient, one has to sum over infinite sequence of scatterings. In [390], it was shown that the maximum transmission coefficient is 1 when $\omega_0 \leq g\phi_0$. The weak coupling limit $\omega_0 > g\phi_0$ leads to a different transmission coefficient and the rate of evaporation is given by [390]

$$\frac{dQ}{dt dA} \simeq 3\pi \left(\frac{g\phi_0}{\omega_0} \right) \left(\frac{\omega_0^3}{192\pi^2} \right) = \frac{g^2 \omega_0^2 \phi_0}{32\pi^2}. \quad (340)$$

The factor $g\phi_0$ determines the penetration width of the fermions. In other words, in the weak coupling limit fermions can penetrate deep inside a Q -ball without completely filling the Fermi sphere. Within Pauli exclusion principle, Q -matter could decay into very weakly coupled fermions within the whole Q -ball volume. This will be discussed in the context of L -balls in Sect. 7.1, which can decay into massless neutrinos throughout the interior.

In [390, 391], it was shown numerically that the evaporation rate is strongly dependent on R but approaches the limiting profile given in [390]. In a realistic case a step-function is not always a good approximation, in particular for a thick wall Q -ball. In the thick wall case the problem has been investigated numerically by Multamäki and Vilja [391], who found that for a sufficiently large Q the evaporation rate decreases with increasing Q . As a result the Q -ball evaporates faster when its size decreases.

6.3.2 The decay temperature

For a thin wall Q -ball the surface area is related to the charge via $A = (36\pi)^{1/3}Q^{2/3}/(\kappa^{2/3})$, where $\kappa = (2\varphi_0^2 U(\varphi_0))^{1/2}$. The lifetime of a Q -ball in this case is given by [49]

$$\tau = 144\pi \left(\frac{4\pi}{3}\right)^{2/3} \frac{\kappa^{2/3}Q^{1/3}}{\omega^3} . \quad (341)$$

For a thick wall case, the area of a Q -ball is independent of its charge, being fixed by its radius $R \approx (|K|^{1/2}m_\phi)^{-1}$. The Q -ball lifetime in this case is then given by [49]

$$\tau = \frac{48\pi Q}{R^2\omega^3} . \quad (342)$$

From the above expression, one can estimate the temperature at which the Q -balls decay. By assuming radiation domination, the decay temperature is defined as [49]

$$T_d = \left(\frac{1}{k_T}\right)^{1/2} \left(\frac{M_{\text{P}}}{2\tau}\right)^{1/2} , \quad (343)$$

where $k_T = \left(\frac{4\pi^3 g(T)}{45}\right)^{1/2}$. By taking $\omega \approx m_\phi \approx 100$ GeV, thick wall Q -balls will decay at a temperature [49]

$$T_d \approx \frac{15}{|K|^{1/2}} \left(\frac{\omega}{100 \text{ GeV}}\right)^{1/2} \left(\frac{10^{15}}{Q}\right)^{1/2} \text{ GeV} , \quad (344)$$

where we have set $k_T \approx 17$. The Q -balls will decay at a temperature less than 100 GeV if $Q \gtrsim 2 \times 10^{13}|K|^{-1}$. For thin wall Q -balls, the right-hand side of Eq. (344) has an additional factor $(Q/Q_c)^{1/3}$, where Q_c is the value at which the thin wall limit becomes valid.

6.3.3 Q -ball decay into a pair of bosons

If the Q -ball forming scalar is not the lightest, as might be the case in the MSSM, it is also possible that the Q -ball decays into lighter bosons. The decay into scalar fields within the Q -ball volume is not blocked by the Pauli exclusion principle. In the case of bosons one should replace $N^{4/3}$ by N in Eq. (336), because the bosons can condense in the lowest mode of the cavity. The ratio Eq. (337) becomes

$$\frac{E_Q}{E_b} \sim \omega_0 L . \quad (345)$$

Cavitation in the case of bosons is always energetically favorable for sufficiently large L . Q -ball decaying into light scalars could be significantly enhanced relative to the decay into fermions.

The decay into light scalars will only be possible near the edge of a thick wall Q -ball. This is because particles coupling directly to the condensate scalars will gain a large effective mass from $\langle\phi\rangle$ inside the Q -ball. As a result, decay into light scalars will only occur via loop diagrams with rates suppressed by the large effective mass. This tends to make MSSM B -balls long-lived, as will be discussed later.

Not all the flat directions would have a scalar decay mode. An example is $\bar{u}\bar{d}\bar{d}$ direction, which is lifted at $d = 6$ and contains the right-handed squarks. For the universal boundary condition for squarks at large scales, RG flow analysis suggests that the left-handed squark masses are typically heavier than the right-handed ones. The Higgs scalar mass could also be heavy compared to the right-handed squark mass and the slepton masses usually come out to be lighter than right-handed squarks. Hence the decay of such baryonic direction would be kinematically forbidden. Even if the right-handed squark decays to Higgses and to sleptons, it would certainly involve either a pair of (light) quarks, gauginos, Higgsinos or leptons in the final state. As a result the core of a $\bar{u}\bar{d}\bar{d}$ Q -ball would be Fermi suppressed. Any final state involving a pair of fermions can arise only from the surface.

Generically Q -balls made up of left-handed squarks and sleptons, such as the $d = 6$ MSSM flat directions $\bar{d}QL$ or $\bar{e}LL$ are expected to decay into a pair of light bosons. Note that these particular directions are good candidates for carrying B charge. One may parameterize by f_s the possible enhancement factor of the scalar decay rate over the fermion decay rate, so that [49]

$$\left(\frac{dQ}{dt}\right)_{boson} = f_s \left(\frac{dQ}{dt}\right)_{fermion} . \quad (346)$$

This gives for the decay temperature [49]

$$T_d \approx \left(\frac{f_s \omega^3 R^2 M_P}{48\pi k_T Q}\right)^{1/2} \approx 0.06 \left(\frac{f_s}{|K|}\right)^{1/2} \left(\frac{m_\phi}{100 \text{ GeV}}\right)^{1/2} \left(\frac{10^{20}}{Q}\right)^{1/2} \text{ GeV} . \quad (347)$$

For $\bar{d}QL$ or $\bar{e}LL$ one may estimate the largest possible enhancement factor using the Gaussian thick wall ansatz as shown in [49]. Within the Q -ball, for $\phi \gg m_\phi$, the

lowest possible dimension operator which could allow the condensate scalars to decay at one loop level to light particles is the one lifted by the $d = 5$ operator

$$\frac{1}{M} \int d^4\theta \varphi \chi^\dagger \eta , \quad (348)$$

where χ and η represent the light particles and $M \approx g\varphi$, where g is the coupling of the heavy particles to φ . The decay rate of the condensate scalars to light scalars will then be given by [49]

$$\frac{dQ}{dt} = - \int \omega \varphi^2(r) \Gamma(r) 4\pi r^2 dr , \quad (349)$$

where $\omega \varphi^2(r)$ is the charge density within the Q -ball, and [49]

$$\begin{aligned} \Gamma(r) &\approx \frac{\alpha^2 m_\phi^3}{\phi^2} ; \quad g\phi > m_\phi \\ &\approx \alpha m_\phi ; \quad g\phi < m_\phi , \end{aligned} \quad (350)$$

with $\alpha = g^2/(4\pi)$ (for simplicity we consider a single coupling constant g). Let r_* be the radius at which $\varphi(r) = m/g$. Then the largest contribution to the decay rate will come from a region of width $\delta r \approx R^2/(4r_*)$ around r_* , over which φ has a roughly constant value $\varphi \approx m/g$, where

$$r^* = \gamma R ; \quad \gamma = \ln^{1/2} \left(\frac{g\varphi(0)}{m_\phi} \right) . \quad (351)$$

From Eq. (349), the rate can be deduced as [49]

$$\frac{dQ}{dt} \approx -4\pi\alpha\omega m_\phi \left(m_\phi^2 \alpha \int_0^{r_*} dr r^2 + \phi^2(0) \int_{r_*}^\infty dr r^2 e^{-\frac{2r^2}{R^2}} \right) \quad (352)$$

$$\approx -\frac{12\pi\gamma}{|K|^{1/2}} \left(1 + \frac{\gamma^2 g^4}{3\pi} \right) \left(\frac{dQ}{dt} \right)_{fermion} . \quad (353)$$

where we have used $\omega \approx m_\phi$. For a thick wall Q -ball and for typical values of the parameters, $g\varphi(0)/m \approx (0.1 - 0.01)Q^{1/2}$, so that $\gamma \approx 4.5$ in Eq. (353). One finds that the enhancement factor is typically [49]

$$f_s \approx \frac{170}{|K|^{1/2}} \left(1 + 2.1g^4 \right) . \quad (354)$$

if g is less than 1 then for $|K| \approx 0.01 - 0.1$, we expect an enhancement factor not much larger than about 10^3 . For g less than 1, most of the enhancement factor comes from the unsuppressed tree level decays occurring at $r > r_*$.

6.4 Cosmological formation of Q -balls

Cosmological formation of Q -balls was initially proposed by Kusenko in [50], by Kusenko and Shaposhnikov in [151], and by Enqvist and McDonald who studied Q -balls from the fragmentation of the flat direction condensates in the gravity mediated case [48, 49]. Subsequently, the problem has been attacked numerically by Kasuya and Kawasaki in case of gravity mediation [350, 351] as well as in gauge mediation [349, 351], and by Multamäki and Vilja in the gravity mediated case [352].

6.4.1 In gravity mediated case

In the gravity mediated case the size of the Q -ball depends on the charge; the larger the charge, the larger is the size of a Q -ball, see Fig. (8). In a cosmological context it is natural to think that the Q -balls form when the the most amplified AD condensate mode is as large as the horizon size just after the AD field starts rotation, such that $H^{-1} \sim \omega^{-1} \sim \varphi_0/m_\phi^2$. The charge of a Q -ball should be given by [351]

$$Q \sim H^{-D} n_\varphi \sim \omega^{-D} \omega \varphi_0^2 \sim m_\phi^{3-D} \left(\frac{\varphi_0}{m_\phi} \right)^{1+D}, \quad (355)$$

where $D = 1, 2, 3$ stands for the number of spatial dimensions, and $\omega \approx m_\phi$. This naive expectation has been verified numerically on lattice by Kasuya and Kawasaki [351], although the formation turns out to be slightly delayed [350]. The maximum charge of a thick wall Q -ball can be written as

$$Q_{max} = \beta_D \varphi_0^{1+D}, \quad (356)$$

where β_D 's are some numerical factors with $\beta_1 \approx 0.1$, $\beta_2 \approx 0.02$, and $\beta_3 \approx 6 \times 10^{-4}$ [351].

Kasuya and Kawasaki [351] also noticed that the charge of a Q -ball depends on the helical motion of the AD condensate, and it is proportional to ε

$$\varepsilon = \frac{n_\varphi(t_{osc})}{n_Q^{max}(t_{osc})} = \frac{m_{3/2} \varphi^2}{\omega \varphi^2} = \frac{m_{3/2}}{\omega} \approx \frac{m_{3/2}}{m_\phi}, \quad (357)$$

where $\varepsilon = 1$ corresponds to a circular motion, and $\varepsilon = 0$ for the radial motion. The numerical calculation in [350, 351] indicates that Q_{max} is constant for small ε where

both positive and negative Q -balls with charges of the same order of magnitude are produced, while linearly dependent on ε , around $\varepsilon \sim 1$. It was noticed that dominantly positive Q balls were formed. Numerical simulations also reveal the presence of small negatively charged Q balls [350, 351].

6.4.2 In gauge mediated case

The potential for the condensate Φ forming a Q -ball is given by Eq. (204). If the AD condensate couples directly to particles in a thermal bath, the potential receives a thermal mass correction of order $\sim T^2|\Phi|^2$ while integrating out the heavy modes yields a contribution $\sim \pm T^4 \log(|\Phi|^2/T^2)$. In the latter case the actual sign depends upon the integrated modes. If the integrated heavy modes mainly belong to the matter multiplet then the sign comes out to be positive, otherwise if it were dominated by the heavy gauge degrees of freedom then it turns out to be negative. In a particular case of LH_u flat direction this sign is positive as shown by Fujii, Hamaguchi and Yanagida in [343]. Combining these gives the relevant part of the effective potential (without non-renormalizable terms) as [351]

$$V(\Phi) \approx m_\phi^4(T) \log \left(1 + \frac{|\Phi|^2}{m_\phi^2(T)} \right). \quad (358)$$

Here we have assumed that the flat direction obtains a positive contribution to the thermal potential. The effective mass can be written as [351]

$$m_\phi(T) = \begin{cases} m_\phi & (T < m_\phi) \\ T & (T > m_\phi) \end{cases}, \quad (359)$$

Note that $T \propto t^{-1/4}$ during inflaton oscillations dominated phase.

At very large amplitudes of the AD condensate, the gravity mediation effects for supersymmetry breaking dominates and a stable Q -ball of hybrid type, as discussed in Sect. 6.2, can form [349, 351]. In this case the AD potential is dominated by the terms

$$V(\Phi) \approx m_{3/2}^2 \left[1 + K \log \left(\frac{|\Phi|^2}{M^2} \right) \right] |\Phi|^2. \quad (360)$$

Since here the curvature of the potential depends weakly on the amplitude, the AD condensate starts rotating when $H \simeq m_{3/2} \ll 1$ TeV. In [351], the authors have

simulated the dynamics of the AD condensate on 1, 2, 3-d lattices, and verified the formation of Q balls for $K = -0.01$.

In a cosmological context one may estimate the maximum charge of a Q-ball from the fragmentation of the AD condensate. The analysis is similar to the gravity mediated case, except for the fact that the times when the AD condensate starts oscillating and the most amplified mode (or instability band) enters the horizon do not coincide. The charge of a Q-ball is then given by [349, 351]

$$Q \sim H_f^{-D} m_\phi \varphi_f^2 \sim (|K|^{1/2} m_\phi)^{-D} m_{3/2} |K| \varphi_0^2 \sim |K|^{1-D/2} m_\phi^{3-D} \left(\frac{\varphi_0}{m_\phi} \right)^2, \quad (361)$$

where the subscript ‘ f ’ denotes the time when the Q-ball forms, and we have assumed $m_{3/2} \sim m_\phi$ in our final expression. Note that $Q \propto \varphi_0^2$, with the proportionality constant to be determined numerically; one finds $Q_{max} \sim \tilde{\beta}_3 \varphi^2$ [351], where $\tilde{\beta}_3 \approx 6 \times 10^{-3}$.

6.5 Q-ball collisions

The dynamics of any extended object carrying charge is quite different from the dynamics of charged point-like objects. In this respect studying the Q-ball collision is important in order to understand the charge distribution of Q-matter in the Universe. Q-ball collisions have been studied by a number of authors [392, 151, 350, 393, 394, 395, 396]. In [351, 395], collisions have been considered in the context of gravity mediated supersymmetry breaking, while in [396], Q-ball collisions have been tackled within gauge mediated supersymmetry breaking. An animation of Q-ball collisions can be found at www.utu.fi/~tuomul/collision.mpg.

The usual ansatz is to take initially two spatially well separated Q-balls so that initial field configuration (in 1d) is [394]

$$\Phi(t, x) = e^{i\omega_1 t + i\alpha} \varphi_{\omega_1}(|x + a|) + e^{i\omega_2 t} \phi_{\omega_2}(|x - a|). \quad (362)$$

The two Q-balls have the profiles φ_{ω_1} and φ_{ω_2} , correspondingly to the frequencies ω_1 and ω_2 , and they are separated by a distance $2a$. The total charge of the configuration

(in 1d) is given by [394]

$$Q = Q_{\omega_1} + Q_{\omega_2} + (\omega_1 + \omega_2) \cos(\alpha) \int_{-\infty}^{+\infty} \varphi_{\omega_1}(|x + a|) \varphi_{\omega_2}(|x - a|) dx. \quad (363)$$

Note that the last term is exponentially small in the separation parameter a , because the profiles of the individual Q -balls die off exponentially outside Q -matter. Usually, for a single Q -ball the phase is unimportant because of the global $U(1)$ symmetry, but for multi- Q -ball case the relative phase α plays an important role and affects the total charge of the configuration.

There are two extreme cases: the two charges are equal; or they have opposite sign. The key parameters are the relative phase, the incident velocity, and the charge [393, 394]. The generic interaction for two Q -balls of equal charges is attractive if the relative phase $\alpha = 0$, and repulsive when $\alpha = \pi$. In case of attraction the two Q -balls coalesce to form one larger Q -ball with a resultant charge less than the sum total of the individual charges. A loss in charge also occurs when the Q -ball suffers a large distortion.

If the initial phase $\alpha \neq 0, \pi$ or if the charges of the Q -balls are not equal, then the dynamics of the Q -ball collisions result in charge transfer. Q -balls tend to repel each other, which happens even after charge transfer. If the incident velocity is extremely high (relativistic), then Q -balls simply pass through each other without losing much charge [394, 393].

A collision of a Q -ball and an anti- Q -ball exhibits several interesting features. A naive expectation would have them annihilating. Instead, they bounce back or pass through each other. Charge is partially annihilated, though. The main reason is the fact that generically Q -balls can transfer their charges only very slowly. The charge transfer is very seldom complete [394].

These conclusions hold mainly for thin-wall Q -balls. There is not much difference between gauge and gravity mediated cases. Because in gravity mediation the Q -ball size is smaller, in a fixed volume the AD condensate tends to break into larger number of Q -balls [350]. As a result the Q -balls can have larger peculiar velocities than in the gauge mediated case. Multamäki and Vilja have studied the gravity mediated case in [395] for 2-d lattice for a range of velocities between $v = 10^{-3}$ and $v = 10^{-2}$.

The authors allowed all possible values of the relative phase and found that the fusion cross section of two Q -balls appears to be smaller than the geometric cross section, whereas the cross section for the charge exchange is larger than the geometric cross section. The probability for charge exchange processes increases with increasing ω . The actual velocity of the Q -ball is an open issue, which however, to some extent, can be determined in a cosmological context [350]. In the gauge mediated case the peculiar velocity of the Q -balls is small and the main interactions are either elastic scattering or partial charge exchange [396].

6.6 Q -balls in a thermal bath

There are several effects which one must take into account when Q -balls are immersed in a thermal bath at a temperature higher than the Q -ball formation scale. The temperature dependent effective potential might not even allow for a Q -ball solution, but if it does, then for a large Q thermal corrections are negligible as argued by Kusenko and Shaposhnikov [151], and Laine and Shaposhnikov [397]. Finite temperature effects always lead to an erosion of the condensate and therefore to a loss of charge from the Q -ball into the ambient plasma. As a result a chemical potential $\mu_{\text{plasma}} \sim \delta Q/(VT^2)$ arises in the surrounding plasma, where V is the Hubble volume and T is the ambient temperature. As pointed out in [397], the process of Q -ball evaporation will stop when the chemical potential $\mu_Q \sim m_\phi(T)(Q - \delta Q)^{-1/4}$ associated with the Q -ball becomes equal to μ_{plasma} . The conclusion is strictly valid for the flat potential studied in [397], but should hold qualitatively for all kinds of Q -balls.

In most cases of interest Q -balls are produced non-adiabatically with a formation time scale much shorter than the evaporation scale. Then it is natural to ask how Q -balls come in thermal and chemical equilibrium. Since the Q -ball does not lose its charge significantly, chemical equilibrium between Q -balls and thermal plasma may never be reached, in which case μ_Q always dominates over μ_{plasma} . It is possible to obtain a thermal equilibrium at least between the soft edge of the Q -ball and the hot plasma.

In general there are three different thermally induced effects: dissociation, diffusion,

and evaporation of the Q -ball.

6.6.1 Dissociation

If Q -balls never reach thermal equilibrium with the plasma, then Q -ball dissociation by the bombardment of thermal particles as discussed in [151, 49] is important. In case of charge dissociation, the thermal particles in the plasma collide with the Q -ball and may even penetrate inside. The penetration width depends on the kinetic energy of the particles. For definiteness, let us focus on the gauge mediated case with thick wall Q -balls. A particle ψ which interacts with φ receives a mass contribution $m_\psi \sim g\langle\varphi(x)\rangle$, while outside $m_\psi \sim gT$. At an ambient temperature T , the particle cannot penetrate the Q -ball beyond x_{st} , known as stopping radius and determined by $g\varphi(x_{st}) \approx 3T$. If ψ imparts sufficient energy to the Q -ball in order to overcome its binding energy within the dynamical time scale then the Q -ball may simply break up. On the other hand, if the energy is delivered to the Q -ball is below the dissociation limit, the Q -ball will be able to radiate the excess energy away adiabatically and will not dissociate.

The rate of dissociation depends on the flux of the incoming thermal particles $f = (g_*(T)/\pi^2)4\pi x_{st}^2 T^3$, and the energy per thermal particle transferred to the Q -ball $\sim \gamma_T T$, where $\gamma_T \leq 3$. Then the rate of energy imparted to the Q -ball is given by [49]

$$\frac{dE}{dt} = \frac{4g_*(T)\gamma_T T^4 \beta^2 R^2}{\pi}, \quad (364)$$

where $x_{st} = \beta R$ is defined by the Gaussian thick wall profile $\varphi(r) = \varphi(0) \exp(-r^2/R^2)$, and therefore $\beta = \sqrt{\log(g\varphi(0)/3T)}$. In order to evade complete dissociation $\Delta E \ll \Delta m_\phi Q$, where $\Delta m_\phi \approx |K|m_\phi$. The dissociation will not be completed provided the temperature of the thermal bath is given by [49]

$$T \leq \left[\frac{\pi |K|^2}{4g_*(T)k_r \gamma_T \beta^2} \right]^{1/4} m_\phi Q^{1/4}, \quad (365)$$

where the dynamical time scale is assumed to be $\sim k_r/m_\phi$, with $k_r > 1$.

In a realistic case dissociation alone cannot erode the Q -ball completely. The Q -ball will rather come into thermal equilibrium with the ambient plasma. In an expanding

Universe the minimum energy is configured in such a way that the Q -charges are always present in the Universe along with the Q -balls. In this case the energy of the Q -balls decreases as the temperature of the Universe decreases.

6.6.2 Diffusion

Diffusion takes place only through the soft edge of the Q -ball at a distance over which φ does not change much. There are two factors which determine the diffusion rate; firstly, how efficiently the edge of the Q -ball diffuses, and secondly how fast the core of the Q -ball readjusts itself in order to compensate for the loss of charge. At large temperatures the diffusion rate is large as the Q -ball tries to relax into chemical equilibrium with a thermal plasma. The net diffusion rate is given by [398, 349, 351]

$$\Gamma_{diff} \equiv \frac{dQ}{dt} \sim -4\pi D R_Q \mu_Q T^2 \sim -4\pi A T, \quad (366)$$

where $D = A/T$ is the diffusion coefficient with $A = \mathcal{O}(1)$, and $\mu_Q \sim \omega$ is the chemical potential of a Q -ball. When the temperature of a thermal bath drops due to expansion of the Universe, surface evaporation rate takes over.

If there is a thermal bath already prior to reheating, the instantaneous temperature of the plasma would be large: $T \sim (M_P^2 \Gamma_I H)^{1/4} \sim (M_P T_{rh}^2 H)^{1/4}$, where Γ_I is the decay rate of the inflaton field. During this period, before reheating, $t \propto T^{-4}$, while after reheating $t \propto T^{-2}$. The diffusion rate can be obtained from Eq. (366) [351]

$$\left(\frac{dQ}{dT}\right)_{diff} \sim \begin{cases} 10 \frac{M_P T_{rh}^2}{T^4} & (T > T_{rh}), \\ 10 \frac{M_P}{T^2} & (T < T_{rh}). \end{cases} \quad (367)$$

6.6.3 Evaporation at finite T

In a thermal bath the surface evaporation rate is no longer given by Eq. (339), since one must take into account of thermal corrections. At finite T the evaporation rate of a Q -ball has been found to be [397, 349, 351]

$$\Gamma_{evap} \equiv \frac{dQ}{dt} = -\zeta(\mu_Q - \mu_{plasma}) T^2 4\pi R_Q^2 \sim e\pi\zeta \frac{T^2}{m(T)} Q^{1/4}, \quad (368)$$

where $\mu_{plasma} \ll \mu_Q \sim \omega \sim m(T)Q^{1/4}$, while ζ and $m(T)$ are given by

$$m(T) = \begin{cases} m_\phi & , \\ T & \end{cases}, \quad \zeta = \begin{cases} \left(\frac{T}{m_\phi}\right)^2 & (T < m_\phi), \\ 1 & (T > m_\phi). \end{cases} \quad (369)$$

The evaporation rate can be calculated as [351]

$$\Gamma_{evap} = \frac{dQ}{dt} = \begin{cases} -4\pi T Q^{1/4} & (T > m_\phi), \\ -4\pi \frac{T^4}{m_\phi^3} Q^{1/4} & (T < m_\phi). \end{cases} \quad (370)$$

In order to determine which rate is dominating, let us consider the ratio $R_{diff} \equiv \Gamma_{diff}/\Gamma_{evap}$. For $T > m_\phi$, the ratio is given by $R_{diff} = A Q^{-1/4}$. If $R_{diff} < 1$, the diffusion rate is the bottle-neck for the charge transfer. This condition is met when the Q -ball charge is large enough:

$$Q > 10^2 \left(\frac{A}{4}\right)^2. \quad (371)$$

On the other hand when $T < m_\phi$, the condition $R_{diff} < 1$ corresponds to

$$T > T_* \equiv A^{1/3} m_\phi Q^{-1/12}, \quad (372)$$

and the transition temperature T_* is lower than m_ϕ for large enough Q -ball charge.

If there exists a thermal bath prior to reheating then there are additional complications regarding the evaporation rate, which now exhibits four different possibilities depending on how the temperature compares with the reheating temperature and the mass of the AD particle [351, 49]. The time-temperature relationship changes at $T = T_{rh}$, while the rate dQ/dt changes at $T = m_\phi$. Combining all the effects, one obtains [351]

$$\left(\frac{dQ}{dT}\right)_{evap} \sim \begin{cases} 10 \frac{M_P T_{rh}^2}{T^4} Q^{1/4} & (T > T_{rh}, m_\phi), \\ 10 \frac{M_P T_{rh}^2}{m_\phi^3 T} Q^{1/4} & (T_{rh} < T < m_\phi), \\ 10 \frac{M_P}{T^2} Q^{1/4} & (m_\phi < T < T_{rh}), \\ 10 \frac{M_P T}{m_\phi^3} Q^{1/4} & (T < m_\phi, T_{rh}). \end{cases} \quad (373)$$

The estimate for the loss charge turns out to be of similar magnitude in all possible regimes with [351]

$$\Delta Q \sim 10 \frac{M_{\text{P}}}{m_{\phi}} Q^{1/12} \sim 2.4 \times 10^{18} \left(\frac{m_{\phi}}{\text{TeV}} \right)^{-1} \left(\frac{Q}{10^{24}} \right)^{1/12}, \quad (374)$$

for any case.

In the gravity mediated hybrid case the evaporation and diffusion rates have the same forms in terms of Q -ball parameters $R_Q \sim |K|^{-1/2} m_{3/2}$, and $\omega \sim m_{3/2}$. The transition temperature at which $\Gamma_{\text{evap}} = \Gamma_{\text{diff}}$, reads $T_* \equiv A^{1/3} |K|^{1/6} (m_{3/2} m_{\phi}^2)^{1/3}$. As in the 'usual' type of Q balls where the potential is dominated by the logarithmic term, the charge evaporation near T_* is dominant and the total evaporated charge is found to be [349, 351]

$$\Delta Q \sim 10^{20} \left(\frac{m_{3/2}}{\text{MeV}} \right)^{-1/3} \left(\frac{m_{\phi}}{\text{TeV}} \right)^{-2/3}. \quad (375)$$

7 Cosmological consequences of Q -balls

In this section we discuss various issues concerning Q -ball cosmology. If the MSSM flat direction carries some combination of B and/or L , the charge will be stored in Q -balls created by the fragmentation of the initial condensate. If Q -balls are stable, like in the case of gauge mediated supersymmetry breaking, then they will be a good candidate for dark matter which can be searched directly. If Q -balls eventually decay, the charge will be released in a form of baryonic quanta, providing an interesting alternative mechanism for baryogenesis which does not necessarily depends on sphalerons transitions. The evaporation of Q -ball also gives rise to supersymmetric dark matter. This is an added advantage of Q -ball cosmology: it provides a physical mechanism for relating the dark matter and the baryon densities.

7.1 L -ball cosmology

Q -balls only carrying a leptonic charge are known as L -balls. They emerge from the LH_u flat direction which is quite different from the rest of the MSSM flat directions. Even though LH_u flat direction might not fragment as already mentioned in Sect. 5.3.1., there are choices for the initial conditions at M_{GUT} which make it possible to obtain a decreasing mass with decreasing μ in the RG equations given in Eq. (201). It has been noticed that $m_{LH_u}^2$ becomes negative for scales typically smaller than 10^8 GeV or so [48, 49, 347]. Depending on the choice of parameters there can be a "hill" in the plot of $m_{LH_u}^2$ versus $|\varphi|$ (representing LH_u), such that m_ϕ^2 starts decreasing with increasing $|\varphi|$ for sufficiently large values of $|\varphi|$. The effect of negative m_ϕ^2 at small enough $|\varphi|$ will generate a minimum for $U(|\phi|)/|\phi|^2$ as required for L -ball formation, typically at $|\varphi_0| \approx 1$ TeV. Effectively such a potential can be given by [49]

$$U(\phi) \approx \frac{m_\phi^2}{2}(2e^{-s\varphi} - 1)\varphi^2, \quad (376)$$

where $s \approx 1 \text{ TeV}^{-1}$. This gives rise to thick wall L -balls with radius $R \approx m_\phi^{-1}$. The charge of a L -ball is bounded, i.e. $L \lesssim m_\phi s^{-2} V$, which tends to become zero once $r \gtrsim m_\phi^{-1}$. Since $L \lesssim (sm_\phi)^{-2}$, the L -balls will have a maximum charge, which for

typical values of s and m_ϕ cannot be larger than 10^3 (with an essentially fixed radius). This stands in contrast to other thin wall Q -balls for which the charge is proportional to the volume.

Inside L -balls the field strength is of order TeV, which is much smaller than the initial amplitude for $d = 4$ AD condensate at $H \approx m_\phi$. This suggests that L -balls cannot form by the collapse of an unstable condensate at $H \approx m_\phi$. Therefore the AD baryogenesis along the LH_u direction will be essentially unaltered from the conventional scenario. Even if there were a primordial formation of L -balls, these objects would decay at $T_d \approx 10^7$ GeV. It is therefore unlikely that such L -balls could have any cosmological consequences. Though it is possible that L -balls, which have a field strength of order 1 TeV or less, could play a role in the physics of the electroweak phase transition (we will discuss about phase transition aided by solitons in Sect. 8.3.1).

7.2 B -ball cosmology

Apart from LH_u flat direction, there are purely baryonic directions such as $\bar{u}\bar{d}\bar{d}$. Q -ball forming along this direction carries only baryonic charge and is dubbed as B -ball. There are also the $\bar{d}QL$ and $\bar{e}LL$ directions. One may expect these directions to be phenomenologically similar to $\bar{u}\bar{d}\bar{d}$ direction. R -parity conservation allows $d = 4$ non-renormalizable superpotential term $(H_u L)^2$ and $d = 6$ term $(\bar{u}\bar{d}\bar{d})^2$. In addition, there is also the $d = 4$, $B - L$ conserving $\bar{u}\bar{u}\bar{d}\bar{e}$ direction (and phenomenologically similar $QQQL$ direction). Although this will not produce a B asymmetry via a direct decay of the AD condensate in case there is subsequent anomalous $B + L$ violation (i.e. sphalerons, see Sect. 2.3.4.), but it can generate a baryon asymmetry via Q -ball decays occurring after the electroweak phase transition [48, 49, 347].

For large enough B , a B -ball cannot decay into the lightest B -carrying fermions (the nucleons), and so it is completely stable. Stable B -balls could have a wide ranging astrophysical [151, 364] and experimental implications [331], as will be discussed in Sects. 7.4 and 7.6.

In the gravity mediated case the B -balls may decay at temperatures below T_{ew} , whence the observed baryon number will be a combination of baryon number originat-

ing from the decay of the B -balls and baryon number from free squarks left over after the break-up of the squark condensate. Since the B -balls are composed of squarks, when they decay they will naturally produce a *number* density of neutralinos which is of the same order of magnitude as the number density of baryons, as was first pointed out in [48]. If the B -balls decay sufficiently below the freeze-out temperature of LSPs, and if the number density of thermal relic neutralinos is less than that from the B -ball decay, then the dark matter density and baryon number densities in the Universe will be naturally related.

The actual ratio of baryons to dark matter will essentially be determined by two variables: (i) the mass of the neutralino LSP, and (ii) the proportion of baryon number trapped in B -balls as compared to baryon number in free squarks; this is usually referred to as the efficiency of B -ball formation.

When a B -ball decays, for each unit of B produced, corresponding to the decay of 3 squarks to quarks, there will be at least three units of R -parity produced, corresponding to at least 3 neutralino LSPs (depending on the nature of the cascade produced by the squark decay and the LSP mass, more LSP pairs could be produced). Let $N_\chi \gtrsim 3$ be the number of LSPs produced per baryon number and f_B be the fraction of the total B asymmetry contained in B -balls. Then the baryon to dark matter ratio is given by [48, 49], see also [399],

$$r_B = \frac{\rho_B}{\rho_{DM}} = \frac{m_n}{N_\chi f_B m_\chi} , \quad (377)$$

where m_n is the nucleon mass and m_χ is the neutralino LSP mass. It is rather natural to have $r_B < 1$. The present LEP lower bound on the neutralino mass in the MSSM (assuming no constraints on the scalar masses) is 17 GeV [400]. If we were to assume radiative electroweak symmetry breaking and universal masses for the squarks and Higgs scalars at the unification scale, then the lower bound would become $m_\chi \gtrsim 40$ GeV for $\tan\beta \lesssim 3$ [400]. For $N_\chi \geq 3$, and with $m_\chi \gtrsim 17$ (40) GeV, we find that $r_B < 1$ occurs for $f_B \gtrsim 0.02$ (0.008). As long as more than 2% of the baryon asymmetry is trapped in B -balls, the observed dominance of dark matter in the Universe can be naturally explained.

Primordial nucleosynthesis [3] bounds the density of baryons in the Universe to satisfy $0.0048 \lesssim \Omega_B h^2 \lesssim 0.013$, where $0.4 < h < 1$ (we adopt the bound based on "reasonable" limits on primordial element abundances [3, 57]). The observed baryon to dark matter ratio, $r_B \approx \Omega_B/(1 - \Omega_B)$ (assuming a flat Universe), satisfies $0.005 \lesssim r_B \lesssim 0.09$. This can be accounted from B -ball baryogenesis, provided [48, 49]

$$3.7 \text{ GeV} \lesssim \left(\frac{N_\chi}{3} \right) f_B m_\chi \lesssim 67 \text{ GeV} . \quad (378)$$

For example, if the LSP mass satisfies $17 \text{ (40) GeV} \lesssim m_\chi \lesssim 500 \text{ GeV}$, then the observed baryon to dark matter ratio can be achieved by a wide range of f_B , i.e. $0.007 \lesssim f_B(N_\chi/3) \lesssim 3.9 \text{ (1.7)}$. Note that if $m_\chi \gtrsim 67 \text{ GeV}$, then we must have $f_B < 1$, implying that the observed baryon asymmetry must come from a *mixture* of decaying B -balls and free baryons.

This all assumes that the asymmetry not trapped in the B -balls can survive down to temperatures below T_{ew} . If we were to consider a $B - L$ conserving condensate or additional L violating interactions in thermal equilibrium above T_{ew} , then the only B asymmetry which could survive anomalous $B + L$ violation is the one associated with the B -balls. In this case f_B would be effectively equal to 1 (we refer to this case as "pure" B -ball baryogenesis (BBB)), so m_χ would have to be less than 67 GeV.

A crucial assumption in all these is that there is effectively no subsequent annihilation of LSPs coming from B -ball decays.

7.3 B -balls in gravity mediated supersymmetry breaking

The attractive force due to logarithmic radiative correction term in the condensate scalar potential is given by Eq. (200). Of particular interest is the $d = 6 \bar{u}d\bar{d}$ squark direction with a non-renormalizable superpotential term of the form $(\bar{u}d\bar{d})^2$ and the $d = 4 \bar{u}d\bar{e}$ direction, which conserves $B - L$. The magnitude of K is important for numerical estimates. From the 1-loop effective potential [34], for the $\bar{u}d\bar{d}$ direction, the correction due to gauginos with supersymmetry breaking masses M_{susy} is given by

(see earlier discussion in Sect. 5.3.1.) [48, 49]

$$K \approx -\frac{1}{3} \sum_{\alpha, \text{gauginos}} \frac{\alpha_{g_\alpha} M_{susy}^2}{8\pi m_\phi^2}, \quad (379)$$

where the sum is over those gauginos which gain a mass from the condensate scalar φ . The main contribution will come from the three gluinos which gain masses from the squark expectation values. For $\alpha_{g_3} \approx 0.1$ we obtain $|K| \approx 0.004(M_3/m)^2$. Depending on the ratio of the supersymmetry breaking gluino mass to the squark mass, we expect $|K|$ to be typically in the range 0.01 to 0.1 (see discussion in Sect. 5.3.1.).

7.3.1 *B*-ball Baryogenesis

Recall that the perturbations of the AD condensate at some scale λ go non-linear once $t \gtrsim m_s^{-1} \sim m_\phi^{-1}$ (see Sect. 5.10.1), causing the AD condensate to collapse into fragments of size λ and trapping inside a baryon density. The fragments then relax into the state of lowest energy *B*-balls with a charge of order B . The charge of *B*-ball prevents the soliton from further collapsing and hence the perturbations do not grow on length scales smaller than the *B*-ball radius. Once the length scale going non-linear is larger than the final *B*-ball radius, one expects *B*-balls to form quite efficiently.

The time at which a perturbation of scale λ goes non-linear is then given by [49]

$$t \approx \frac{\alpha_k}{2\pi} \left(\frac{2}{|K|} \right)^{1/2} \lambda, \quad (380)$$

where

$$\alpha_k = \log \left(\frac{\phi_i}{\delta\phi_{i\ k}} \right). \quad (381)$$

One finds that $\alpha_k \approx 34$ (44), for $d = 4$ ($d = 6$) directions. In practice *B*-balls will typically turn out to have thick walls with radius $R \approx (|K|^{1/2} m_\phi)^{-1}$. Perturbations on this scale which have the largest possible growth in time H^{-1} will go non-linear at

$$t \approx \frac{10}{|K| m_\phi}, \quad (382)$$

corresponding to $H \approx 0.1|K|m_\phi$. Assuming that the charge asymmetry corresponds to the presently observed baryon asymmetry; $\eta_B \approx 10^{-10}$ prior to reheating, then the

baryon density is given by

$$n_B \approx \left(\frac{\eta_B}{2\pi}\right) \frac{H^2 M_p^2}{T_R}. \quad (383)$$

The charge contained inside a region within a radius $R \approx (|K|^{1/2} m_\phi)^{-1}$ is given by [48, 347, 49]

$$B \approx 10^{15} |K|^{1/2} \left(\frac{\eta_B}{10^{-10}}\right) \left(\frac{10^9 \text{ GeV}}{T_R}\right) \left(\frac{100 \text{ GeV}}{m_\phi}\right). \quad (384)$$

Hence with $|K| \gtrsim 0.01$, we see that B -balls of charge larger than 10^{14} are likely to form.

It is quite natural for the MSSM B -balls to decay after the electroweak phase transition, see Eq. (344). It is therefore possible to generate the observed B asymmetry from the decay of B -balls occurring at relatively low temperatures. This is true even if there are rapid L violating interactions or $B - L$ conservation. In a thermal bath B -balls could lose their charges by the processes discussed in Sect. 6.5. Therefore we naturally have some kind of constraint on the ambient thermal temperature. Combining Eqs. (384,375), one obtains an upper bound on the final reheat temperature of the Universe [49], which is given by

$$T_{\text{rh}} \leq 10^5 |K|^{1/2} \left(\frac{\eta_B}{10^{-10}}\right) \left(\frac{1 \text{ TeV}}{m_\phi}\right)^{1/3} \text{ GeV}. \quad (385)$$

The reheating temperature therefore cannot be larger than 10^5 GeV , in order that the B -balls have enough charge left over for the purposes of baryogenesis.

7.3.2 LSP dark matter from B -ball decay

B -ball formation from even dimensional $d = 4, 6$ operators is a good candidate for generating dark matter in the Universe through B -ball decay into LSPs [48, 49, 361, 362, 360]. When the B -balls decay there will be $N_\chi \gtrsim 3$ LSPs produced per baryon number or equivalently

$$\Omega_\chi \approx 3 f_B \left(\frac{m_\chi}{m_n}\right) \Omega_B, \quad (386)$$

where f_B is the efficiency parameter which denotes the fraction of the total charge stored in the B -ball, which ought to be less than one in a realistic situation, and m_n is the nucleon mass. Considering the bounds on baryon number density from

nucleosynthesis $0.004 \leq \Omega_B h^2 \leq 0.013$ [3], and the conservative bound from CMB $0.004 \leq \Omega_B h^2 \leq 0.023$, one obtains the upper limits on the LSP mass [49]

$$m_\chi \leq (17.6 - 20.8) f_B^{-1} \left(\frac{\Omega_\chi}{0.4} \right) \left(\frac{h}{0.8} \right)^2 \text{ GeV}. \quad (387)$$

The direct experimental lower limit on the neutralino as a LSP comes from ALEPH $m_\chi > 32.2$ GeV [401], which requires $f_B \leq 0.64$. LEP and Tevatron constraints for the universal A -term and gaugino masses lead to $m_\chi > 46$ GeV, and for the universal scalar masses $m_\chi > 51$ GeV, which implies that $f_B < 0.45$ and $f_B < 0.41$, respectively. This discourages the maximally charged condensate hypothesis for a B -ball formation.

If one adopts Eq. (386) as it stands, without taking into account of possible LSP annihilations, then the amount of dark matter density would be given by [49]

$$\begin{aligned} \Omega_\chi|_{\text{no ann}} &= 3 \left(\frac{N_\chi}{3} \right) f_B \left(\frac{m_\chi}{m_n} \right) \Omega_B \\ &\gtrsim 2.6 f_B \times \left(\frac{N_\chi}{3} \right) \left(\frac{m_\chi}{100 \text{ GeV}} \right) \left(\frac{0.7}{h} \right)^2, \end{aligned} \quad (388)$$

where $m_n \simeq 1$ GeV is the nucleon mass, and we have used the bound $\Omega_B h^2 \gtrsim 0.004$. In case of a bino-like LSP, the B -ball formation could be a serious obstacle for the AD baryogenesis.

In this regard a Q -ball with a smaller charge would be beneficial. One option was considered in [380], where the authors proposed a gauged $B - L$ symmetry to make Q -ball small enough (for a discussion on Q -balls with a local gauge symmetry, see 6.2.6). The gauged $U(1)_{B-L}$ was assumed to be broken at a scale $\sim 10^{14}$ GeV. It was argued that in the gravity-mediated models the D-term from $U(1)_{B-L}$ helps forming smaller B -balls from the oscillations of the flat directions at weak scale. As noticed by the authors [380], their mechanism fails to ameliorate the problem for the gauge-mediated models. Another solution has been invoked in [402], where it was argued that if one takes into account of the Hubble induced radiative corrections to the flat direction then for a range of gaugino masses $3H \lesssim m_{1/2} \lesssim 5H$, the amplitude of the AD condensate oscillations is redshifted and leads to a formation of considerably smaller Q -balls at low scales. The advantage is that the mechanism works for both gravity and gauge mediated type AD potentials.

Otherwise, because of a large annihilation cross section, Wino and Higgsino like LSP can be rather more promising candidates for CDM if ultimately originating from an AD condensate, as pointed out by Fujii and Hamaguchi in [361, 362]. If the LSP is not very heavy, say $m_\chi \lesssim 200$ GeV, so that their freeze-out temperature is about $m_\chi/20 \simeq 10$ GeV, then B -balls should decay at temperatures below 10 GeV for all the produced LSPs to survive. It is quite possible that B -balls are not the only source of LSPs and that some of the LSPs produced in B -ball decay will be annihilated with the LSPs in the background. Using Eq. (384), one finds that if the B -ball decay temperature $T_d \leq 10$ GeV, there is an upper limit on the reheating temperature [49]

$$T_{rh} \lesssim 5 \times 10^8 \frac{|K|^{3/2}}{f_s} \left(\frac{100 \text{ GeV}}{m_\phi} \right)^2 \left(\frac{40}{\alpha_k} \right)^2 \left(\frac{T_d}{10 \text{ GeV}} \right)^2 \left(\frac{\eta_B}{10^{-10}} \right) \text{ GeV} . \quad (389)$$

With $|K| \approx (0.01 - 0.1)$, and $\alpha_k \sim 40$, $T_{rh} \lesssim (5 \times 10^5 - 2 \times 10^7) f_s^{-1} \text{ GeV}$.

7.3.3 The LSP abundance

The LSPs produced in B -ball decays will collide with themselves and with other weakly interacting particles in the background and settle locally into a kinetic equilibrium. Thermal contact can be maintained until $T_f \sim m_\chi/20$ [285, 18], and a rough freeze-out condition for LSPs (if they were initially in thermal equilibrium) will be given by [18]

$$n_{\text{LSP}} \langle \sigma_{\text{ann}} v \rangle \approx H_f \frac{m_\chi}{T_f} , \quad (390)$$

where σ_{ann} is the LSP annihilation cross-section and the subscript f refers to the freeze-out values. The thermally averaged cross section can be written as $\langle \sigma_{\text{ann}} v \rangle = a + bT/m_\chi$, where a and b depend on the couplings and the masses of the light fermions [18]. For a light neutralino with $m_\chi < m_W$, neglecting the final state fermion masses and assuming an efficient LSP production, so that $f_B = 1$, one finds for the LSP density from Eq. (390), that $b \approx H m_\chi^2 T_f^{-2} n_f^{-1}$, and

$$n_f = \frac{1}{(2\pi)^{3/2}} (m_\chi T_f)^{3/2} e^{-m_\chi/T_f} \approx 1.46 \times 10^{-12} m_\chi^3 . \quad (391)$$

The LSPs produced in B -ball decays will spread out by a random walk with a rate ν determined by the collision frequency divided by a thermal velocity $v_{th} \approx \sqrt{T/m_\chi}$.

Since the decay is spherically symmetric, it is very likely that the LSPs have a Gaussian distribution around the central region of radius as pointed out in [48]

$$\bar{r} \approx \left(\frac{\nu x}{\Gamma_B} \right)^{1/2} \quad (392)$$

Within this central region annihilation is significant if $\bar{n}_{\text{LSP}} \langle \sigma_{\text{ann}} v_{\text{rel}} \rangle \gtrsim H$. In [48], it was shown that the annihilation of LSPs is insignificant provided the Q -ball decay temperature is given by

$$T_d \ll 21 \left(\frac{m_\chi}{100 \text{ GeV}} \right)^{3/16} \left(\frac{10^{20}}{N_{\text{LSP}}^{\text{tot}}} \right)^{1/8} \left(\frac{100}{g(T)} \right)^{3/16} \text{ GeV} . \quad (393)$$

The main conclusion is that typically most of the LSPs will survive if the B -ball decay temperature is less than a few GeVs.

The final abundance of the LSPs can be approximately expressed by a simple analytical form [361, 362]. Solving the Boltzmann equation analytically for the LSP, one finds [362]

$$Y_\chi(T) \simeq \left[\frac{1}{Y_\chi(T_d)} + \sqrt{\frac{8\pi^2 g_*(T_d)}{45}} \langle \sigma v \rangle M_{pl}(T_d - T) \right]^{-1}, \quad (394)$$

where $Y_\chi = n_\chi/s$. In Fig. (11), we present the numerical solution of Fujii and Hamaguchi for two cases: with and without large entropy production from the B -balls. In the latter case n_Q need not be directly related to the present baryon asymmetry. This case can be simulated by choosing a very small ε and including the radiation energy density generated by the B -ball decay in the Boltzmann equation [362].

If initial abundance $Y_\chi(T_d)$ is large enough, the final abundance $Y_{\chi 0}$ for $T \ll T_d$ is given by [361, 362]

$$Y_{\chi 0} \simeq Y_\chi^{\text{approx}} \equiv \left[\sqrt{\frac{8\pi^2 g_*(T_d)}{45}} \langle \sigma v \rangle M_{pl} T_d \right]^{-1}. \quad (395)$$

In this case the final abundance $Y_{\chi 0}$ is determined only by the Q -ball decay temperature T_d and the annihilation cross section of the LSP $\langle \sigma v \rangle$ (independently of the initial value $Y_\chi(T_d)$ as long as $Y_\chi(T_d) \gg Y_\chi^{\text{approx}}$). In terms of the density parameter Ω_χ , the neutralino abundance can be rewritten as [362]

$$\Omega_\chi \simeq 0.5 \left(\frac{0.7}{h} \right)^2 \times \left(\frac{m_\chi}{100 \text{ GeV}} \right) \left(\frac{10^{-7} \text{ GeV}^2}{\langle \sigma v \rangle} \right) \times \left(\frac{100 \text{ MeV}}{T_d} \right) \left(\frac{10}{g_*(T_d)} \right)^{1/2}. \quad (396)$$




Figure 11: The evolution of the neutralino dark matter abundance generated from the Q -ball decay for $T_d = 1$ GeV and $\langle\sigma v\rangle = 10^{-9}$, 10^{-7} , and 10^{-5}GeV^{-2} , represented by thick solid lines. The abundances estimated from the analytic formula in Eq. (395) are shown in dashed lines. In the first figure the energy density of the Q -balls has been assumed to be small enough with respect to radiation; the parameter values are $m_\varphi = 1$ TeV, $m_\chi = 100$ GeV, $\epsilon = 0.1$ and $N_\chi = 10$. In the second plot Q -balls have been assumed to dominate the energy density before their decay; here the parameters are $m_\varphi = 1$ TeV, $m_\chi = 100$ GeV, $\epsilon < 0.01$ and $N_\chi = 100$, from [362].

In case of $Y_\chi(T_d) < Y_\chi^{\text{approx}}$, the final abundance is given by

$$Y_{\chi 0} \simeq Y_\chi(T_d) \gtrsim \varepsilon^{-1} \left(\frac{n_B}{s} \right)_0, \quad (397)$$

This is the case where the LSP annihilation cross section is small enough, which holds for a bino-like LSP, where the relic abundance of the LSPs is directly related to the observed baryon asymmetry [48]. Unfortunately, a bino-like neutralino will overclose the Universe unless we assume an extremely light bino (which is experimentally excluded).

7.3.4 Which direction?

Consider the $\bar{u}d\bar{d}$ and $\bar{u}\bar{u}d\bar{d}\bar{e}$ directions, which are both lifted by $d = 4$ and $d = 6$ non-renormalizable operators. For $d = 6$ case the initial value of the field when the condensate oscillations begin is given by $\varphi_0 = 5.8 \times 10^{14} \lambda^{-1/4} (m_\varphi/100 \text{ GeV})^{1/4} \text{ GeV}$, and for $d = 4$ case $\varphi_0 = 3.2 \times 10^{10} \lambda^{-1/2} (m_\varphi/100 \text{ GeV})^{1/2} \text{ GeV}$. Let us compare the

lower bound on the reheating temperature for the two cases. For $d = 6$, it has been estimated [49] to be $T_{rh} \gtrsim 0.23 \lambda^{1/2} (m_\phi/100 \text{ GeV})^{1/2} \text{ GeV}$ (for $d = 6$, $\lambda \sim 0.003$ if the strength of the non-renormalizable interactions is set by M_P). Such bound on T_{rh} fulfills all the requirements for the survival of B -balls from thermal dissociation, diffusion, etc. Repeating the same analysis for the $d = 4$ case leads to an upper bound $T_{rh} \gtrsim 8 \times 10^7 \lambda \text{ GeV}$ (with $\lambda \sim 0.1$). This is rather hard to satisfy. The robust conclusion appears to be that for an efficient B -ball production which also gives rise to dark matter, should involve the $d = 6$ $\bar{u}d\bar{d}$ direction. This particular direction is also favored in the sense that $f_s = 1$ can be possible. As long as the reheating temperature does not exceed 10^{3-5} GeV , the baryon to dark matter ratio can be naturally accounted by B -ball decays in case of the MSSM flat directions which are lifted by $d = 6$ operators [48].

In the case of the $d = 4$ directions B -balls do not form efficiently [49]. For $d = 4$ and along the $\bar{u}d\bar{d}$ direction $B - L$ is actually conserved. In this case only the baryon number trapped in the B -balls will survive. Even though it might appear that the observed B asymmetry could be obtained if the initial condensate had a large B , this would be a rather ambitious program because $\bar{u}d\bar{d}$ -balls disappear rather quickly in a thermal bath as argued in [48].

7.3.5 Direct LSP searches and B -balls

If dark matter is a neutralino, then it should be possible to confirm its existence by direct and indirect searches. The direct detection involves the interaction of neutralino with matter, which is usually dominated by scalar couplings of relatively heavy nuclei $A \gtrsim 20$ [403, 18]. The counting rate of the elastic neutralino-nucleon scattering is given by [404]

$$R = \left(\frac{\sigma \xi}{m_\chi m_N} \right) \left(\frac{1.8 \times 10^{11} \text{ GeV}^4}{\text{Kg} \cdot \text{days}} \right) \left(\frac{\rho_\chi}{0.3 \text{ GeV/cm}^3} \right) \left(\frac{v_\chi}{320 \text{ Km/s}} \right) \text{ events}, \quad (398)$$

where ρ_χ is the density, v_χ the average velocity of the neutralino, m_n the mass of the target nucleon, and ξ a nuclear form factor. The cross section σ is the neutralino-nucleon cross-section at zero momentum transfer. These interactions are mediated

by heavy Higgs exchanges, or by a sfermion exchange. The forthcoming experiments, such as CDMS [405], CRESST [406], EDELWEISS II [407], GENIUS [408] and ZEPLIN [409], have high hopes in reaching cross-sections up to 10^{-10} pb.

Indirect detection relies on astrophysics. The annihilation of neutralino LSPs in any astrophysical sources can give rise to fluxes of anti-protons and positrons which are usually not seen in the cosmic rays. Other possible ways are neutralino annihilation into 2γ final states or $Z\gamma$ final states [410]. The monoenergetic gamma rays with energy $\sim m_\chi$ might not have any competitive background from other astrophysical sources, and has a possibility of being detected in the next generation air Cherenkov telescopes observing the galactic center, such as VERITAS [411], HESS [412] and MAGIC [413].

Fujii and Hamaguchi [362], have studied in detail the parameter space where neutralino production from the decays of B -balls gives the correct dark matter abundance. The allowed region is found where the dominant contribution to the LSP is provided by $\widetilde{H}$. A large $\widetilde{H}$ content of the LSP enhances the neutralino annihilation cross section into W bosons via chargino exchange. There is a small difference in thermally produced $\widetilde{H}$ and non-thermally produced ones. In the latter case the mass of the neutralino is much smaller than thermally produced ones. The annihilation rate of neutralinos into 2γ final states is enhanced for $\widetilde{H}$ -like neutralino (this was previously shown in the context of thermally distributed neutralinos in [410, 414]), and seems to hold true even when the neutralinos are created non-thermally [361, 362]. A typical B -ball decay temperature which leads to the desired CDM density is $100 \text{ MeV} \lesssim T_d \lesssim (\text{a few}) \text{ GeV}$. In the anomaly mediation and in the no-scale supersymmetry breaking models, it was found that $\widetilde{W}$ is the most promising candidate for the LSP in a wide region of the parameter space [362]. The conclusion is that if AD baryogenesis is successful within minimal SUGRA, then Higgsino and Wino like neutralinos are perhaps the likely dark matter candidates [362].

7.4 Q -balls and gauge mediated supersymmetry breaking

As pointed out by Kusenko and Shaposhnikov, in gauge mediated supersymmetry breaking scenarios the salient feature is that the Q -balls are stable against decaying into nucleons [151], because the Q -ball energy per unit charge is given by $E_Q/Q \simeq m_\phi Q^{-1/4} < 1$ GeV for $m_\phi \sim 1$ TeV. For sufficiently large Q , the Q -ball itself could then be a candidate for CDM [331, 151, 356, 351, 415].

7.4.1 Baryogenesis and gauge mediation

A sufficiently large Q -ball in the gauge mediated supersymmetry breaking case will be absolutely stable against decaying into nucleons. Although baryogenesis by evaporation of a Q -ball is not effective, Q -balls nevertheless lose some charge to the ambient plasma by thermal dissociation and diffusion, which might create asymmetry in the nucleons. An ambitious possibility, but one that would be hard to realize, is to work out whether Q -ball evaporation could also lead to dark matter as in the case of gravity mediated supersymmetry breaking.

In the gauge mediated case the baryon asymmetry can be related to the dark matter density via

$$\eta_B = \frac{n_B}{n_\gamma} \simeq \frac{\varepsilon n_Q \Delta Q}{n_\gamma} \simeq \frac{\varepsilon \rho_Q \Delta Q}{n_\gamma M_Q} \simeq \frac{\varepsilon \rho_{c,0} \Omega_Q \Delta Q}{n_{\gamma,0} M_Q}, \quad (399)$$

where $\varepsilon \ll 1$ displays the departure from a circular orbit, ΔQ is the evaporated charge, Ω_Q is the density parameter for the Q -balls, $\rho_{c,0} \sim 8h_0^2 \times 10^{-47} \text{ GeV}^4$ is the present critical density, and $n_{\gamma,0} \sim 3.3 \times 10^{-39} \text{ GeV}^3$ is the present photon number density, with $h_0 \sim 0.7$. The evaporated charge should yield the baryons while the remaining charge is in Q -ball dark matter. The baryon-to-dark matter ratio is given by [351]

$$r_B \equiv \frac{\Delta Q}{Q} \sim \eta_B \frac{m_\phi n_{\gamma,0}}{\varepsilon \rho_{c,0} \Omega_Q} Q^{-1/4} \sim 10^{11} \varepsilon^{-1} \eta_B \Omega_Q^{-1} \left(\frac{m_\phi}{\text{TeV}} \right) Q^{-1/4}. \quad (400)$$

The total evaporated charge from Q -balls in a thermal bath is given by Eq. (374). Requiring that $Q > \Delta Q$, the charge of the Q -ball should be sufficiently large [351]:

$$Q \geq 1.2 \times 10^8 \eta_B^{-3/2} \varepsilon^{3/2} \Omega_Q^{3/2} \left(\frac{m_\phi}{\text{TeV}} \right)^{-3}. \quad (401)$$

The value of ε is not an independent quantity and can be related to the charge of a Q -ball. At the beginning of the condensate rotation $H \sim m_\phi^2(T)/\varphi_0 \sim T^2/\varphi_0$, where $T \sim (MT_{rh}^2 H)^{1/4}$ is the instantaneous temperature prior to reheating. Combining these two pieces of information, one obtains $T \sim T_{rh} \sqrt{M/\varphi_0}$. The net charge of the Q -ball is then related to the baryon number density through $n_B \sim r_B \varepsilon n_\varphi \sim r_B \varepsilon \omega \varphi_0^2$, where $\omega \sim m_\phi^2(T)/\varphi_0 \sim T^2/\varphi_0$ is the rotation frequency of the condensate. With the help of Eq. (400), one finds [351]

$$n_B \sim 10^{11} \eta_B \Omega_Q^{-1} \left(\frac{m_\phi}{\text{TeV}} \right) T^2 \varphi_0 Q^{-1/4}, \quad (402)$$

Kasuya and Kawasaki also estimated the net charge of the Q -ball, given by the initial amplitude of the condensate and the reheating temperature, as [351]

$$Q \sim 10^{44} \left(\frac{m_\phi}{\text{TeV}} \right)^4 \frac{\varphi_0^{16}}{T_{rh}^4 M^{12}} \Omega_Q^{-4}. \quad (403)$$

From Eq. (356) one obtains $Q \sim \beta (\varphi_0/T)^4$ for $\varepsilon \sim 1$ (for $\varepsilon \ll 1$, one replaces β by $\beta' = \gamma\beta$ with $\gamma \sim 0.1$).

From all these considerations one can estimate the amplitude of the condensate and the charge of the Q -ball. The amplitude is given by [351]

$$\phi_0 \sim 4.6 \times 10^{13} \varepsilon^{1/10} \Omega_Q^{2/5} \times \left(\frac{\beta}{6 \times 10^{-4}} \right)^{1/10} \left(\frac{m_\phi}{\text{TeV}} \right)^{-2/5} \text{GeV}. \quad (404)$$

Inserting this in the expression $T \sim T_{rh} \sqrt{M/\varphi_0}$, one obtains [351]

$$T \sim 2.3 \times 10^7 \varepsilon^{-1/20} \Omega_Q^{-1/5} \left(\frac{T_{rh}}{10^5 \text{GeV}} \right) \times \left(\frac{\beta}{6 \times 10^{-4}} \right)^{-1/20} \left(\frac{m_\phi}{\text{TeV}} \right)^{1/5} \text{GeV}, \quad (405)$$

so that the charge of the Q -ball reads [351]

$$Q \sim 9.3 \times 10^{21} \varepsilon^{8/5} \Omega_Q^{12/5} \left(\frac{T_{rh}}{10^5 \text{GeV}} \right)^{-4} \times \left(\frac{\beta}{6 \times 10^{-4}} \right)^{8/5} \left(\frac{m_\phi}{\text{TeV}} \right)^{-12/5}. \quad (406)$$

Now, with the help of Eqs. (401) and (406), one obtains [351]

$$\varepsilon \sim 1.5 \times 10^{-2} \Omega_Q^{3/5} \left(\frac{\eta_B}{10^{-10}} \right) \left(\frac{T_{rh}}{10^5 \text{GeV}} \right)^{-8/3} \times \left(\frac{\beta}{6 \times 10^{-5}} \right)^{16/15} \left(\frac{m_\phi}{\text{TeV}} \right)^{2/5}. \quad (407)$$

The above equations determine the parameter space for Q -ball baryogenesis to coexist with Q -ball dark matter in gauge mediated models.

7.4.2 Generic gauge mediated models

Now the task is to consider the parameter space for a realistic MSSM flat direction. A detailed analysis can be found in [351], where it was pointed out that the $d = 5, 6$ directions are favorable. For $d = 4$, the charge does not accumulate enough to survive as a dark matter relic while $d = 7$ requires unnaturally small values of ε . It is worth pointing out that as long as the ambient temperature of the plasma is sufficiently high with $T \geq (m_{3/2}\varphi_0)^{1/2} [\log(\varphi_0^2/T^2)]^{-1/4}$, the logarithmic term dominates over the gravity-mediation term. For $d = 5, 6$, the requirement on temperature is thus $T \geq 10^{5-6}$ GeV.

On the other hand, if the Q -ball forming scalar field has a large vev, or if the reheating temperature is extremely low, then gravity mediation effects should also be taken into account. However, as Kasuya and Kawasaki pointed out in [351], in order to produce enough baryon asymmetry and simultaneously produce surviving Q -balls for dark matter, one requires a considerably large gravitino mass $m_{3/2} \geq 10^3$ GeV, which is an unacceptable value for the gravitino mass within the gauge mediated supersymmetry breaking [39].

One could also consider a generic model for gauge mediation where the scale of the logarithmic potential is larger than m_ϕ , such that [349, 351]

$$V \sim \begin{cases} M_F^4 \log\left(\frac{\varphi^2}{M_S^2}\right) & (\varphi \gg M_S), \\ m_\phi^2 \varphi^2 & (\varphi \ll M_S), \end{cases} \quad (408)$$

where M_S is the messenger mass scale. In this particular case the condensate will start oscillating at large field amplitudes. The Q -ball will form at a large vev. The mass and the size of the Q -ball are now given by [349, 351]

$$M_Q \sim M_F Q^{3/4}, \quad R \sim M_F^{-1} Q^{1/4}, \quad \omega \sim \frac{M_F^2}{\varphi}, \quad (409)$$

If $M_F \gtrsim T$, then the Q -ball could be stable against decaying into the nucleons, provided the Q -ball mass per unit charge is smaller than 1 GeV. This condition holds for

$$Q \gtrsim 10^{24} \left(\frac{M_F}{10^6 \text{ GeV}} \right)^4. \quad (410)$$

The ambient temperature is smaller than M_F , but could be larger than m_ϕ . In this case the Q -ball would rather evaporate. The total evaporated charge can be estimated to be [351]

$$\Delta Q \sim 10^{15} \left(\frac{m_\phi}{10^2 \text{ GeV}} \right)^{-2/3} \left(\frac{M_F}{10^6 \text{ GeV}} \right)^{-1/3} \times Q^{1/12}. \quad (411)$$

The survival condition: $Q \geq \Delta Q$ then imposes a bound on an initially accumulated charge to be

$$Q \geq 10^{17} \left(\frac{m_\phi}{10^2 \text{ GeV}} \right)^{-8/11} \left(\frac{M_F}{10^6 \text{ GeV}} \right)^{-4/11}. \quad (412)$$

While combining with Eq. (399), one obtains a relationship between the baryon number and the amount of dark matter as [351]

$$Q \sim 10^{17} \Omega_Q^{3/2} \varepsilon^{3/2} \left(\frac{\eta_B}{10^{-10}} \right)^{-3/2} \left(\frac{m_\phi}{10^2 \text{ GeV}} \right)^{-1} \left(\frac{M_F}{10^6 \text{ GeV}} \right)^{-2}. \quad (413)$$

The consistency condition requires the initial amplitude of the AD condensate to be $\varphi_0 \sim 10^{11} \text{ GeV}$. Note that when all the above conditions are taken into account, then there is hardly any region in the parameter space which allows for the required baryon asymmetry from evaporation of Q -balls together with enough surviving Q -balls to provide the dark matter, as was concluded by Kasuya and Kawasaki in [351].

7.4.3 Late formation of gauged Q -balls

In [351], the authors have also considered the late formation of Q -ball at a scale when gravity mediation and thermal logarithmic correction terms dominate the potential at large and small scales, respectively. The Q -ball forms when the instability band enters the horizon with a wavelength k_{eq}^{-1} and the angular velocity $\omega_{eq} \sim (T_{eq}^2/\varphi_{eq})^{-1} \sim m_{3/2}^{-1}$. The number density of the Q -balls is found to be $n_{eq} \sim T_{eq}^4/m_{3/2}$, while the charges are $Q \sim (\varphi/T_{eq})^4$ [351]. The exact relationship depends on the helicity of the condensate. In this late formation scenario the only realistic flat directions turn out to be $d = 6$ with $Q \sim 10^{24} - 10^{21}$, $T_{rh} \sim 1.0 \times 10^7 - 30 \text{ GeV}$, and $M_F \sim 10^2 - 10^4 \text{ GeV}$; and $d = 7$ with $Q \sim 10^{25} - 10^{22}$, $T_{rh} \sim 60 \text{ GeV} - 10 \text{ MeV}$, and $M_F \sim 10^2 - 10^3 \text{ GeV}$ [351].

Kasuya and Kawasaki also repeated their analysis when temperature effects are negligible. In this case the Q -ball forms when $V \sim M_F^4 \log(\varphi_{eq}^2/M_s^2) \sim m_{3/2}^2 \varphi_{eq}^2$, where $\varphi_{eq} \sim M_F^2/m_{3/2}$. The AD condensate fragment just after its amplitude becomes

smaller than φ_{eq} . Consistent scenarios arise only in $d = 6, 7, 8$ cases. For the allowed parameter space, one should require $m_{3/2} \sim 0.1$ GeV, $M_F \sim 10^4$ GeV, $T_{rh} \sim 5$ GeV, and $Q \sim 10^{20}$.

These many proposals indicate very clearly that in order to find a consistent cosmological scenario where Q -ball evaporation leads to baryon asymmetry while the survived Q -balls act as a dark matter candidate requires a stringent condition on $M_F \leq 10^6$ GeV. Note that $M_F \sim 10^6$ GeV is required to provide the right spectrum for sparticle masses, see [297, 39].

7.5 Q -balls as self-interacting dark matter

Recently, Q -ball has been proposed as a candidate for self-interacting dark matter by Kusenko and Steinhardt [357]. Such a consideration is motivated by the fact that collisionless CDM appears to have certain discrepancies between numerical simulations and observations. The halo density profiles, and the number density of satellite galaxies, do not match well with the observations [416, 417]. A possible remedy is dark matter that has fairly strong self-interactions [417], a situation which however is not easily achieved in the standard particle physics models.

The self-interaction cross section and the mass of the dark matter particles undergoing elastic scattering should satisfy the relation $s = \sigma_{DD}/m_{DM} \sim 2 \times 10^3 - 3 \times 10^4$ GeV⁻³. This might change if one considers other processes such as dark matter annihilation.

Q -balls, being extended objects, can certainly have a large geometric cross-section, but it was found that in order to match the required cross-section to mass ratio s , the AD particles should have a very low mass scale $\sim \mathcal{O}(1)$ MeV. This requires that the charge of the Q -ball should also be very small, i.e. $Q \leq 10^5$. Such Q -balls would not originate within MSSM but could be possible in some extended theories. There are however a number of issues concerning production of such a small charged Q -balls, their thermal distribution, their evaporation, annihilation and scattering should be taken into account consistently. It has been argued that thick wall Q -balls do not seem to have much admissible parameter space [358], while thin wall Q -balls have a

slightly better chance to succeed with a vev $\varphi_0 \geq \mathcal{O}(\text{MeV})$. In any case, such a small AD condensate amplitude seems hard to obtain, although not completely impossible, in order to reconcile with the existing particle physics models.

Recently, it has been pointed out [418] that future experiments should be able to discern the spatial extent of the dark-matter particle. It was noticed that the extended objects such as Q -balls leave its distinct imprint on the spectrum which falls off very fast with increasing energy. The signal is primarily dominated by the low-energy events near the threshold.

7.6 Direct searches for gauge mediated Q -balls

If the initial charge of a Q -ball is larger than the evaporated charge, the Q -ball survives and contributes to the energy density of the Universe. Let us take as an example B -balls, such as $\bar{u}_2 \bar{d}_1 \bar{d}_2$, which in gauge mediated case are unstable because of the presence of baryon number violating operators (required in the first place to charge up the condensate) with dimensions larger than 5 [415]. Nevertheless for $Q \geq 10^{20}$, the lifetime of the Q -ball is in fact greater than the age of the Universe ($t_0 \sim 10^{10}$ years) [415]. Such Q -balls are potential candidates for CDM.

From Eq.(374), we can read the limit on the initial charge of a Q -ball

$$Q_{\text{init}} \geq 7.4 \times 10^{17} \left(\frac{m_\phi}{\text{TeV}} \right)^{-12/11}. \quad (414)$$

On the other hand, in order for the Q -ball to be stable against decaying into nucleons, i.e. $E_Q/Q \leq 1 \text{ GeV}$, one finds [351]

$$Q \geq 10^{12} \left(\frac{m_\phi}{\text{TeV}} \right)^4. \quad (415)$$

The condition relating baryon number and dark matter is given by Eq.(401), and can be written as

$$Q \leq 10^{23} \varepsilon^{3/2} \left(\frac{m_\phi}{\text{TeV}} \right)^{-3}, \quad (416)$$

where we have taken $\eta_B \sim 10^{-10}$ and $\Omega_Q \leq 1$.

If we do not impose the condition that the evaporated charge accounts for the baryons in the Universe, the only constraint is that the energy density of Q -balls must

not exceed the critical density. As mentioned in the previous Section, this condition is given by Eq.(406).

Let us now assume that the gauge mediated Q -balls indeed make up the dark matter in the galactic halo. Then the corresponding number density is given by [355]

$$n_Q \sim \frac{\rho_{dm}}{M_Q} \sim 5 \times 10^{-5} Q_B^{-3/4} \left(\frac{1 \text{ TeV}}{m_\phi} \right) \text{ cm}^{-3}, \quad (417)$$

and the Q -ball flux will be determined by $\mathcal{F} \sim (1/4\pi)n_Q v \sim 10^2 Q_B^{-3/4} (1 \text{ TeV}/m_\phi) \text{ cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$, where we have taken $v \sim 10^{-3}c$. The number of events will obviously depend on the surface area of the detector. In case of Super-Kamiokande the number of events could be estimated to be $N \sim (10^{24}/Q_B)^{3/4} (1 \text{ TeV}/m_\phi) \text{ yr}^{-1}$ [355], where the surface area of the water tank is $\sim 7.5 \times 10^7 \text{ cm}^2$ [419].

The Q -balls can be detected via Kusenko-Kuzmin-Shaposhnikov-Tinyakov (KKST) process [355]. Note that in the absence of any fundamental singlet a B -ball would necessarily break $SU(3)$ while the electroweak symmetry could be restored inside, provided the Higgses do not obtain vevs or if the flat direction is already a $SU(2)$ singlet such as $\bar{u}\bar{d}\bar{d}$. The instanton mediated baryon number violation will be much weaker in these cases because of the limitation on the size of the instanton, which cannot be larger than the size of the Q -ball. When nucleons collide with a Q -ball, they enter the surface layer of the Q ball, and dissociate into quarks, which are converted into squarks via gluino exchange. In this process, Q -balls release $\sim 1 \text{ GeV}$ energy per collision by emitting soft pions.

For an electrically neutral Q -ball the absorption cross-section is quite large and is determined by the soliton size to be [355] $\sigma_B \sim 10^{-33} Q_B^{1/2} (1 \text{ TeV}/m_\phi)^2 \text{ cm}^2$. The estimated mean free path in matter is given by $\lambda \sim 10^{-3} A (10^{24}/Q_B)^{1/2} (m_\phi/1 \text{ TeV}) (1 \text{ g/cm}^3/\rho) \text{ cm}$ [355, 51], where A is the weight of the atomic nucleus with a density distribution ρ . Obviously absorption of quarks takes place at a higher rate than the collisions of Q -balls with nuclei. The BAIKAL experiment [420] sets a limit on the monopole flux which also gives a lower bound on the charge of a Q -ball as $Q_B \geq 10^{22}$ for $m_\phi \sim 1 \text{ TeV}$.

If the Q -balls are electrically charged, for instance when the selectrons obtain a large vev along the $QQQLLL\bar{e}$ flat direction, the detection prospects worsen, because of two reasons. First, there will be a Coulomb barrier which will prevent absorption




Figure 12: The bounds on the charge of an electrically neutral Q -ball with respect to the supersymmetry breaking scale. Various shaded regions of the parameter space are already ruled out. The line "B-Ball Stability Limit" marks the region below which B -balls are not stable. The allowed region $Q > 10^{22}$ can be found only in the upper part of the figure, from [51].

of the incoming nuclei. Secondly, the absorption cross section will be determined by the Bohr radius $\sigma \sim \pi r_B^2 \sim 10^{-16} \text{ cm}^{-2}$, and therefore the corresponding mean free path length will decrease to $\lambda \sim 10^{-8} A(1 \text{ g/cm}^3/\rho) \text{ cm}$ [355]. The present limit on electrically charged Q -ball comes from the MACRO search [421] with a flux $\mathcal{F} \leq 1.1 \times 10^{-14} \text{ cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$, which constraints the charge of a B -ball to $Q_B \geq 10^{21}$.

A wide range of charges are already ruled out, see Fig. (12) from [51]. The only allowed range of charges is $Q \sim 10^{22} - 10^{26}$ for $m_\phi \sim 1 \text{ TeV} - 100 \text{ GeV}$. The range however depends on ε , which one tacitly assumes to be of order one in Eq. (416). For smaller values of ε the range of allowed charges will further reduce. In a generic logarithmic potential a charge $Q \sim 10^{24}$ and $M_F \sim 10^2 \text{ GeV}$ is still allowed [351]. In many cases such as in delayed Q -ball formation [351], the charge of a Q -ball is large $\sim 10^{26}$. Such Q -balls could be detectable in the Telescope Array Project or the OWL-AIRWATCH detector.

There are also astrophysical ways for detecting stable Q -balls. Charged Q -balls can dissipate a large amount of energy quite efficiently before they can be detected. The Q -balls might leave a track behind whose stopping range is roughly 1000 m

for a charge $Q_B \leq 10^{13}(m_\phi/1 \text{ TeV})^{-4/3}$ [355]. Electrically neutral Q -balls would hardly have any impact when they pass through a planet like ours. The loss in Q -ball kinetic energy would be almost undetectable; the velocity would decrease by $\delta v/v \sim 10^{-2}Q_B^{-1/4}(1 \text{ TeV}/m_\phi)^3$.

Q -balls could also be captured in sufficiently dense stars such as inside the core of neutron stars. This will result in an increase in the temperature of the neutron star studied in [355], but an insignificant enhancement $\sim 0.01(Q_B/10^{24})^{-1/16}$ may not result in any observable consequences. There is still a possibility that the gradual accumulation of Q -balls may seal the fate of a neutron star by decreasing its mass and thereby reaching the critical condition which may lead to a supernovae explosion [355].

8 Flat directions other than MSSM

Flat directions, fragmentation of scalar condensates and Q -balls are generic features that could be encountered in many cosmological models with scalar fields. Examples include such cornerstones of modern cosmology as inflation, as well as the currently popular particle physics models with extra dimensions. It has also been suggested that the motion of a complex flat direction condensate field, spinning in a $U(1)$ -symmetric potential, could give a dynamical explanation for the dark energy [422], although the situation could be complicated by copious Q -ball formation [423].

8.1 Fragmentation of the inflaton condensate

By definition, the inflaton is a homogeneous scalar condensate with a small quantum induced spatial fluctuations. Reheating of the Universe, which is a consequence of the inflaton condensate break-up and decay, may take place via ordinary smooth perturbative decay of the condensate [229, 22] or via a non-perturbative process dubbed as preheating [236, 237], which typically involves an amplification in some of the fluctuation modes as well as the fragmentation of the inflaton condensate. Reheating dynamics depends very much on the form assumed for the inflaton potential, and for some choices, the inflaton condensate may also form Q -balls, see [353, 354].

8.1.1 Reheating as a surface effect

Usually the process of reheating is taken to be entirely a volume effect. This can be problematic, however, especially if the scale of inflation is high, i.e. $H_{inf} \sim 10^{15} - 10^{16}$ GeV, as it is sometimes assumed in order to provide the right magnitude for the density perturbations, and also for reasons that have to do with non-thermal heavy dark matter production or exciting right-handed Majorana neutrinos for leptogenesis, etc. [136]. As it is well known, the entropy thus dumped into the Universe may pose a problem for big bang nucleosynthesis by overproducing gravitinos from a thermal bath; an often quoted bound on the reheat temperature is $T_{rh} \leq 10^9$ GeV [258]. Obtaining such a low reheat temperature is a challenge for high scale inflation models. (One

way to solve the gravitino problem is to dilute them via a brief period of late thermal inflation [172]).

A novel way to avoid the gravitino and other moduli problems is reheating via the surface evaporation of an inflatonic soliton. Compared with the volume driven inflaton decay, the surface evaporation naturally suppresses the decay rate by a factor

$$\frac{area}{volume} \propto L^{-1}, \quad (418)$$

where L is the effective size of an object whose surface is evaporating. The larger the size, the smaller is the evaporation rate, and therefore the smaller is the reheat temperature.

Reheating as a surface phenomenon has been considered [353, 354] in a class of chaotic inflation models where the inflaton field is not real but complex. As the inflaton should have coupling to other fields, the inflaton mass should in general receive radiative corrections [9], resulting in a running inflaton mass and in the simplest case in the inflaton potential that can be written as

$$V = m^2 |\Phi|^2 \left[1 + K \log \left(\frac{|\Phi|^2}{M^2} \right) \right], \quad (419)$$

where the coefficient K could be negative or positive, and m is the bare mass of the inflaton. The logarithmic correction to the mass of the inflaton is something one would expect because of the possible Yukawa and/or gauge couplings to other fields. Though it is not pertinent, we note that the potential Eq. (419) can be generated in a supersymmetric theory if the inflaton has a gauge coupling [49, 348, 330] where $K \sim -(\alpha/8\pi)(m_{1/2}^2/m_{\tilde{\ell}}^2)$, where $m_{1/2}$ is the gaugino mass and $m_{\tilde{\ell}}$ denotes the slepton mass and α is a gauge coupling constant. It is also possible to obtain the potential Eq. (419) in a non-supersymmetric (or in a broken supersymmetry) theory, provided the fermions live in a larger representation than the bosons. In this latter situation the value of K is determined by the Yukawa coupling h with $K = -C(h^2/16\pi^2)$, where C is some number.

As long as $|K| \ll 1$, during inflation the dominant contribution to the potential comes from $m^2 |\Phi|^2$ term, and inflationary slow roll conditions are satisfied as in the case of the standard chaotic model. COBE normalization then implies $m \sim 10^{13}$ GeV.

If $K < 0$, the inflaton condensate feels a negative pressure (see, Sect. 5.9.1, for the discussion on negative pressure of the AD condensate) and it is bound to fragment into lumps of inflatonic matter. Moreover, since the inflation potential Eq. (419) respects a global $U(1)$ symmetry and since for a negative K it is shallower than $m^2|\Phi|^2$, it admits a Q -ball solution (see Sect. 5.9.3). Comparing with Q -balls along the MSSM flat directions, here the major difference is that the inflatonic condensate has no classical motion along the imaginary direction as usually required for a Q -ball solution.

8.1.2 Q -balls from the inflaton condensate

As pointed out in [353], there are quantum fluctuations along both the real and imaginary directions which may act as the initial seed that triggers on the condensate motion in a whole complex plane. The fluctuations in the real direction grow and drag the imaginary direction along via mode-mode interactions, as illustrated by 2 dimensional lattice simulation in [354], see Figs. (13). The first plot shows the linear fluctuations without rescattering effects; scattering effects are accounted for in the second plot. The late time formation of inflatonic solitons is shown in Fig. (14). Q -balls were observed to form with both positive and negative charges, as can be seen in the first plot of Fig. (14), while keeping the net global charge conserved. Inflatonic Q -balls are of same size because the running mass potential resembles the MSSM flat direction potential in the gravity mediated case, where the Q -ball radius is independent of the charge.

Q -balls of size $R \sim |K|^{-1/2}m^{-1}$ form when the fluctuations grow nonlinear (see Eq. (324) [353, 354]). Since the growth rate of fluctuation is $\sim |K|m$, the Hubble parameter at the formation time can be estimated as $H_f \sim \gamma|K|m$, where γ is a numerical coefficient less than one. For $|K| \ll 1$, we can approximate the decrease in the amplitude of the oscillations by $\phi_f \sim \phi_i(H_f/H_i)$ as in the matter dominated era, where $\phi_i \simeq M_P$ denotes the amplitude at the end of inflation in the chaotic model, and $H_i \sim m$ when the oscillations begin. The total charge of a Q -ball is given by $Q \sim (4\pi/3)R^3n_q \sim (1/9)\beta\zeta^2\gamma^2|K|^2R^3mM_P^2$, where $n_q = \beta\omega\phi_0^2$, $\phi_0 \simeq \zeta\phi_f$, and $\beta \ll 1$ and $\zeta \geq 1$ are numerical factors.




Figure 13: The first plot from left shows the instability bands of the homogeneous mode of the inflaton along the real (solid) and imaginary (dotted) directions. The second plot shows the result of lattice simulation in the real and imaginary directions, together with the evolution of the homogeneous mode. The third plot shows the power spectra of fluctuations at late times. All plots assume $K = -0.02$, from [354].

Given an inflaton coupling to fermions of the type $h\phi\bar{\psi}\psi$ it has been shown that [353, 354] reheating is driven by surface evaporation of inflatonic Q -balls for relatively large Yukawa couplings $h \leq 1$. In general K and h are not independent quantities but are related to each other by $|K| \sim C(h^2/16\pi^2)$. If the inflaton sector does not belong to the hidden sector, it is very natural that the inflaton coupling to other matter fields is relatively large, i.e. $h \geq (m/M_P)$. In this regime the evaporation rate is saturated by Eq. (340) and [354]

$$\Gamma_Q = \frac{1}{Q} \frac{dQ}{dt} \simeq \frac{3}{16\pi\beta\zeta^2\gamma^2|K|^{3/2}} \left(\frac{m}{M_P}\right)^2 m. \quad (420)$$

Note that the decay rate is determined by the ratio $m/M_P \simeq 10^{-6}$, which is fixed by the anisotropies seen in the cosmic microwave background radiation. Even though we are




Figure 14: The first plot shows the charge density distribution in a small sub-lattice at late times. The second plot shows inflatonic solitons forming in 3D lattice. Here $K = -0.02$, from [354].

in a relatively large coupling limit, the decay rate mimics that of a Planck suppressed interaction of the inflatonic Q -ball with matter fields. Fermionic preheating [246] is not a problem in this case because the whole inflaton energy is not transferred in this process and the energy density stored in the fermions remains small compared to the inflaton energy density, as argued in [354]. Fermions cannot scatter inflaton quanta off the condensate [246], unlike in the case of bosonic preheating [237]. Inflatonic Q -ball formation is reminiscent of bosonic preheating [424] due to the presence of attractive self coupling of the inflaton, which stems from the logarithmic term in the potential.

8.2 Affleck-Dine baryogenesis without MSSM flat directions

8.2.1 Leptogenesis with sneutrino

It is interesting to note that it is possible to mimic the AD baryogenesis even without MSSM flat directions. This can happen in F-term hybrid inflation with a superpotential [137]

$$W = -\Lambda^2 S + \lambda S \Phi^2 + \kappa \Phi \Psi^2, \quad (421)$$

where $\Lambda \approx 6.5 \cdot 10^{16} \epsilon^{1/4}$ GeV, or equivalently $\Lambda \approx 1.3 \cdot 10^{15} |\eta| \lambda^{-1/2}$ GeV, where $\epsilon, \eta \ll 1$ are the slow roll parameters. The superpotential Eq. (421) extends the usual hybrid

case Eq. (135) in order to provide a natural initial condition for Φ , which should be zero within the accuracy 10^{-5} [224], while the inflaton should have an initial value close to the Planck scale. Ψ is identified with the $U(1)_{B-L}$ carrying right-handed neutrino which alleviates the problem of initial conditions through the dynamics of Eq. (421) [224]. During inflation the system is trapped in a false vacuum with $\phi = 0$. Once ϕ decays, sneutrino $\tilde{\Psi}$ obtains a vev of order of the Hubble parameter at the end of inflation. It then starts oscillating around the origin and due to dynamical breaking of $U(1)_{B-L}$, a net $B - L$ asymmetry is generated [137]. Right after inflation the real and imaginary component of Ψ , ψ_1 and ψ_2 have a relative sign difference which induces a helical motion on Ψ similar to the MSSM flat direction condensate. In this model the neutrino mass is generated by the ordinary see-saw mechanism [32]

$$m_\nu = \frac{g^2 \langle \varphi_2 \rangle^2}{M_\Psi} \simeq \frac{g^2 \lambda}{\kappa \eta} \times 1.2 \cdot 10^{-2} \text{ eV}, \quad (422)$$

(where we have taken $\langle \varphi_2 \rangle \simeq 170 \text{ GeV}$). The Hubble parameter during inflation is

$$H_{inf} \simeq \frac{\Lambda^2}{\sqrt{3} M_P} \simeq \frac{\eta^2}{\lambda} \times 4 \cdot 10^{11} \text{ GeV}. \quad (423)$$

Consideration of the detailed dynamics and decay of the condensate leads to an estimation of the lepton asymmetry, which is given by [137]

$$B - L \sim X \frac{T_r}{M_P} \simeq X \left(\frac{T_r}{10^9 \text{ GeV}} \right) \times 10^{-10}, \quad (424)$$

where the numerical factor is $X = \min\{1, x\} C \kappa^{-1}$, with $C \sim 1$ coefficient if the field $\tilde{\Psi}$ gets a supergravity induced mass term during inflation, otherwise $C = 3/N_e$. The reheat temperature appears to respect the gravitino bound.

8.2.2 AD baryogenesis in theories with low scale quantum gravity

In low scale gravity models the weakness of gravity arises because of extra spatial dimensions, the scale of which could be as large as mm , [425]. In the simplest models, the SM fields live on a three dimensional brane while gravity can also permeate the bulk [425, 426]. The fundamental scale is $M_* \sim \mathcal{O}(\text{TeV})$ in $4 + d$ dimensions, which is related to the the volume suppression V_d and to the effective four dimensional Planck

mass M_p by a simple relationship [425]

$$M_p^2 = M_*^{2+d} V_d. \quad (425)$$

This automatically sets a *common size* for all the extra dimensions at b_0 . For two extra dimensions with $M_* = 1$ TeV, the size is 0.2 mm, just below the current experimental limits from the search for deviations in Newtonian gravity [427]. Recent astrophysical bounds based on neutron stars suggests $M_* \geq 500$ TeV for two extra dimensions [428]. Naturally, low scale gravity models have important implications for collider experiments [429] and for cosmology [430, 431] (for a review on large extra dimensions, see [432]).

Although quite attractive from a particle physics point of view, large extra dimensions bring along a host of cosmological problems. There are dynamical questions regarding the stabilization of the size of the extra dimension(s), or equivalently the vev of the radion field, whose mass can be as small as $\mathcal{O}(\text{eV})$ (for two large extra dimensions) [425]. Cosmologically stabilization should take place very early by some trapping mechanism as discussed in [174]. Another challenge is how to realize inflation in these models. There have been many proposals [431], such as invoking a SM singlet scalar living in the bulk [433]. There is the problem of the Kaluza Klein (KK) states of the graviton and any other fields residing in the bulk, which above a certain temperature known as the *normalcy temperature*, should fill the Universe. The normalcy temperature is constrained by cosmological considerations to lie in the range 1 MeV to 100 MeV [425, 430, 434, 435, 436], and the reheat temperature should be lower than the normalcy temperature.

In addition, one must not only forbid dangerous higher order operators which can mediate proton decay, but also ensure that such operators are not being reintroduced by whatever mechanism is responsible for baryogenesis. This tends to make baryogenesis in low scale gravity models quite difficult. Moreover, it has been argued [434, 435] that leptogenesis is not a viable option; obviously at very low reheat temperatures required by the normalcy temperature, the sphalerons cannot be activated (a possible way out of this problem could be to increase the reheat temperature by increasing the number of large extra dimensions to six [437]).

One solution to this predicament was provided in [435] (see also [438]). There $U(1)_\chi$ carried by a gauge singlet χ was broken dynamically in order to provide a small asymmetry in the current density. Baryon asymmetry is produced by the decays of χ and $\bar{\chi}$ into SM quarks and leptons, analogously to the old AD baryogenesis. The decay channels are constrained because quarks and leptons must carry a non zero global χ charge. This prevents χ - $\bar{\chi}$ asymmetry to be transferred into non-baryon-number violating interactions such as interactions involving the Higgses.

A consistent model for baryogenesis can be constructed [435] along the lines of sneutrino leptogenesis discussed in Sect. 8.2.1. It requires a hybrid inflaton sector, described by the fields ϕ and N , and the flat direction field χ , which are promoted to the bulk. This ensures the right amplitude for the density perturbations and provides enough baryon asymmetry towards the end of reheating [434, 435]. The potential for the zero modes in 4 dimensions can be written as [435]

$$V_{\text{AD}}(\phi, N, \chi_1, \chi_2) = \kappa_1^2 \left(\frac{M_*}{M_{\text{p}}} \right)^2 N^2 (\chi_1^2 + \chi_2^2) + \frac{\kappa_2^2}{4} \left(\frac{M_*}{M_{\text{p}}} \right)^2 (\chi_1^2 + \chi_2^2)^2 + \kappa_3^2 \left(\frac{M_*}{M_{\text{p}}} \right)^2 \phi N (\chi_1^2 - \chi_2^2), \quad (426)$$

where $\kappa_1, \kappa_2, \kappa_3$ are order one constants, and χ_1 and χ_2 are the real and imaginary components of the complex field χ . Note that all the terms are Planck mass suppressed because Eq. (426) is an effective 4d potential derived from a higher dimensional Lagrangian by integrating out the extra spatial dimensions. Since during inflation the auxiliary field $N = 0$, the flat direction condensates χ_1 and χ_2 are massless and the potential is almost flat.

The final χ asymmetry was found to be given by [435]

$$\frac{n_\chi}{s} \approx \frac{2\kappa_3^2}{27\lambda^2} \left(\frac{|\chi(0)|}{M_{\text{p}}} \right)^2 \left(\frac{T_{\text{r}}}{H_0} \right) \leq \frac{2\sqrt{6}\pi}{27} \left(\frac{\kappa_3}{\kappa_2} \right)^2 \left(\frac{1}{\lambda N_{\text{e}}} \right) \left(\frac{T_{\text{r}}}{M_*} \right), \quad (427)$$

which at the same time provides an upper bound on the baryon to entropy ratio. As an example, taking $T_{\text{r}} \sim 100$ MeV, $M_* \sim 100$ TeV, the number of e-foldings $N_{\text{e}} \sim 100$, and couplings of order one, yields an asymmetry of order $\sim 10^{-10}$.

8.3 Solitosynthesis

Solitosynthesis is a mechanism of charge accretion with the help of pre-existing small Q -balls in a charge asymmetric background [439, 440, 441, 443, 444]. The accumulation of charge and forming a large charged Q -ball has been shown to be quite efficient especially in a finite temperature thermal bath.

8.3.1 Accretion of charge by Q -balls

Consider thermodynamics of Q -balls surrounded by massive non-relativistic fermions ψ . The number densities of Q -balls and ψ particles are governed by the Boltzmann distributions $n_Q(T) = g_Q(M_Q T/2\pi)^{3/2} \exp[(\mu_Q - M_Q)/T]$, and $n_\psi = g_\psi(m_\psi T/2\pi)^{3/2} \exp[(\mu_Q - m_\psi)/T]$, where g_Q is the partition function for the Q -ball and $g_\psi = 2$. The respective chemical potentials are denoted by μ . In chemical equilibrium the absorption and evaporation of charge is equally possible, i.e. $(Q) + \psi \leftrightarrow (Q + 1)$, which relates the two chemical potentials through $\mu_Q = Q\mu_\psi$. The number density of Q -balls can be expressed in terms of the number density of the fermions [440, 444]

$$n_Q = \frac{g_Q}{g_\psi} n_\psi \left(\frac{M_Q}{m_\psi} \right)^{3/2} \left(\frac{2\pi}{m_\psi T} \right)^{3(Q-1)/2} e^{B_Q/T}, \quad (428)$$

where $B_Q \equiv Qm_\psi - M_Q$ is the binding energy per charge of a Q -ball. When B_Q grows with Q , the formation of large charged Q -ball is likely. The interactions between Q -ball and ψ quanta leads to a chemical equilibrium when [440, 444]

$$n_\psi v_\psi \sigma_{abs}(Q) = n_{Q+1} r_{evap}(Q + 1). \quad (429)$$

The accretion and evaporation rate from a charge Q -ball is given by Saha equation:

$$\begin{aligned} \frac{dQ}{dt} &= r_{abs}(Q) - r_{evap}(Q), \\ &= n_\psi v_\psi \left[\sigma_{abs}(Q) - \frac{n_Q - 1}{n_Q} \sigma_{abs}(Q - 1) \right], \end{aligned} \quad (430)$$

where $v_\psi = (T/2\pi m_\psi)^{1/2}$ is the mean velocity of ψ particles. The charge of a Q -ball grows when $r_{abs}(Q) > r_{evap}(Q)$. The necessary charge asymmetry in ψ quanta affects the Q -ball abundance. Charge conservation requires

$$N = n_\psi - n_{\psi^*} + \sum Q n_Q + \sum Q^* n_{Q^*} = \eta_\psi n_\gamma, \quad (431)$$

where $\eta_\psi = n_\psi - n_{\psi^*}/n_\gamma$ is the charge asymmetry. If η is close to zero, annihilations of ψ quanta will be effective and there will be virtually no solitosynthesis. The number density of the stable ψ quanta is $n_\psi = \eta_\psi n_\gamma$, and the charge asymmetry is given by [440]

$$\eta_\psi = 2.5 \times 10^{-8} \Omega_\psi h^2 \frac{\text{GeV}}{m_\psi}. \quad (432)$$

Obviously the over-closure limit $\Omega_\psi h^2 \leq 1$ yields an upper bound on η_ψ in terms of m_ψ . At high temperatures large Q -balls are suppressed by a small asymmetry factor $\sim \eta_\psi^{Q-1}$. At lower temperatures the abundance is dominated by $Q^{5/2}$, where $M_Q \propto m_\psi Q$ [440].

The Q -ball starts growing at a temperature T_s when $r_{abs}(Q) > r_{evap}(Q)$. This can be estimated from the Saha equation [443, 444]

$$T_s = \frac{m_\psi + M_{Q-1} - M_Q}{-\frac{3}{2} \ln\left(\frac{T_s}{m_\psi}\right) - \ln(c\xi^3 \eta_\psi)}, \quad (433)$$

where $c \sim \mathcal{O}(1)$ number for large enough Q .

When the absorption of Q -charge from the surroundings freezes out, solitosynthesis stops. There are two distinct era which one may consider; solitosynthesis freezes out during the radiation dominated epoch, or during the matter dominated epoch. The demarcation temperature is $T_{eq} \approx 5.5(\Omega_0 h^2)^{-1} \xi^{-1}$ eV. The freeze-out temperature is defined by $\Gamma[(Q) + \psi \rightarrow (Q+1)] \leq H(T)$. For a geometric cross section $\sigma_{abs} \sim \pi R_Q^2$, the freeze-out temperature is given by [444]

$$\frac{T_F}{m_\psi} \leq \begin{cases} \left[\frac{10^{-9}}{\xi \beta_Q Q^{2/3}} \left(\frac{m_\psi}{\text{GeV}} \right)^2 \left(\frac{0.3}{\Omega_\psi h^2} \right) \left(\frac{g_*^{1/2}}{10} \right) \right]^{2/3} & (T > T_{eq}) \\ \left[\frac{10^{-13}}{\xi \beta_Q Q^{2/3}} \left(\frac{m_\psi}{\text{GeV}} \right)^{3/2} \left(\frac{0.3}{\Omega_\psi h^2} \right)^{1/2} \left(\frac{g_*^{1/2}}{10} \right) \right]^{1/2} & (T < T_{eq}). \end{cases} \quad (434)$$

where $\xi \equiv (g_*(T_D) - g_*(T))^{1/3}$, the temperature T_D signifies the decoupling temperature of a Q -ball with a rest of the plasma, and $\beta_Q = (3m_\phi^3/4\pi\omega\varphi_0^2)^{1/3}$. One should require that $T_F < T_s$, which is satisfied for a very low ψ -mass. In order to have some feeling for the numbers involved, we note that for the smallest charged Q -ball with $Q = 2$, the freeze-out temperature is smaller than T_s only when $m_\psi \leq \mathcal{O}(1)$ GeV. During the matter dominated epoch, due to the gravitational clustering, Q -ball synthesis is favorable but for ambient temperature $T_\gamma \leq \mathcal{O}(1)$ eV, $m_\psi \leq \mathcal{O}(1)$ MeV.

8.3.2 Phase transition via solitogenesis

An interesting observation has been made in [442, 443], where it was pointed out that solitosynthesis may also lead to a first order phase transition. The gradual absorption of charge may lead to a critical charge $Q = Q_c$ for which the false vacuum inside the Q -ball becomes unstable and expands as the whole space is filled with the true vacuum. The value of the critical charge is determined by minimizing the energy of the Q -ball with respect to the radius when $dE/dR = 0$ and $d^2E/dR^2 = 0$ are satisfied simultaneously. In a thin wall limit when $Q_c \gg 1$, the critical charge is given by [443]

$$Q_c = \frac{100\pi\sqrt{10}}{81} \frac{\varphi_0 S_1^3}{U(\varphi_0)^{5/2}}, \quad (435)$$

where $S_1 = \int_0^{\varphi_0} d\varphi \sqrt{2(U(\varphi) - (\omega^2/2)\varphi^2)}$.

One can also enquire how large the critical charge should be in order to facilitate the destabilization of the otherwise cosmologically stable false vacuum. It has been found that the decay of a metastable false vacuum at zero temperature requires a small charge $Q_C \sim 28(\varphi_0/U(\varphi_0)^{1/4})$ for $A \sim (100 \text{ GeV})^4$ [443]. Similar considerations at finite temperature result in a larger critical charge $Q > Q_c \sim 146\varphi_0 T/U(\varphi_0, T)^{1/2}$ for the phase transition to proceed.

Solitogenesis-catalyzed phase transitions in supersymmetric models would require a local violation of lepton or baryon number. As an example (see [443]), one could consider a lepton number violating local minimum along $H_d \tilde{L}_L \tilde{L}_R \neq 0$. This false vacuum can decay into the standard true vacuum but the sphaleron induced transition rate is almost negligible compared to the cosmological time scale because $SU(2) \times U(1)$ is broken. If the typical mass scale of squarks and sleptons is considerably heavier than 1 TeV, L -balls will accumulate when $T_s > T_F \approx m_\phi/40$, and can catalyze a phase transition within one Hubble time at temperatures T_s [442, 443].

Acknowledgements

The authors are thankful to Rouzbeh Allahverdi, Mar Bastero-Gil, Zurab Berezhiani, Ed Copeland, Masaaki Fujii, Katrin Heitmann, Asko Jokinen, Shinta Kasuya, Alex Kusenko, Mikko Laine, Andrew Liddle, John McDonald, Tuomas Multamäki, Altug Özpineci, Abdel Pérez-Lorezana and Iiro Vilja for useful discussions.

A.M. was partially supported by *The Early Universe Network*: HPRN-CT-2000-00152. A.M. is also thankful to the Helsinki Institute of Physics where part of the project has been carried out. K.E. acknowledges the Academy of Finland grant no. 51433.

References

- [1] S. Weinberg, *The quantum theory of fields. v..2: Modern applications*, Cambridge, University Press, (1995).
- [2] S. Sarkar, Rep. Prog. Phys. **59**, 1493 (1996). [arXiv:hep-ph/9602260].
- [3] K. A. Olive, G. Steigman, and T. P. Walker, Phys. Rep. **333**, 389 (2000). [arXiv:astro-ph/9905320].
- [4] <http://map.gsfc.nasa.gov/>
- [5] <http://aether.lbl.gov/www/projects/cosa/>
- [6] P. J. E. Peebles, *Principles of Physical Cosmology*, Princeton, Princeton University Press, (1993).
- [7] Cosmic Background Explorer homepage:
<http://space.gsfc.nasa.gov/astro/cobe/>
- [8] G.F. Smoot, et.al., Ap J. **396**, L1 (1992).
- [9] A. R. Liddle, and D. H. Lyth, *Cosmological Inflation and Large-Scale Structure*, Cambridge, Cambridge University Press (2000).
- [10] C.B. Netterfield et al, BOOMERANG Collaboration, Astrophys. J. **571**, 604 (2002) [arXiv:astro-ph/0104460].
<http://www.physics.ucsb.edu/~boomerang/>
- [11] S. Hanany, et al, MAXIMA Collaboration, Astrophys. J. **545**, L5 (2000) [arXiv:astro-ph/0005123].
<http://cosmology.berkeley.edu/group/cmb/>
- [12] C. Pryke, et al, DASI Collaboration, Astrophys. J. **568**, 46 (2002) [arXiv:astro-ph/0104490].
<http://astro.uchicago.edu/dasi/>

- [13] A. H. Guth, Phys. Rev. D **23**, 347 (1981).
- [14] A. D. Linde, Phys. Lett. B **108**, 389 (1982).
- [15] A. D. Linde, *Particle Physics and Inflationary Cosmology*, New York, Harwood Academy (1990).
- [16] V. F. Mukhanov, H. A. Feldman, and R. H. Brandenberger, Phys. Rept. **215**, 203 (1992).
- [17] A. G. Cohen, A. De Rujula, and S. L. Glashow, Astrophys. J. **495**, 539 (1998) [arXiv:astro-ph/9707087].
- [18] G. Jungman, M. Kamionkowski, and K. Griest, Phys. Rept. **267**, 195 (1996) [arXiv:hep-ph/9506380].
- [19] A. R. Liddle, and D. H. Lyth, Phys. Rept. **231**, 1 (1993) [arXiv:astro-ph/9303019].
- [20] S. Perlmutter, et.al., Astrophys. J. **483**, 55 (1997) [arXiv:astro-ph/9608192].
- [21] A. G. Riess, et.al., Astron. J **116**, 1009 (1998) [arXiv:astro-ph/9805201].
- [22] E. W. Kolb, and M. S. Turner, *The Early Universe*, Adison-Wesley (1990).
- [23] A. D. Sakharov, Pis'ma ZhETF **5**, 32 (1967).
- [24] A. G. Cohen, D. B. Kaplan, and A. E. Nelson, Ann. Rev. Nucl. Part. Sci. **43**, 27 (1993) [arXiv:hep-ph/9302210].
- [25] L. G. Yaffe, *The Electroweak Phase Transition: A Status Report*, arXiv: hep-ph/9512265.
- [26] V. A. Rubakov, and M. E. Shaposhnikov, Usp. Fiz. Nauk **166**, 5 (1996) [arXiv:hep-ph/9603208].
- [27] K. Kajantie, M. Laine, K. Rummukainen, and M. E. Shaposhnikov, Phys. Rev. Lett. **77**, 2887 (1996) [arXiv:hep-ph/9605288].

- [28] A. D. Dolgov, Phys. Rept. **370**, 333 (2002) [arXiv:hep-ph/0202122].
- [29] S. A. Bonometto, F. Gabiani, and A. Masiero, Phys. Rev. D. **49**, 3918 (1994) [arXiv:hep-ph/9305237].
- [30] Y. Fukuda, et. al, Phy. Rev. Lett. **81**, 1158 (1998), Erratum-ibid., **81**, 4279 (1998); ibid. 1562 [arXiv:hep-ex/9805021]. Super-Kamiokande Collaboration, Phys. Rev. Lett. **85**, 3999 (2000). [arXiv:hep-ex/0009001].
- [31] Q. R. Ahmad, et al., SNO Collaboration, Phys. Rev. Lett. **89**, 011302 (2002) [arXiv:nucl-ex/0204009].
- [32] M. Gell-Mann, P. Ramond, and R. Slansky, in *Supergravity*, eds. P. van Nieuwenhuizen and D.Z. Freedman (North Holland 1979); T. Yanagida, Proceedings of *Workshop on Unified Theory and Baryon number in the Universe*, eds. O. Sawada and A. Sugamoto (KEK 1979).
- [33] R.N. Mohapatra, and G. Senjanović, Phys. Rev. Lett. **44**, 912 (1980).
- [34] H. P. Nilles, Phys. Rept. **110**, 1 (1984).
- [35] H. E. Haber, and G. L. Kane, Phys. Rept. **117**, 75 (1985).
- [36] S. P. Martin, in *Kane, G.L. (ed.): Perspectives on supersymmetry*, 1, *A Supersymmetry Primer*, arXiv:hep-ph/9709356.
- [37] D. Bailin, and A. Love, *Supersymmetric Gauge Field Theory and String Theory*, Bristol, Institute of Physics (1994).
- [38] R. McPherson, SUSY Working Group at LEP *SUSY Searches: Lessons and loopholes from LEP*, SUSY02, DESY Hamburg, (2002).
- [39] G. F. Giudice, and R. Rattazzi, Phys. Rep. **322**, 419 (1999) [arXiv:hep-ph/9801271].
- [40] J. M. Cline, M. Joyce, and K. Kainulainen, JHEP **0007**, 018 (2000) [arXiv:hep-ph/0006119].

- [41] M. Quirós, Nucl. Phys. Proc. Suppl. **101**, 401 (2001) [arXiv:hep-ph/0101230].
- [42] M. Carena, M. Quirós, M. Seco, and C. E. M. Wagner, arXiv:hep-ph/0208043.
- [43] T. Gherghetta, C. Kolda, and S. P. Martin, Nucl. Phys. B **468**, 37 (1996) [arXiv:hep-ph/9510370].
- [44] I. Affleck, and M. Dine, Nucl. Phys. B **249**, 361 (1985).
- [45] M. Dine, L. Randall, and S. Thomas, Phys. Rev. Lett. **75**, 398 (1995) [arXiv:hep-ph/9503303].
- [46] M. Dine, L. Randall, and S. Thomas, Nucl. Phys. B **458**, 291 (1996) [arXiv:hep-ph/9507453].
- [47] S. Coleman, Nucl. Phys. B **262**, 263 (1985); Erratum-ibid. B **269**, 744 (1986).
- [48] K. Enqvist, and J. McDonald, Phys. Lett. B **425**, 309 (1998) [arXiv:hep-ph/97111514].
- [49] K. Enqvist, and J. McDonald, Nucl. Phys. B **538**, 321 (1999) [arXiv:hep-ph/9803380].
- [50] A. Kusenko, Phys. Lett. B **405**, 108 (1997) [arXiv:hep-ph/9704273].
- [51] J. Arafune, T. Yoshida, S. Nakamura, and K. Ogure, Phys. Rev. D **62**, 105013 (2000) [arXiv:hep-ph/0005103].
- [52] K. Enqvist, and J. McDonald, Phys. Rev. Lett. **83**, 2510 (1999) [arXiv:hep-ph/9811412].
- [53] K. Enqvist, and J. McDonald, Phys. Rev. D **62**, 043502 (2000) [arXiv:hep-ph/9912478]; Phys. Rev. D **65**, 043002 (2002).
- [54] M. Kawasaki, and F. Takahashi, Phys. Lett. B **516**, 388 (2001) [arXiv:hep-ph/0105134].

- [55] E. A. Bogomolov, et. al., *Proc. of the 16th Intl. Cosmic Ray Conference*, Kyoyo, Vol. **1**, 330 (1979); R. L. Golden, et. al., *Phys. Rev. Lett.* **43**, 1196 (1979).
- [56] C. J. Copi, D. N. Schramm, and M. S. Turner, *Science* **267**, 192 (1995) [arXiv:astro-ph/9407006]; *Phys. Rev. Lett.* **75**, 3981 (1995) [arXiv:astro-ph/9508029].
- [57] K. A. Olive, arXiv: astro-ph/0202486.
- [58] A. Riess, W. Press, and R. Kirshner, *Astrophys. J* **473**, 88 (1996) [arXiv:astro-ph/9604143].
- [59] I. N. Reid, *Astron. J* **114**, 161 (1997). arXiv:astro-ph/9704078.
- [60] D. Tytler, S. Burles, and D. Kirkham, arXiv: astro-ph/9612121.
- [61] J. Levin, *Phys. Rep.* **365**, 251 (2002). [arXiv:gr-qc/0108043].
- [62] E. D. Masso, and F. Rota, *Phys. Lett. B* **545**, 221 (2002) [arXiv:astro-ph/0201248].
- [63] E. R. Harrison, *Ann. Rev. Astron. Astrophys.* **11**, 155 (1973), *Astrophys. J* **403**, 28 (1993).
- [64] M. Srednicki, W. Watkins, and K. A. Olive, *Nucl. Phys. B.* **310**, 693 (1988).
- [65] See, *Review of Particle Physics*, K. Hagiwara et al., *Phys. Rev. D* **66**, 010001 (2002).
- [66] C. M. Will, *Theory and Experiment in Gravitational Physics*, Cambridge, Cambridge University Press, (1993).
- [67] R. Stompor et. al., *Ap. J.* **545**, L1 (2001) [arXiv:astro-ph/0005124].
- [68] R. Omne, *Phy. Rev. Lett.* **23**, 38 (1969); *Phys. Rev. D* **1**, 723 (1970).
- [69] K. Sato, *Phys. Lett. B* **99**, 66 (1981).
- [70] G. t' Hooft, *Phys. Rev. Lett.* **37**, 8 (1976).

- [71] G. t' Hooft, Phys. Rev. D **14**, 3432 (1976); Erratum-ibid. D **18**, 2199 (1978).
- [72] S. L. Adler, Phys. Rev. **177**, 2426 (1969); J. S. Bell, and R. Jackiw, Nuovo Cimento A **60**, 47, (1969).
- [73] N. S. Manton, Phys. Rev. D **28**, 2019 (1983).
- [74] K. Enqvist, and D. V. Nanopoulos, Prog. Part. Nucl. Phys. **16**, 1 (1986); K. Enqvist, A. Masiero, and D. V. Nanopoulos, Phys. Lett. B **156**, 209 (1985); R. Barbieri, and A. Masiero, Nucl. Phys. B **267**, 679 (1986); S. Dimopoulos, and L. Hall, Phys. Lett. B **207**, 210 (1987); R. Godbole, P. Roy, and X. Tata, Nucl. Phys. B **401**, 67 (1992) [arXiv:hep-ph/9209251].
- [75] R. D. Peccei, and H. R. Quinn, Phys. Rev. Lett. **38**, 1440 (1977); Phys. Rev. D **16**, 1791 (1977).
- [76] A. Vilenkin, and E. P. S. Shellard, *Cosmic Strings and Other Topological Defects*, Cambridge, Cambridge University Press (1994).
- [77] S. Weinberg, Phys. Rev. Lett. **42**, 860 (1979).
- [78] A. D. Dolgov, and Ya. B. Zeldovich, Rev. Mod. Phys. **53**, 1 (1981).
- [79] E. W. Kolb, and M. S. Turner, Ann. Rev. Nucl. Part. Sci. **33**, 645 (1983).
- [80] J. Kapusta, *Finite Temperature Field Theory*, Cambridge, Cambridge University Press, (1988).
- [81] F. Karsch, E. Laermann, and A. Peikrt, Phys. Lett. B **478**, 447 (2000) [arXiv:hep-lat/0002003]. S. Ejiri, et. al, *The QCD Phase Transition at High Temperature and Low Density*, arXiv: hep-lat/0209012.
- [82] A. Zhitnitsky, R. H. Brandenberger, and I. E. Halperin, AIP. Conf. Proc. **478**, 251 (1999).
- [83] F. R. Klinkhamer, and N. S. Manton, Phys. Rev. D **30**, 2212 (1984).
- [84] J. Kunz, B. Kleihaus, and Y. Brihaye, Phys. Rev. D **46**, 3587 (1992).

- [85] Y. Brihaye, and J. Kunz, Phys. Rev. D **48**, 3884 (1993) [arXiv:hep-ph/9308290].
- [86] J. M. Moreno, D. H. Oaknin, and M. Quirós, Nucl. Phys. B **483**, 267 (1997) [arXiv:hep-ph/9612212].
- [87] A. I. Bochkarev, and M. E. Shaposhnikov, Mod. Phys. Lett. **A2**, 417 (1987).
Erratum-ibid. A **4**, 1495 (1989).
- [88] A. D. Dolgov, Phys. Rep. **222**, 309 (1992).
- [89] M. Trodden, Rev. Mod. Phys. **71**, 1463 (1999) [arXiv:hep-ph/9803479].
- [90] A. Riotto, and M. Trodden, Ann. Rev. Nucl. Part. Sci. **49**, 35 (1999) [arXiv:hep-ph/9901362].
- [91] A. Riotto, *Talk given at ICTP Summer School in High-Energy Physics and Cosmology*, arXiv: hep-ph/9807454.
- [92] A. Polyakov, Nucl. Phys. B **120**, 429 (1977).
- [93] V. A. Kuzmin, V. A. Rubakov, and M. E. Shaposhnikov, Phys. Lett. B **155**, 36 (1985).
- [94] L. Carson, Xu-Li, L. McLerran, and R.-T. Wang, Phys. Rev. D **42**, 2127 (1992).
- [95] M. Dine, R. G. Leigh, P. Huet, A. Linde, and D. Linde, Phys. Lett. B **283**, 319 (1992) [arXiv:hep-ph/9203201]; Phys. Rev. D **46**, 550 (1992) [arXiv:hep-ph/9203203].
- [96] P. Arnold, D. Son, and L. G. Yaffe, Phys. Rev. D **55**, 6264 (1997) [arXiv:hep-ph/9609481]; G. D. Moore, Phys. Rev. D **59**, 014503 (1999) [arXiv:hep-ph/9810313]; D. Bodeker, Phys. Lett. B **426**, 351 (1998) [arXiv:hep-ph/9801430].
- [97] G. D. Moore, C. Hu, and B. Muller, Phys. Rev. D **58**, 045001 (1998) [arXiv:hep-ph/9710436].
- [98] K. A. Olive, Phys. Rept. **190**, 307 (1990).

- [99] K. Enqvist, and K. J. Eskola, *Mod. Phys. Lett. A* **5**, 1919 (1990)
- [100] P. Elmfors, K. Enqvist, and I. Vilja, *Phys. Lett. B* **326**, 37 (1994) [arXiv:hep-ph/9312265].
- [101] K. Enqvist, and J. Sirkka, *Phys. Lett. B* **314**, 298 (1993) [arXiv:hep-ph/9304273].
- [102] J. Ellis, J. E. Kim, and D. V. Nanopoulos, *Phys. Lett. B* **145**, 181 (1984)
- [103] V. A. Kuzmin, V. A. Rubakov, and M. E. Shaposhnikov, *Phys. Lett. B* **191**, 171 (1987).
- [104] J. Ambjorn, M. Laursen, and M. Shaposhnikov, *Phys. Lett. B* **197**, 49 (1989); J. Ambjorn, T. Askaard, H. Porter, and M. E. Shaposhnikov, *Phys. Lett. B* **244**, 479 (1990); *Nucl. Phys. B* **353**, 346 (1991); J. Ambjorn, and K. Farakos, *Phys. Lett. B* **294**, 248 (1992) [arXiv:hep-lat/9207020]; J. Ambjorn, and A. Krasnitz, *Phys. Lett. B* **362**, 97 (1995) [arXiv:hep-ph/9508202]; L. McLerran, M. E. Shaposhnikov, N. Turok, and M. Voloshin, *Phys. Lett. B* **256**, 451 (1991); E. Farhi, J. Goldstone, S. Gutmann, K. Rajagopal, and R. Singleton Jr, *Phys. Rev. D* **51**, 4561 (1995) [arXiv:hep-ph/9410365]; A. Lue, K. Rajagopal, and M. Trodden, *Phys. Rev. D* **55**, 1250 (1997) [arXiv:hep-ph/9612282]. G. D. Moore, and N. Turok, *Phys. Rev. D* **55**, 6538 (1997) [arXiv:hep-ph/9608350]; **56**, 6533 (1997) [arXiv:hep-ph/9703266].
- [105] N. Turok, and J. Zadrozny, *Phys. Rev. Lett.* **65**, 2331 (1990); N. Turok, and J. Zaroni, *Nucl. Phys. B* **358**, 471 (1991).
- [106] M. McLerran, *Phys. Rev. Lett.* **62**, 1075 (1989).
- [107] J. F. Gunion, H. E. Haber, G. L. Kane, and S. Dawson, *The Higgs Hunters Guide*, Addison-Wesley, (1989).
- [108] A. G. Cohen, and A. E. Nelson, *Phys. Lett. B* **297**, 111 (1992) [arXiv:hep-ph/9209245].

- [109] J. Ellis, S. Ferrara, and D. V. Nanopoulos, Phys. Lett. B **114**, 231 (1982).
- [110] P. Huet, and A. E. Nelson, Phys. Rev. D **53**, 4578 (1996) [arXiv:hep-ph/9506477].
- [111] M. Carena, J. M. Moreno, M. Quirós, M. Seco, and C. E. M. Wagner, Nucl. Phys. B **599**, 158 (2001) [arXiv:hep-ph/0011055].
- [112] S. J. Huber, P. John, and M. G. Schmidt, Eur. Phys. J. C **20**, 695 (2001) [arXiv:hep-ph/0101249].
- [113] J. M. Cline, M. Joyce, and K. Kainulainen, hep-ph/0110031.
- [114] M. E. Shaposhnikov, Nucl. Phys. B **287**, 757 (1987); Nucl. Phys. B **299**, 797 (1988).
- [115] A. I. Bochkarev, V. Kuzmin, and M. E. Shaposhnikov, Phys. Rev. D **43**, 369 (1991).
- [116] J. Garcia-Bellido, D. Y. Grigoriev, A. Kusenko and M. E. Shaposhnikov, Phys. Rev. D **60**, 123504 (1999) [arXiv:hep-ph/9902449].
- [117] L. M. Krauss and M. Trodden, Phys. Rev. Lett. **83**, 1502 (1999) [arXiv:hep-ph/9902420].
- [118] M. Joyce and T. Prokopec, JHEP **0010**, 030 (2000) [arXiv:hep-ph/0003190].
- [119] M. Gurtler, E. M. Ilgenfritz, and A. Schiller, Phys. Rev. D **56**, 3888 (1997) [arXiv:hep-lat/9704013].
- [120] K. Rummukainen, M. Tsy-pin, K. Kajantie, M. Laine, and M. E. Shaposhnikov, Nucl. Phys. B **532**, 283 (1998) [arXiv:hep-lat/9805013].
- [121] F. Csikor, Z. Fodor, and J. Heitger, Phys. Rev. Lett. **82**, 21 (1999) [arXiv:hep-ph/9809291].

- [122] G. F. Giudice, Phys. Rev. D **45**, 3177 (1992); S. Myint, Phys. Lett. B **287**, 325 (1992) [arXiv:hep-ph/9206266]; J. R. Espinosa, M. Quirós, and F. Zwirner, Phys. Lett. B **307**, 106 (1993) [arXiv:hep-ph/9303317]; A. Brignole, J. R. Espinosa, M. Quirós, and F. Zwirner, Phys. Lett. B **324**, 181 (1994) [arXiv:hep-ph/9312296]; D. Delepine, J. M. Gerard, R. G. Felipe, and J. Weyers, Phys. Lett. B **386**, 183 (1996) [arXiv:hep-ph/9604440]; J. Espinosa, Nucl. Phys. B **475**, 273 (1996) [arXiv:hep-ph/9604320]; D. Bodeker, P. John, M. Laine, and M. G. Schmidt, Nucl. Phys. B **497**, 387 (1997) [arXiv:hep-ph/9612364]; B. de Carlos, and J. R. Espinosa, Nucl. Phys. B **503**, 24 (1997). [arXiv:hep-ph/9703212].
- [123] M. Carena, M. Quirós, and C. E. M. Wagner, Phys. Lett. B **380**, 81 (1996) [arXiv:hep-ph/9603420].
- [124] M. Carena, M. Quirós, and C. E. M. Wagner, Nucl. Phys. B **524**, 3 (1998) [arXiv:hep-ph/9710401].
- [125] J. M. Cline, M. Joyce, and K. Kainulainen, Phys. Lett. B **417**, 79 (1998) Erratum-ibid. B **448**, 321 (1999) [arXiv:hep-ph/9708393].
- [126] M. Laine, Nucl. Phys. B **481**, 43 (1996) [arXiv:hep-lat/9604011].
- [127] J. M. Cline, and K. Kainulainen, Nucl. Phys. B **482**, 73 (1996) [arXiv:hep-ph/9605235].
- [128] M. Laine, and K. Rummukainen, Nucl. Phys. B **535**, 423 (1998) [arXiv:hep-ph/9804255].
- [129] P. Lutz, for the LEP Working Group on Higgs boson searches, see:
<http://lephiggs.web.cern.ch/LEPHIGGS/talks/index.html>
- [130] J. M. Cline, and K. Kainulainen, Phys. Rev. Lett. **85**, 5519 (2000) [arXiv:hep-ph/0002272]; K. Kainulainen, T. Prokopec, M. G. Schmidt, and S. Weinstock, JHEP **0106**, 031 (2001) [arXiv:hep-ph/0105295]; K. Kainulainen, T. Prokopec, M. G. Schmidt, and S. Weinstock, Phys. Rev. D **66**, 043502 (2002) [arXiv:hep-ph/0202177].

- [131] J. A. Harvey, and E. W. Kolb, Phys. Rev. D **24**, 2090 (1981).
- [132] M. Fukugita, and T. Yanagida, Phys. Lett. B **174**, 45 (1986); M. Fukugita and T. Yanagida, Phys. Rev. D **42**, 1285 (1990).
- [133] W. Buchmüller, and M. Plümacher, Int. J. Mod. Phys. A **15**, 5047 (2000) [arXiv:hep-ph/0007176].
- [134] P. Langacker, Phys. Rept. **72**, 185 (1981).
- [135] R. N. Mohapatra, *Unification and Supersymmetry*, Springer-Verlag (1991).
- [136] F. Giudice, A. Riotto, I. Tkachev, and M. Peloso, JHEP, 9908 014, (1999) [arXiv:hep-ph/9905242].
- [137] Z. Berezhiani, A. Mazumdar, and A. Pérez-Lorenzana, Phys. Lett. B **518**, 282 (2001) [arXiv:hep-ph/0107239].
- [138] M. A. Luty, Phys. Rev. D **45**, 455 (1992).
- [139] M. Plümacher, Z. Phys. C **74**, 549 (1997). [arXiv:hep-ph/9604229].
- [140] S. Yu. Khlebnikov, and M. E. Shaposhnikov, Nucl. Phys. B **308**, 885 (1998).
- [141] K.-I. Aoki, Phys. Lett. B **174**, 371 (1986); J. A. harvey, and M. S. Turner, Phys. Rev. D **42**, 3344 (1990); A. E. Nelson, and S. M. Barr, Phys. Lett. B **246**, 141 (1990)
- [142] P. Langacker, R. D. Peccei, and T. Yanagida, Mod. Phys. Lett. A **1**, 541 (1986); L. Covi, E. Roulet, and F. Vissani, Phys. Lett. B **384**, 169 (1996) [arXiv:hep-ph/9605319]; **424**, 101 (1998) [arXiv:hep-ph/9712468].
- [143] W. Buchmüller, and M. Plümacher, Phys. Lett. B **389**, 73 (1996); [arXiv:hep-ph/9608308]. W. Buchmüller, and M. Plümacher, Phys. Lett. B **431**, 354 (1998) [arXiv:hep-ph/9710460].
- [144] A. Pilaftsis, Int. J. Mod. Phys. A **14**, 1811 (1999) [arXiv:hep-ph/9812256].

- [145] W. Buchmüller, P. Di Bari, and M. Plümacher, Nucl. Phys. B **643**, 367 (2002) [arXiv:hep-ph/0205349].
- [146] T. Hambye, Nucl. Phys. B **633**, 171 (2002) [arXiv:hep-ph/0111089].
- [147] G. Lazarides, Q. Shafi, Phys. Lett. B **258**, 305 (1991); Phys. Lett. B **258**, 305 (1991). K. Kumekawa, T. Moroi, and T. Yanagida, Prog. Theor. Mod. Phys. **92**, 437 (1994) [arXiv:hep-ph/9405337]; ; G. Lazarides, *Springer Tracts Mod. Phys.* **163**, 227 (2000) [arXiv:hep-ph/9904428]; T. Asaka, K. Hamaguchi, M. Kawasaki, and T. Yanagida, Phys. Lett. B **464**, 12 (1999) [arXiv:hep-ph/9906366]; Phys. Rev. D **61**, 083512 (2000) [arXiv:hep-ph/9907559]; M. Fujii, K. Hamaguchi, and T. Yanagida, Phys. Rev. D **65**, 115012 (2002) [arXiv:hep-ph/0202210].
- [148] R. Allahverdi, and A. Mazumdar, *Non-thermal leptogenesis with almost super-heavy neutrinos*, arXiv:he-ph/0208268
- [149] L. Bento, and Z. Berezhiani, Phys. Rev. Lett. **87**, 231304 (2001) [arXiv:hep-ph/0107281].
- [150] B. A. Campbell, S. A. Davidson, and K. A. Olive, Nucl. Phys. B **399**, 111 (1993) [arXiv:hep-ph/9302223]. H. Murayama, and T. Yanagida, Phys. Lett. B **322**, 349 (1994) [arXiv:hep-ph/9310297]. K. Hamaguchi, H. Murayama, and T. Yanagida, Phys. Rev. D **65**, 043512 (2002) [arXiv:hep-ph/0109030].
- [151] A. Kusenko, and M. E. Shaposhnikov, Phys. Lett. B **418**, 46 (1998) [arXiv:hep-ph/9709492].
- [152] K. Wang Ng, Nucl. Phys. B **321**, 528 (1989); D. Morgan, Nucl. Phys. B **364**, 401 (1991).
- [153] J. Ellis, D. V. Nanopoulos, and K. A. Olive, Phys. Lett. B **184**, 37 (1987); J. Ellis, K. Enqvist, D. V. Nanopoulos, and K. A. Olive, Phys. Lett. B **191**, 343 (1987).

- [154] A. A. Starobinsky, Phys. Lett. B **91**, 99 (1980).
- [155] D. H. Lyth, and A. Riotto, Phys. Rept. **314**, 1 (1998) [arXiv:hep-ph/9807278].
- [156] T. Vachaspati, and M. Trodden, Phys. Rev. D **61**, 023502 (2000) [arXiv:gr-qc/9811037].
- [157] A. Albrecht, R. H. Brandenberger, and R. A. Matzner, Phys. Rev. D **32**, 1280 (1985); J. H. Kung, and R. H. Brandenberger, Phys. Rev. D **42**, 1008 (1990); I. Yi, and E. T. Vishniac, Phys. Rev. D **47**, 5280 (1993).
- [158] A. D. Linde, Phys. Lett. B **129**, 177 (1983).
- [159] A. D. Linde, Phys. Lett. B **175**, 395 (1986).
- [160] A. D. Linde, D. Linde, and A. Mezhlumian, Phys. Rev. D **49**, 1783 (1994) [arXiv:gr-qc/9306035].
- [161] A.R. Liddle, A. Mazumdar, and F.E. Schunck, Phys. Rev D **58**, 061301 (1998) [arXiv:astro-ph/9804177].
- [162] K. Malik, and D. Wands, Phys. Rev. D **59**, 123501 (1999) [arXiv:astro-ph/9812204]; E.J. Copeland, A. Mazumdar, and N. Nunes, Phys. Rev. D **60**, 83508 (1999) [arXiv:astro-ph/9904309]; P. Kanti, and K.A. Olive, Phys. Rev. D **60**, 043502 (1999) [arXiv:hep-ph/9903524]; A. M. Green, and J. E. Lidsey, Phys. Rev. D **61**, 067301 (2000) [arXiv:astro-ph/9907223].
- [163] T. S. Bunch, and P. C. W. Davies, Proc. R. Soc. London A **360**, 117 (1978).
- [164] A. Vilenkin, Phys. Lett. B **115**, 91 (1982).
- [165] A. Vilenkin, and L. H. Ford, Phys. Rev. D **26**, 1231 (1982).
- [166] A. D. Linde, Phys. Lett. B **116**, 335 (1982).
- [167] K. Enqvist, K. W. Ng, and K. Olive, Nucl. Phys. B **303**, 713 (1988).
- [168] A. Vilenkin, Nucl. Phys. B **226**, 527 (1983).

- [169] A. D. Linde, Phys. Rev. D **49**, 748 (1994) [arXiv:astro-ph/9307002].
- [170] T. Damour, and V. F. Mukhanov, Phys. Rev. Lett. **80**, 3440 (1998) [arXiv:gr-qc/9712061]; A. R. Liddle, and A. Mazumdar, Phys. Rev. D **58**, 083508 (1998) [arXiv:astro-ph/9806127].
- [171] D. S. Salopek, and J. R. Bond, Phys. Rev. D **42**, 3936 (1990); A. R. Liddle, Phys. Rev. D **49**, 739 (1994) [Erratum-ibid. D **51**, 4603 (1995)] [arXiv:gr-qc/9307036].
- [172] D. H. Lyth, and E. D. Stewart, Phys. Rev. Lett. **75**, 201 (1995) [arXiv:hep-ph/9502417]; Phys. Rev. D **53**, 1784 (1996) [arXiv:hep-ph/9510204].
- [173] A. Mazumdar, Phys. Lett. B **469**, 55 (1999) [arXiv:hep-ph/9902381].
- [174] A. Mazumdar, and A. Pérez-Lorezana, Phys. Lett. B **508**, 340 (2001). [arXiv:hep-ph/0102174].
- [175] E. Lifshitz, Zh. Eksp. Teor. Fiz **16**, 587 (1946); E. Lifshitz, and I. Khalatnikov, Adv. Phys. **12**, 185 (1963); S. Hawking, Astrophys. J **145**, 544 (1966); G. Field, and L. Shapley, Astrophys. Space Sci. **1**, 309 (1968); J. Bardeen, Bull. Am. Astron. Soc. **73**, (1968); U. Gerlach, and U. Sengupta, Phys. Rev. D **18**, 1789 (1978); J. Bardeen, Phys. Rev. D **22**, 1882 (1980).
- [176] V. Mukhanov, and G. Chibisov, Pis'ma Zh. Eksp. Teor. Fiz. **33**, 549 (1981); JETP Lett. **33**, 532 (1981).
- [177] A. Guth, and S.-Y. Pi, Phys. Rev. Lett. **49**, 1110 (1982).
- [178] S. W. Hawking, Phys. Lett. B **115**, 295 (1982).
- [179] A. Starobinsky, Phys. Lett. B **117**, 175 (1982).
- [180] J. Bardeen, P. J. Steinhardt, and M. S. Turner, Phys. Rev. D **28**, 679 (1983).
- [181] A. Starobinsky, JETP Lett. **42**, 152 (1985); Stron. Lett. **11**, 271 (1985).
- [182] R. H. Brandenberger, Rev. Mod. Phys. **57**, 1 (1985).

- [183] V. Lukash, Zh. Eksp. Teor. Fiz. **79**, 1601 (1980); Sov. Phys. JETP. **52**, 807 (1980); D. H. Lyth, Phys. Rev. D **31**, 1792 (1985).
- [184] R. K. Sachs, and A. M. Wolfe, Astrophys. J **147**, 73 (1967).
- [185] R. Harrison, Phys. Rev. D **1**, 2726 (1970); Ya. B. Zeldovich, Astron. Astrophys. **5**, 84 (1970).
- [186] A. R. Liddle, and D. H. Lyth, Phys. Lett. B **291**, 391 (1992) [arXiv:astro-ph/9208007].
- [187] A. Kosowsky, and M. S. Turner, Phys. Rev. D **52**, 1739 (1995) [arXiv:astro-ph/9504071].
- [188] V. F. Mukhanov, JETP Lett. **41**, 493 (1985). [Pisma Zh. Eksp. Teor. Fiz. **41**, 402 (1985)].
- [189] M. Sasaki, Prog. Theor. Phys. **76**, 1036 (1986).
- [190] V. F. Mukhanov, Phys. Lett. B **218**, 17 (1989).
- [191] J. E. Lidsey, A. R. Liddle, E. W. Kolb, E. J. Copeland, T. Barrerio, and M. Abney, Rev. Mod. Phys. **69**, 373 (1997) [arXiv:astro-ph/9508078].
- [192] L. P. Grishchuk, Zh. Eksp. Teor. Fiz 67, 825 (1974); Sov. Phys. JETP **40**, 409 (1974).
- [193] L. Grishchuk, and Yu. Sidorov, Class. Quant. Grav. **6**, L161 (1989); Yu. Sidorov, Europhys. Lett. **10**, 415 (1989); L. Grishchuk, and Yu. Sidorov, Phys. Rev. D **42**, 3413 (1990); B. Allen, Phys. Rev. D **37**, 2078 (1988); V. Sahni, Phys. Rev. D **42**, 453 (1990);
- [194] M. Maggiore, Phys. Rept. **331**, 283 (2000) [arXiv:gr-qc/9909001].
- [195] M. Sasaki, and E. D. Stewart, Prog. Theor. Phys. **95**, 71 (1996) [arXiv:astro-ph/9507001].
- [196] D. S. Salopek, Phys. Rev. D **52**, 5563 (1995) [arXiv:astro-ph/9506146].

- [197] H. Kodama, and M. Sasaki, Prog. Theor. Phys. Supp. **78**, 1 (1984).
- [198] A. A. Starobinsky, and V. Sahni, *Modern Theoretical and Experimental Problems of General Relativity*, Moscow University Press, in Russian; J. R. Bond, and G. Efstathiou, Astrophys. J. **285**, L45 (1984); H. Kodama, and M. Sasaki, Int. J. Mod. Phys. A **2**, 491 (1987).
- [199] A. R. Liddle, and A. Mazumdar, Phys. Rev. D **61**, 123507 (2000) [arXiv:astro-ph/9912349].
- [200] A. D. Linde, and V. F. Mukhanov, Phys. Rev. D **56**, 535 (1997) [arXiv:astro-ph/9610219].
- [201] R. Stompor, K. M. Gorski, and A. J. Banday, Astrophys. J **463**, 9 (1996) [astro-ph/9511087].
- [202] K. Enqvist, H. Kurki-Suonio, and J. Valiviita, Phys. Rev. D **62**, 103003 (2000) [arXiv:astro-ph/0006429]; Phys. Rev. D **65**, 043002 (2002) [arXiv:astro-ph/0108422].
- [203] M. Kawasaki, N. Sugiyama, and T. Yanagida, Phys. Rev. D **54**, 2442 (1996) [arXiv:hep-ph/9512368].
- [204] K. Enqvist, and H. Kurki-Suonio, Phys. Rev. D **61**, 043002 (2000) [arXiv:astro-ph/9907221].
- [205] E. Pierpaoli, J. Garcia-Bellido, and S. Borgani, JHEP 9910:015 (1999) [arXiv:hep-ph/9909420]. L. Amendola, C. Gordon, and D. Wands, Phys. Rev. Lett. **88** 211302 (2002) [arXiv:astro-ph/0107089].
- [206] D. Langlois, Phys. Rev. D **59**, 123512 (1999); D. Langlois, and A. Riazuelo, Phys. Rev. D **62**, 043504 (2000). [arXiv:astro-ph/9912497].
- [207] M. A. Bucher, K. Moodley, and N. G. Turok, Phys. Rev. D **62**, 083508 (2000) [arXiv:astro-ph/9904231]; M. A. Bucher, K. Moodley, and N. G. Turok, Phys. Rev. Lett. **87**, 191301 (2001) [arXiv:astro-ph/0012141]; ; M. A. Bucher, K.

- Moodley, and N. G. Turok, Phys. Rev. D **66**, 023528 (2002) [arXiv:astro-ph/0007360].
- [208] T. Kanazawa, M. Kawasaki, N. Sugiyama, and T. Yanagida, Prog. Theor. Phys. **102**, 71 (1999) [arXiv:astro-ph/9805102].
- [209] J. S. Dowker, and R. Critchley, Phys. Rev. D **13**, 3224 (1976).
- [210] A. Vilenkin, Phys. Rev. D **32**, 2511 (1985).
- [211] A. A. Starobinsky, Sov. Astron. Lett. **9**, 302 (1983).
- [212] H. Buchdahl, Nuovo. Cimento **23**, 141 (1962); J. Phys. A **11**, 871 (1978); G. V. Bicknell, J. Phys. A **7**, 1061 (1974).
- [213] B. Whitt, Phys. Lett. B **145**, 176 (1984); J. D. Barrow, and S. Cotsakis, Phys. Lett. B **214**, 515 (1988); K. Maeda, Phys. Rev. D **39**, 3159 (1989); D. Wands, Class. Quant. Grav. **11**, 269 (1994) [arXiv:gr-qc/9307034].
- [214] D. La., and P. J. E. Steinhardt, Phys. Rev. Lett. **62**, 376 (1989) [Erratum-ibid. **62**, 1066 (1989)]; J. D. Barrow, and K. Maeda, Nucl. Phys. B **341**, 294 (1990).
- [215] A. D. Linde, Phys. Lett. B **259**, 38 (1991).
- [216] J. Ellis, D. V. Nanopoulos, K. A. Olive, and K. Tamvakis, Phys. Lett. B **118**, 335 (1982); Nucl. Phys. B **221**, 224 (1983).
- [217] E. J. Copeland, A. R. Liddle, D. H. Lyth, E. D. Stewart, and D. Wands, Phys. Rev. D **49**, 6410 (1994) [arXiv:astro-ph/9401011].
- [218] G. Dvali, Q. Shafi, and R. Schaefer, Phys. Rev. Lett. **73**, 1886 (1984) [arXiv:hep-ph/9406319].
- [219] S. Coleman, and E. J. Weinberg, Phys. Rev. D **7**, 1888 (1973).
- [220] E. D. Stewart, Phys. Lett. B **391**, 34 (1997); [arXiv:hep-ph/9606241]. E. D. Stewart, Phys. Rev. D **56**, 2019 (1997) [arXiv:hep-ph/9703232].

- [221] L. Covi, and D. H. Lyth, Phys. Rev. D **59**, 063515 (1999) [arXiv:hep-ph/9809562].
- [222] P. Binetruy, and G. Dvali, Phys. Lett. B **388**, 241 (1996) [arXiv:hep-ph/9606342].
- [223] M. Dine, N. Seiberg, and E. Witten, Nucl. Phys. B **289**, 585 (1987); M. Dine, I. Ichinose, and N. Seiberg, Nucl. Phys. B **293**, 253 (1987); J. J. Atick, L. J. Dixon, and A. Sen, Nucl. Phys. B **292**, 109 (1987).
- [224] Z. Berezhiani, D. Comelli, and N. Tetradis, Phys. Lett. B. **431**, 286 (1998) [arXiv:hep-ph/9803498].
- [225] M. Dine, W. Fischler, and D. Nemeschansky, Phys. Lett. B **136**, 169 (1984); G. D. Coughlan, R. Holman, P. Ramond, and G. G. Ross, Phys. Lett. B **140**, 44 (1984); A. S. Goncharov, A. D. Linde, and M. I. Vysotsky, Phys. Lett. B **147**, 279 (1984); O. Bertolami, and G. G. Ross, Phys. Lett. B **183**, 163 (1987).
- [226] E. D. Stewart, Phys. Rev. D **51**, 6847 (1995) [arXiv:hep-ph/9405389]; Phys. Lett. B **345**, 414 (1995) [arXiv:astro-ph/9407040].
- [227] R. Jeannerot, Phys. Rev. D **56**, 6205 (1997) [arXiv:hep-ph/9706391]; E. Halyo, Phys. Lett. B **387**, 48 (1996) [arXiv:hep-ph/9606423].
- [228] C. Kolda, and J. March-Russell, Phys. Rev. D **60**, 023504 (1998) [arXiv:hep-ph/9802358].
- [229] A. Albrecht, P. J. Steinhardt, M. S. Turner, and F. Wilczek, Phys. Rev. Lett. **48**, 1437 (1982); A. D. Dolgov, and A. D. Linde, Phys. Lett. B **116**, 329 (1982); L. F. Abbott, E. Farhi, and M. Wise Phys. Lett. B **117**, 29 (1982).
- [230] M. S. Turner, Phys. Rev. D **28**, 1243 (1983).
- [231] J. Ellis, K. Enqvist, D. V. anopoulos, and K. A. Olive, Phys. Lett. B **191**, 343 (1987); S. Dodelson, Phys. Rev. D **37**, 2059 (1988); W. Zimdahl, D. Pavon, and R. Maartens, Phys. Rev. D **55**, 4681 (1997); [arXiv:astro-ph/9611147].

- [232] D. J. H. Chung, E. W. Kolb, and A. Riotto, Phys. Rev. D **60**, 063504 (1999) [arXiv:hep-ph/9809453]; J. McDonald, Phys. Rev. D **61**, 083513 (2000) [arXiv:hep-ph/9909467]. R. Allahverdi, Phys. Rev. D **62**, 063509 (2000) [arXiv:hep-ph/0004035].
- [233] S. Davidson, and S. Sarkar, JHEP **0011**, 012 (2000) [arXiv:hep-ph/0009078].
- [234] R. Allahverdi, and M. Drees, Phys. Rev. D **66**, 063513 (2002) [arXiv:hep-ph/0205246]. Phys. Rev. Lett. **89**, 091302 (2002) [arXiv:hep-ph/0203118].
- [235] P. Jaikumar, and A. Mazumdar, *Post-inflationary thermalization and hadronization: QCD based approach* [arXiv:hep-ph/0212265].
- [236] J. Traschen, R. Brandenberger, Phys. Rev. D **42**, 2491 (1990); Y. Shtanov, Ukr. Fiz. Zh. **38**, 1425 (1993). (in Russian); Y. Shtanov, J. Traschen, and R. Brandenberger, Phys. Rev. D **51**, 5438 (1995) [arXiv:hep-ph/9407247].
- [237] L. Kofman, A. Linde, and A. Starobinsky, Phys. Rev. Lett. **73**, 3195 (1994). [arXiv:hep-th/9405187].
- [238] F. Cooper, S. Habib, Y. Kluger, E. Mottola, J. P. Paz, and P. R. Anderson, Phys. Rev. D **50**, 2848 (1994) [arXiv:hep-ph/9405352]; F. Cooper, S. Habib, Y. Kluger, and E. Mottola, Phys. Rev. D **55**, 6471 (1997) [arXiv:hep-ph/9610345].
- [239] D. Boyanovsky, H. J. de Vega, R. Holman, D. S. Lee, and A. Singh, Phys. Rev. D **51**, 4419 (1995); [arXiv:hep-ph/9408214]. D. Boyanovsky, M. D' Attanasio, H. J. de Vega, R. Holman, D. S. Lee, and A. Singh, Phys. Rev. D **52**, 6805 (1995); [arXiv:hep-ph/9507414]. D. Boyanovsky, H. J. de Vega, R. Holman, D. S. Lee, A. Singh, and J. F. J. Salgado, Phys. Rev. D **54**, 7570 (1996); [arXiv:hep-ph/9608205]. Boyanovsky, D. Cormier, H. J. de Vega, R. Holman, D. S. Lee, A. Singh, and M. Srednicki, Phys. Rev. D **56**, 1939 (1997). [arXiv:hep-ph/9703327].
- [240] L. A. Kofman, A. D. Linde, and A. A. Starobinsky, Phys. Rev. Lett. **76**, 1011 (1996) [arXiv:hep-th/9510119]; L. Kofman, A. D. Linde and A. A. Starobinsky,

- Phys. Rev. D **56**, 3258 (1997) [arXiv:hep-ph/9704452]; P. B. Greene, L. A. Kofman, A. D. Linde, and A. A. Starobinsky, Phys. Rev. D **56**, 6175 (1997) [arXiv:hep-ph/9705347].
- [241] R. Allahverdi, and B. A. Campbell, Phys. Lett. B **395**, 169 (1997). [arXiv:hep-ph/9606463].
- [242] M. Bastero-Gil, S. F. King, and J. Sanderson, Phys. Rev. D **60**, 103517 (1999) [arXiv:hep-ph/9904315].
- [243] J. Baacke, and K. Heitmann, Phys. Rev. D **62**, 105022 (2000) [arXiv:hep-ph/0003317]; D. Cormier, K. Heitmann, and A. Mazumdar, Phys. Rev. D **65**, 083521 (2001) [arXiv:hep-ph/0105236].
- [244] S. Khlebnikov, and I. Tkachev, Phys. Rev. Lett. **77**, 219 (1996) [arXiv:hep-ph/9603378]; Phys. Lett. B **390**, 80 (1997) [arXiv:hep-ph/9608458]; Phys. Rev. D **56**, 653 (1997) [arXiv:hep-ph/9701423]; Phys. Rev. Lett. **79**, 1607 (1997); [arXiv:hep-ph/9610477]. T. Prokopec, and T. G. Roos, Phys. Rev. D **55**, 3768 (1997); [arXiv:hep-ph/9610400]. S. Kasuya, and M. Kawasaki, Phys. Lett. B **388**, 686 (1996) [arXiv:hep-ph/9603317].
- [245] A. D. Dolgov, and D. P. Kirilova, Sov. J. Nucl. Phys. **51**, 172 (1990) [Yad. Fiz. **51**, 273 (1990)].
- [246] J. Baacke, K. Heitmann, C. Patzold, Phys. Rev. D **58**, 125013 (1998); [arXiv:hep-ph/9806205]; P.B. Greene, and L. Kofman, Phys. Lett. B **448**, (1999); [arXiv:hep-ph/9807339]; P. B. Greene and L. Kofman, Phys. Rev. D **62**, 123516 (2000) [arXiv:hep-ph/0003018]; M. Peloso, and L. Sorbo, JHEP **0005**, 016 (2000); [arXiv:hep-ph/0003045]; J. Garcia-Bellido, S. Mollerach, and E. Roulet, JHEP **0002**, 034 (2000). [arXiv:hep-ph/0002076].
- [247] A. L. Maroto, and A. Mazumdar, Phys. Rev. Lett. **84**, 1655 (2000) [arXiv:hep-ph/9904206].

- [248] M. Bastero-Gil, and A. Mazumdar, Phys. Rev. D **62**, 083510 (2000) [arXiv:hep-ph/0002004].
- [249] R. Kallosh, L. Kofman, A. Linde, and A. Van Proyen, Phys. Rev. D **61**, 103503 (2000).[arXiv:hep-th/9907124].
- [250] G.F. Giudice, I. Tkachev, A. Riotto, JHEP **9908**, 009 (1999) **9911**, 036 (1999).[arXiv:hep-ph/9907510].
- [251] R. Jeannerot, and J. Lesgourgues, Phys. Rev. D **62**, 023514 (2000) [arXiv:hep-ph/9905208].
- [252] S. Deser, and B. Zumino, Phys. Rev. Lett. **38**, 1433 (1977).
- [253] P. Nath, R. Arnowitt, and A. Chamseddine, *Applied N=1 Supergravity*, Singapore (1984).
- [254] R. Barbieri, S. Ferrara, and C. A. Savoy, Phys. Lett. B **119**, 343 (1982); A. Chamseddine, R. Arnowitt, and P. Nath, Phys. Rev. Lett. **49**, 970 (1982); H. P. Nilles, M. Srednicki, and D. Wyler, Phys. Lett. B **120**, 346 (1983); L. Alvarez-Gaume, J. Polchinsky, M. Wise, Nucl. Phys. B **221**, 495 (1984).
- [255] P. Fayet, Phys. Lett. B **86**, 272 (1979).
- [256] S. Weinberg, Phys. Rev. Lett. **48**, 1303 (1982).
- [257] J. Ellis, and K. A. Olive, Nucl. Phys. B **223**, 252 (1983); L. M. Krauss, Nucl. Phys. B **227**, 177 (1983).
- [258] J. Ellis, D. V. Nanopoulos, and S. Sarkar, Phys. Lett. B **167**, 457 (1986).
- [259] J. Ellis, G. B. Gelmini, D. V. Nanopoulos, J. Lopez and S. Sarkar, Nucl. Phys. B **373**, 399 (1992); R. Juskiewicz, J. Silk, and A. Stebbins, Phys. Lett. B **158**, 463 (1985); S. Dimopoulos, R. Esmailzadeh, L. J. Hall, and G. D. Starkman, Astrophys. J. **330**, 545 (1988); Nucl. Phys. B **311**, 699 (1989).

- [260] M. Kawasaki, and T. Moroi, Prog. Theor. Phys. **93**, 879 (1995); [arXiv:hep-ph/9403364]. Astrophys. J **452**, 506 (1995); Phys. Lett. B **346**, 27 (1995). [arXiv:astro-ph/9412055]. T. Moroi, arXiv:hep-ph/9503210.
- [261] W. Fischler, Phys. Lett. B **332**, 277 (1994) [arXiv:hep-th/9404044].
- [262] R. G. Leigh, and R. Rattazzi, Phys. Lett. B **352**, 20 (1995) [arXiv:hep-ph/9503402].
- [263] J. Ellis, D. V. Nanopoulos, K. A. Olive, and S-J. Rey, Astropart. Phys. **4**, 371 (1996) [arXiv:hep-ph/9505438].
- [264] M. Bolz, A. Brandenburg and W. Buchmuller, Nucl. Phys. B **606**, 518 (2001) [arXiv:hep-ph/0012052].
- [265] M. Dine, A. E. Nelson, Phys. Rev. D **48**, 1277 (1993). [arXiv:hep-ph/9303230].
- [266] E. Pierpaoli, S. Borgani, A. Masiero, and M. Yamaguchi, Phys. Rev. D **57**, 2089 (1998) [arXiv:astro-ph/9709047].
- [267] J. R. Ellis, K. Enqvist, and D.V. Nanopoulos, Phys. Lett. B **147**, 99 (1984); J. R. Ellis, C. Kounnas, and D.V. Nanopoulos, Phys. Lett. B **143**, 410 (1984).
- [268] L. Randall, and R. Sundrum, Nucl. Phys. B **557**, 79 (1999). [arXiv:hep-th/9810155].
- [269] M.Yu. Khlopov, and A. D. Linde, Phys. Lett. B **138**, 265 (1984).
- [270] R. Kallosh, L. Kofman, A. Linde, and A. Van Proyen, Class. Quant. Grav. **17**, 4268 (2000) [arXiv:hep-th/0006179].
- [271] H. P. Nilles, M. Peloso, and L. Sorbo, Phys. Rev. Lett. **87**, 051302 (2001); [arXiv:hep-ph/0102264]. JHEP **0104**, 004 (2001) [arXiv:hep-th/0103202].
- [272] R. Allahverdi, M. Bastero-Gil, and A. Mazumdar, Phys. Rev. D **64**, 023516 (2001) [arXiv:hep-ph/0012057].

- [273] H. P. Nilles, K. A. Olive, and M. Peloso, Phys. Lett. B **522**, 304 (2001) [arXiv:hep-ph/0107212].
- [274] R. Allahverdi, K. Enqvist, and A. Mazumdar, Phys. Rev. D **65**, 103519 (2002) [arXiv:hep-ph/0108225].
- [275] M. Grisaru, W. Siegel, and M. Rocek, Nucl. Phys. B **159**, 429 (1979); N. Seiberg, Phys. Lett. B **318**, 469 (1993). [arXiv:hep-ph/9309335].
- [276] M. Dine, and N. Seiberg, Nucl. Phys. B **306**, 137 (1988).
- [277] B. de Carlos, J.A. Casas, F. Quevedo, and E. Roulet, Phys. Lett. B **318**, 447 (1993) [arXiv:hep-ph/9308325]. T. Banks, M. Berkooz, G. Moore, S. Shenker, and P. J. Steinhardt, Phys. Rev. D **52**, 3548 (1995)[arXiv:hep-th/9503114].
- [278] J. E. Kim, and H. P. Nilles, Phys. Lett. B **138**, 150 (1984);
- [279] G. F. Giudice, and A. Masiero, Phys. Lett. B **206**, 480 (1988); J. E. Kim, and H. P. Nilles, Phys. Lett. B **263**, 79 (1991); J. A. Casas, and C. Munoz, Phys. Lett. B **306**, 288 (1993); [arXiv:hep-ph/9302227]. G.F. Giudice and E. Roulet, Phys. Lett. B **315**, 107 (1993). [arXiv:hep-ph/9307226].
- [280] E. J. Chun, J. E. Kim, and H. P. Nilles, Nucl. Phys. B **370**, 105 (1992).
- [281] G. Dvali, G. F. Giudice, and A. Pomarol, Nucl. Phys. B **478**, 31 (1996). [arXiv:hep-ph/9603238].
- [282] S. Dimopoulos, and H. Georgi, Nucl. Phys. B **193**, 150 (1981).
- [283] R. N. Mohapatra, Phys. Rev. D **34**, 3457 (1986); A. Font, L. E. Ibàñez, and F. Quevedo, Phys. Lett. B **228**, 79 (1989); S. P. Martin, Phys. Rev. D **46**, 2769 (1992); [arXiv:hep-ph/9207218]. Phys. Rev. D **54**, 2340 (1996)[arXiv:hep-ph/9602349].
- [284] H. Goldberg, Phys. Rev. Lett. **50**, 1419 (1983).

- [285] J. Ellis, J. Hagelin, D. V. Nanopoulos, K. Olive, and M. Srednicki, Nucl. Phys. B **238**, 453 (1984).
- [286] F. Buccella, J. P. Derendinger, S. Ferrara, and C. A. Savoy, Phys. Lett. B **115**, 375 (1982).
- [287] I. Affleck, M. Dine, and N. Seiberg, Nucl. Phys. B **241**, 493 (1984);
- [288] I. Affleck, M. Dine, and N. Seiberg, Nucl. Phys. B **256**, 557 (1985).
- [289] M. A. Luty, and W. Taylor, Phys. Rev. D **53**, 3399 (1996). [arXiv:hep-th/9506098].
- [290] S. Dimopoulos, and D. Sutter, Nucl. Phys. B **452**, 496 (1995) [arXiv:hep-ph/9504415]. F. Gabbiani, E. Gabrielli, A. Masiero, and L. Silvestrini, Nucl. Phys. B **477**, 321 (1996)[arXiv:hep-ph/9604387].
- [291] P. Fayet, and J. Iliopoulos, Phys. Lett. B **51**, 461 (1974); P. Fayet, Nucl. Phys. B **90**, 104 (1975).
- [292] L. O’Raifeartaigh, Nucl. Phys. B **96**, 331 (1975).
- [293] E. Witten, Nucl. Phys. B **202**, 253 (1982).
- [294] I. Affleck, M. Dine, and N. Seiberg, Nucl. Phys. B **241**, 493 (1984); Nucl. Phys. B **256**, 557 (1986).
- [295] M. Dine, and W. Fischler, Phys. Lett. B **110**, 227 (1982).
- [296] M. Dine, A. E. Nelson, and Y. Shirman, Phys. Rev. D **51**, 1362 (1995) [arXiv:hep-ph/9408384].
- [297] M. Dine, A. E. Nelson, Y. Nir, and Y. Shirman, Phys. Rev. D **53**, 2658 (1996)[arXiv:hep-ph/9507378].
- [298] Y. Shadmi, and Y. Shirman, Rev. Mod. Phys. **72**, 25 (2000). [arXiv:hep-th/9907225].

- [299] G. F. Giudice, M. Luty, H. Murayama, and R. Rattazzi, JHEP **9812**, 027 (1998) [arXiv:hep-ph/9810442].
- [300] H. P. Nilles, Phys. Lett. B **115**, 193 (1982); S. Ferrara, L. Griardello, and H. P. Nilles, Phys. Lett. B **125**, 457 (1983)
- [301] V. Kaplunovsky, and J. Louis, Phys. Lett. B **306**, 269 (1993) [arXiv:hep-th/9303040]; R. Barbieri, J. Louis, and M. Moretti, Phys. Lett. B **312**, 451 (1993) [Erratum-ibid. B **316**, 632 (1993)] [arXiv:hep-ph/9305262]; A. Brignole, L. E. Ibàñez, and C. Munoz, Nucl. Phys. B **422**, 125 (1994); erratum Nucl. Phys. B **436**, 747 (1995). [arXiv:hep-ph/9308271].
- [302] J. Polonyi, Hungary Central Research Institute Report KFKI-77-93 (1977) (unpublished).
- [303] A. B. Lahanas, and D. V. Nanopoulos, Phys. Rep. **145**, 1 (1987).
- [304] C. R. Nappi, and B. A. Ovrut, Phys. Lett. B **113**, 175 (1982); L. Alvarez-Gaume, M. Claudson, and M. B. Wise, Nucl. Phys. B **207**, 96 (1982).
- [305] M. K. Gaillard, H. Murayama, and K. A. Olive, Phys. Lett. B **355**, 71 (1995) [arXiv:hep-ph/9504307].
- [306] J. Bagger, E. Poppitz, and L. Randall, Nucl. Phys. B **455**, 59 (1995). [arXiv:hep-ph/9505244].
- [307] B. A. Campbell, M. K. Gaillard, H. Murayama, and K. A. Olive, Nucl. Phys. B **538**, 351 (1999). [arXiv:hep-ph/9805300].
- [308] R. Allahverdi, M. Drees, and A. Mazumdar, Phys. Rev. D **65**, 065010 (2002) [arXiv:hep-ph/0110136].
- [309] R. Allahverdi, B. A. Campbell, and J. R. Ellis, Nucl. Phys. B **579**, 355 (2000) [arXiv:hep-ph/0001122].
- [310] A. Anisimov, and M. Dine, Nucl. Phys. B **619**, 729 (2001). [arXiv:hep-ph/0008058].

- [311] N. D. Birrell, and P. C. W. Davies, *Quantum fields in curved space*, Cambridge, Cambridge University Press, (1982); S. A. Fulling, *Aspects of Quantum Field Theory in Curved Space-Time*, Cambridge: Cambridge University Press, (1989).
- [312] Il. Buchbinder, S. D. Odintsov, and I. L. Shapiro, *Effective Action in Quantum Gravity*, Institute of Physics Publishing, Bristol (1992).
- [313] L. Parker, and D. J. Toms, Phys. Rev. Lett. **52**, 1269 (1984); Phys. Rev. D **29**, 1584 (1984); Phys. Rev. D **31**, 2424 (1985); **32**, 1409 (1985).
- [314] M. Drees, and S. P. Martin, *Report of Subgroup 2 of the DPF Working Group on 'Electroweak Symmetry Breaking and Beyond the Standard Model*, Barklow, T.L. (ed.) et al.: Electroweak symmetry breaking and new physics at the TeV scale* 146: arXiv:hep-ph/9504324.
- [315] L.E. Ibàñez, and G.G. Ross, Phys. Lett. B **110**, 215 (1982); L.E. Ibàñez, Phys. Lett. B **118**, 73 (1982); J. Ellis, D.V. Nanopoulos, and K. Tamvakis, Phys. Lett. B **121**, 123 (1983); L. Alvarez-Gaumè, J. Polchinski, and M.B. Wise, Nucl. Phys. B **221**, 495 (1983). L. E. Ibàñez, and C. Lòpez, Nucl. Phys. B **233**, 511 (1984).
- [316] C. T. Hill, Phys. Rev. D **24**, 691 (1981); B. Pendleton, and G.G. Ross, Phys. Lett. B **98**, 291 (1981); V. Barger, M.S. Berger, P. Ohmann, and R.J.N. Phillips, Phys. Lett. B **314**, 351 (1993) [arXiv:hep-ph/9304295].
- [317] M. Carena, and C.E.M. Wagner, Nucl. Phys. B **452**, 45 (1995). [arXiv:hep-ph/9408253].
- [318] The LEP collaborations, hep-ex/0107029.
- [319] S. Mollerach, Phys. Lett. B **242**, 158 (1990).
- [320] K. Enqvist and M. S. Sloth, Nucl. Phys. B **626**, 395 (2002) [arXiv:hep-ph/0109214].

- [321] D. H. Lyth and D. Wands, Phys. Lett. B **524**, 5 (2002) [arXiv:hep-ph/0110002].
- [322] D. H. Lyth, C. Ungarelli and D. Wands, arXiv:astro-ph/0208055; N. Bartolo and A. R. Liddle, Phys. Rev. D **65**, 121301 (2002) [arXiv:astro-ph/0203076]; T. Moroi and T. Takahashi, Phys. Rev. D **66**, 063501 (2002) [arXiv:hep-ph/0206026]; M. S. Sloth, arXiv:hep-ph/0208241; M. Bastero-Gil, V. Di Clemente and S. F. King, arXiv:hep-ph/0211012. T. Moroi and H. Murayama, arXiv:hep-ph/0211019. C. Gordon and A. Lewis, arXiv:astro-ph/0212248; M. Postma, arXiv:hep-ph/0212005.
- [323] K. Enqvist, S. Kasuya and A. Mazumdar, arXiv:hep-ph/0211147.
- [324] J. McDonald, Phys. Lett. B **456**, 118 (1999) [hep-ph/9902416].
- [325] P. Binetruy, and M. K. Gaillard, Phys. Lett. B **195**, 382 (1987); P. Binetruy, and M. K. Gaillard, Phys. Lett. B **232**, 83 (1989).
- [326] J. Ellis, A. B. Lahanas, D. V. Nanopoulos, and K. Tamvakis, Phys. Lett. B **134**, 429 (1984); J. Ellis, C. Kounnas, and D. V. Nanopoulos, Nucl. Phys. B **241**, 406 (1984); **247**, 373 (1984).
- [327] M. K. Gaillard, and V. Jain, Phys. Rev. D **49**, 49 (1994). [arXiv:hep-th/9308090].
- [328] K. Enqvist, and J. McDonald, Nucl. Phys. B **570**, 407 (2000) [arXiv:hep-ph/9908316].
- [329] K. Enqvist, A. Jokinen, and J. McDonald, Phys. Lett. B **483**, 191 (2000) [arXiv:hep-ph/0004050].
- [330] K. Enqvist, A. Jokinen, T. Multamäki, and I. Vilja, Phys. Rev. D **63**, 083501 (2001) [arXiv:hep-ph/0011134].
- [331] G. R. Dvali, A. Kusenko, and M. E. Shaposhnikov, Phys. Lett. B **417**, 99 (1998) [arXiv:hep-ph/9707423].

- [332] K. Koyama, and J. Soda, Phys. Rev. Lett. **82**, 2632 (1999). [arXiv:astro-ph/9810006].
- [333] J. A. Peacock, and S. J. Dodds, MNRAS, **267**, 1020 (1994). [arXiv:astro-ph/9311057].
- [334] J. Casas, and G. B. Gelmini, Phys. Lett. B **410**, 36 (1997). [arXiv:hep-ph/9706439].
- [335] M. Senami, and K. Yamamoto, Phys. Lett. B **524**, 332 (2002). [arXiv:hep-ph/0105054].
- [336] A. D. Dolgov, and D. P. Kirilova, Sov. J. Nucl. Phys. **50**, 1006 (1989), Yad. Fiz. **50**, 1621 (1989); J. Moscow. Phys. Soc. **1**, 217 (1991). A. D. Dolgov, and J. Silk, Phys. Rev. D **47**, 4244 (1993); M. Chizhov, and D. Kirilova, Astron. Astro. Trans. **10**, 69 (1996); T. Matsuda, Phys. Rev. D **65**, 103501 (2002) [arXiv:hep-ph/0202211].
- [337] T. Asaka, M. Fujii, K. Hamaguchi, and T. Yanagida, Phys. Rev. D **62**, 123514 (2000). [arXiv:hep-ph/0008041].
- [338] T. Asaka, Phys. Lett. B **521**, 329 (2001). [arXiv:hep-ph/0110073].
- [339] E. D. Stewart, M. Kawasaki, and T. Yanagida, Phys. Rev. D **54**, 6032 (1996)[arXiv:hep-ph/9603324].
- [340] S. Kasuya, M. Kawasaki, and F. Takahashi, Phys. Rev. D **65**, 063509 (2002) [arXiv:hep-ph/0108171].
- [341] J. McDonald, Phys. Rev. D **55**, 4240 (1997). [arXiv:hep-ph/9610322].
- [342] A. Anisimov, *Thermal effects and flat direction baryogenesis*, arXiv:hep-ph/0111233.
- [343] M. Fujii, K. Hamaguchi, and T. Yanagida, Phys. Rev. D **63**, 123513 (2001). [arXiv:hep-ph/0102187].

- [344] A. Jokinen, *Analytical and numerical properties of Affleck-Dine Condensate formation*. arXiv: hep-ph/0204086.
- [345] G. Rosen, J. Math. Phys. **9**, 996 (1986).
- [346] T. D. Lee, and G. C. Wick, Phys. Rev. D **9**, 2291 (1974).
- [347] K. Enqvist, and J. McDonald, Phys. Rev. Lett. **81**, 3071 (1998) [arXiv:hep-ph/9806213].
- [348] S. Kasuya, and M. Kawasaki, Phys. Rev. D **61**, 041301 (2000) [arXiv:hep-ph/9909509].
- [349] S. Kasuya, and M. Kawasaki, Phys. Rev. Lett. **85**, 2677 (2000) [arXiv:hep-ph/0006128].
- [350] S. Kasuya, and M. Kawasaki, Phys. Rev. D **62**, 023512 (2000) [arXiv:hep-ph/0002285].
- [351] S. Kasuya, and M. Kawasaki, Phys. Rev. D **64**, 123515 (2001) [arXiv:hep-ph/0106119].
- [352] T. Multamäki, and I. Vilja, Phys. Lett. B **535**, 170 (2002). [arXiv:hep-ph/0203195].
- [353] K. Enqvist, S. Kasuya, and A. Mazumdar, Phys. Rev. Lett. **89**, 091301 (2002) [arXiv:hep-ph/0204270].
- [354] K. Enqvist, S. Kasuya, and A. Mazumdar, Phys. Rev. D **68**, 043505 (2002) [arXiv:hep-ph/0206272].
- [355] A. Kusenko, V. Kuzmin, M. E. Shaposhnikov, and P.G. Tinyakov, Phys. Rev. Lett. **80**, 3185 (1998) [arXiv:hep-ph/9712212].
- [356] D. A. Demir, Phys. Lett. B **450**, 215 (1999). [arXiv:hep-ph/9810453].
- [357] A. Kusenko, and P. J. Steinhardt, Phys. Rev. Lett. **87**, 141301 (2001) [arXiv:astro-ph/0106008].

- [358] K. Enqvist, A. Jokinen, T. Multamäki, and Iiro Vilja, Phys. Lett. B **526**, 9 (2002) [arXiv:hep-ph/0111348].
- [359] K. Enqvist, and J. McDonald, Phys. Lett. B **440**, 59 (1998) [arXiv:hep-ph/9807269].
- [360] J. McDonald, JHEP **0103**, 022 (2001). [arXiv:hep-ph/0012369].
- [361] M. Fujii, and K. Hamaguchi, Phys. Lett. B **525**, 143 (2002). [arXiv:hep-ph/0110072].
- [362] M. Fujii, and K. Hamaguchi, Phys. Rev. D **66**, 083501 (2002) [arXiv:hep-ph/0205044].
- [363] T. Shiromizu, Phys. Rev. D **58**, 107301 (1998). [arXiv:hep-ph/9811420].
- [364] A. Kusenko, M. E. Shaposhnikov, P.G. Tinyakov, and I. I. Tkachev, Phys. Lett. B **423**, 104 (1998) [arXiv:hep-ph/9801212].
- [365] T. D. Lee, and Y. Pang, Phys. Rep. **221**, 251 (1992).
- [366] B. W. Lynn, Nucl. Phys. B **321**, 465 (1989).
- [367] A. Prikas, Phys. Rev. D **66**, 025023 (2002) [arXiv:hep-th/0205197].
- [368] R. Friedberg, T. D. Lee, and A. Sirlin, Phys. Rev. D **13**, 2739 (1976).
- [369] T. F. Morris, Phys. Lett. B **76**, 337 (1978), **78**, 87 (1978).
- [370] G. 't Hooft, Nucl. Phys. B **79**, 276 (1974).
- [371] A. M. Polyakov, JETP Lett. **20**, 194 (1974), [Pisma Zh. Eksp. Teor. Fiz. **20** (1974) 430].
- [372] L. Wilets, Nontopological solitons, Lecture Notes in Physics, Vol. **24**, (World Scientific, Singapore, 1989).
- [373] A. M. Safian, S. Coleman, and M. Axenides, Nucl. Phys. B **297**, 498 (1988);
A. M. Safian. Nucl. Phys. B **304**, 392 (1988).

- [374] M. Axenides, Int. J. Mod. Phys. A **7**, 7169 (1992)
- [375] M. Axenides, E. Floratos, and A. Kehagias, Phys. Lett. B **444**, 190 (1998) [arXiv:hep-ph/9810230].
- [376] K. Lee, J. A. Stein-Schabes, R. Watkins, and L. M. Widrow, Phys. Rev. D. **39**, 1665 (1989).
- [377] A. Khare, and S. Rao, Phys. Lett. B **227** 424 (1989).
- [378] T. Shiromizu, T. Uesugi, and M. Aoki, Phys. Rev. D **59**, 125010 (1999) [arXiv:hep-ph/9811420].
- [379] K. N. Anagnostopoulos, M. Axenides, E. G. Floratos, and N. Tetradis, Phys. Rev. D **64**, 125006 (2001). [arXiv:hep-ph/0109080].
- [380] M. Fujii, K. Hamaguchi, and T. Yanagida, Phys. Rev. D **64**, 123526 (2001); [arXiv:hep-ph/0104186]. **65**, 043511 (2002). [arXiv:hep-ph/0109154].
- [381] T. S. Levi, and M. Gleiser, Phys. Rev. D **66**, 087701 (2002) [arXiv:hep-ph/0110395].
- [382] A. Kusenko, Phys. Lett. B **404**, 285 (1997) [arXiv:hep-th/9704073].
- [383] F. P. Correia, and M.G. Schmidt, Eur. Phys. J. C **21** 181 (2001). [arXiv:hep-th/0103189].
- [384] I. Yu. Kobzarev, L. B. Okun, and M. B. Voloshin, Yad. Fiz. **20**, 1229 (1974), Sov. J. Nucl. Phys. **20**, 644 (1975); P. H. Frampton, Phys. Rev. D **15**, 2922 (1977).
- [385] A. Kusenko, M. E. Shaposhnikov, and P.G. Tinyakov, Pisma Zh. Eksp. Teor. Fiz. **67**, 229 (1998), JETP Lett. **67**, 247 (1998) [arXiv:hep-th/9801041].
- [386] N. Graham, Phys. Lett. B **513**, 112 (2001). [arXiv:hep-th/0105009].
- [387] A. de Gouvêa, T. Moroi, and H. Murayama, Phys. Rev. D **56**, 1281 (1997) [arXiv:hep-ph/9701244].

- [388] T. Multamäki, and I. Vilja, Phys. Lett. B **542**, 137 (2002) [arXiv:hep-ph/0205302].
- [389] R. Friedberg, and T. D. Lee, Phys. Rev. D **15**, 1694 (1977); **16**, 1096 (1977); **18**, 2623 (1978).
- [390] A. Cohen, S. Coleman, H. Georgi, and A. Manohar, Nucl. Phys. B **272**, 301 (1986).
- [391] T. Multamäki, and I. Vilja, Nucl. Phys. B **574**, 130 (2000) [arXiv:hep-ph/9908446].
- [392] V. G. Makhankov, G. Kummer, and A. B. Shvachka, Phys. Lett. A **70**, 171 (1979); J. K. Drohm, L. P. Kok, Y. A. Simonov, J. A. Tjon and A. I. Veselov, Phys. Lett. B **101**, 204 (1981); T. I. Belova, and A. E. Kudryavtsev, Sov. Phys. JETP **68**, 7 (1989) [Zh. Eksp. Teor. Fiz. **95**, 13 (1989)].
- [393] M. Axenides, S. Komineas, L. Perivolaropoulos, and M. Floratos, Phys. Rev. D **61**, 085006 (2000). [arXiv:hep-ph/9910388].
- [394] R. A. Battye, and P. M. Sutcliffe, Nucl. Phys. B **590**, 329 (2000). [arXiv:hep-th/0003252].
- [395] T. Multamäki, and I. Vilja, Phys. Lett. B **482**, 161 (2000). [arXiv:hep-ph/0003270].
- [396] T. Multamäki, and I. Vilja, Phys. Lett. B **484**, 283 (2000). [arXiv:hep-ph/0005162].
- [397] M. Laine, and M. Shaposhnikov, Nucl. Phys. B **532**, 376 (1998). [arXiv:hep-lat/9805013].
- [398] R. Banerjee, and K. Jedamzik, Phys. Lett. B **484**, 278 (2000). [arXiv:hep-ph/0005031].
- [399] M. Fujii, and T. Yanagida, Phys. Lett. B **542**, 80 (2002). [arXiv:hep-ph/0206066].

- [400] J. Ellis, T. Falk, K. A. Olive, and M. Schmitt, Phys. Lett. B **413**, 355 (1997) [arXiv:hep-ph/9705444].
- [401] D. Hutchcroft, Nucl. Phys. Proc. Suppl. **87**, 99 (2000). [arXiv:hep-ex/9912015].
- [402] R. Allahverdi, A. Mazumdar, and A. Özpineci, Phys. Rev. D **65**, 125003 (2002) [arXiv:hep-ph/0203062].
- [403] V. A. Bednyakov, H. V. Klapdor-Kleingrothaus, and S. Kovalenko, Phys. Rev. D **50**, 7128 (1994). [arXiv:hep-ph/9401262].
- [404] M. Drees, and M. Nojiri, Phys. Rev. D **48**, 3483 (1993). [arXiv:hep-ph/9307208].
- [405] R. W. Schnee et. al., Phys. Rept. **307**, 283 (1998).
- [406] M. Bravin et.al., Astropart. Phys. **12**, 107 (1999). [arXiv:hep-ex/9904005].
- [407] A. Juillard et.al., Nucl. Phys. Proc. Suppl. **110**, 70 (2002). [arXiv:astro-ph/0104192].
- [408] H. V. Klapdor-Kleingrothaus et.al., Nucl. Phys. Proc. Suppl. **100**, 350 (2001)[arXiv:hep-ph/0102277].
- [409] D. B. Cline, *ZEPLIN IV: A one ton Ultimate WIMP detector*, arXiv: astro-ph/0111098.
- [410] L. Bergstrom, and P. Ullio, Nucl. Phys. B **504**, 27 (1997); [arXiv:hep-ph/9706232]. P. Ullio, and L. Bergstrom, Phys. Rev. D **57**, 1962 (1998). [arXiv:hep-ph/9707333].
- [411] S..M. Bradbury et. al., Proc. of 26th ICRC, Salt Lake City, 1999, OG. 4. 3. 28.
- [412] A. Kohnle et. al., Proc. of 26th ICRC, Salt Lake City, 1999, OG. 4. 3. 13.
- [413] M. Martinez et. al., Proc. of 26th ICRC, Salt Lake City, 1999, OG. 4. 3. 08.

- [414] L. Bergstrom, P. Ullio, and J. H. Buckley, *Astropart. Phys.* **9**, 137 (1998).
[arXiv:astro-ph/9712318].
- [415] J. Hisano, K. Kohri, and M. M. Nojiri, *Phys. Lett. B* **505**, 169 (2001).
[arXiv:hep-ph/0011216].
- [416] B. Moore, *Nature* **370**, 629 (1994) [astro-ph/9402009]; J. F. Navarro, C. Frenk, and S. White, *Astrophys. J.* **462**, 536 (1996); [arXiv:astro-ph/9508025]. B. Moore, et.al., *Astrophys. J. Lett.* **499**, L5 (1998) [astro-ph/9709051]; A. Klypin et.al., *Astrophys. J.* **522**, 82 (1999) [arXiv:astro-ph/9901240]; B. Moore, et.al., *Astrophys. J. Lett.* **524**, L19 (1999) [astro-ph/9907411].
- [417] D. N. Spergel, and P. J. Steinhardt, *Phys. Rev. Lett.* **84**, 3760 (2000).
[arXiv:astro-ph/9909386].
- [418] G. Gelmini, A. Kusenko, and S. Nussinov, *Phys. Rev. Lett.* **89**, 101302 (2002)
[arXiv:hep-ph/0203179].
- [419] S. Desai, *4th International Symposium on Sources and Detection of Dark Matter*, Universe at Marina Del Ray, Feb. 2000.
- [420] S. G. Bayburina et.al., *Proc. 16th Intern. Cosmic Ray Conf.*, **7**, 279 (1979).
- [421] S. Ahlen et.al., *Phys. Rev. Lett.* **69**, 1860 (1992).
- [422] L. A. Boyle, R. R. Caldwell and M. Kamionkowski, *Phys. Lett. B* **545**, 17 (2002) [arXiv:astro-ph/0105318].
- [423] S. Kasuya, *Phys. Lett. B* **515**, 121 (2001) [arXiv:astro-ph/0105408].
- [424] E. J. Copeland, S. Pascoli, and A. Rajantie, *Phys. Rev. D* **65**, 103517 (2002)
[arXiv:hep-ph/0202031].
- [425] N. Arkani-Hamed, S. Dimopoulos, and G. Dvali, *Phys. Lett. B* **429**, 263 (1998)
[arXiv:hep-ph/9803315]; I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and

- G. R. Dvali, Phys. Lett. B **436**, 257 (1998) [arXiv:hep-ph/9804398]; N. Arkani-Hamed, S. Dimopoulos and G. R. Dvali, Phys. Rev. D **59**, 086004 (1999) [arXiv:hep-ph/9807344].
- [426] For some early ideas see also: I. Antoniadis, Phys. Lett. B **246**, 377 (1990); I. Antoniadis, K. Benakli, and M. Quirós, Phys. Lett. B **331**, 313 (1994); [arXiv:hep-ph/9403290]; K. Benakli, Phys. Rev. D **60**, 104002 (1999) [arXiv:hep-ph/9809582]; K. Benakli, Phys. Lett. B **447**, 51 (1999) [arXiv:hep-th/9805181].
- [427] C. D. Hoyle *et al*, Phys. Rev. Lett. **86**, 1418 (2001). [arXiv:hep-ph/0011014].
- [428] S. Hannestad, and G. G. Raffelt, Phys. Rev. Lett. **88**, 071301 (2002). [arXiv:hep-ph/0110067].
- [429] For experimental bounds see for instance: T. G. Rizzo, Phys. Rev. D **59**, 115010 (1999); G. F. Giudice, R. Rattazzi, and J. D. Wells, Nucl. Phys. **B544**, 3 (1999); E. A. Mirabelli, M. Perelstein, and Michael E. Peskin, Phys. Rev. Lett. **82**, 2236 (1999); J. L. Hewett, Phys. Rev. Lett. **82**, 4765 (1999); V. Barger, T. Han, C. Kao, and R. J. Zhang, *Phys. Lett.* **B461** (,) 34 (1999).
- [430] K. Benakli, and S. Davidson, *Phys. Rev. D* **60** (,) 025004 (1999); L. J. Hall, and D. Smith. Phys. Rev. D **60**, 085008 (1999).
- [431] D. H. Lyth, Phys. Lett **B 448**, 191 (1999) [arXiv:hep-ph/9810320]; G. Dvali, S. H. H. Tye, Phys. Lett. **B450**, 72 (1999) [arXiv:hep-ph/9812483]; N. Kaloper and A. Linde, Phys. Rev **D 59**, 101303 (1999) [arXiv:hep-th/9811141]. N. Arkani-Hamed, *et al.*, Nucl. Phys. **B567**, 189 (2000) [arXiv:hep-ph/9903224]; J. Cline, Phys. Rev. D **61**, 023513 (2000). [arXiv:hep-ph/9904495].
- [432] A. Pérez-Lorenzana, AIP Conf. Proc. **562**, 53 (2001) [arXiv:hep-ph/0008333]; for brane world scenarios, see: V. A. Rubakov, [Usp. Fiz. Nauk **171**, 913 (2001)] [arXiv:hep-ph/0104152].

- [433] R.N. Mohapatra, A. Pérez-Lorenzana and C.A. de S. Pires, Phys. Rev. **D 62**, 105030 (1999) [arXiv:hep-ph/0003089].
- [434] A. Mazumdar, and A. Pérez-Lorenzana, Phys. Rev. D **65**, 107301 (2002) [arXiv:hep-ph/0103215].
- [435] R. Allahverdi, K. Enqvist, A. Mazumdar, and A. Pérez-Lorenzana, Nucl. Phys. B **618**, 277 (2001) [arXiv:hep-ph/0108225].
- [436] A. M. Green, and A. Mazumdar, Phys. Rev. D **65**, 105022 (2002) [arXiv:hep-ph/0201209].
- [437] A. Pilaftsis, Phys. Rev. D **60**, 105023 (2000). [arXiv:hep-ph/9906265].
- [438] A. Mazumdar, Phys. Rev. D **64**, 027304 (2001) [arXiv:hep-ph/0007269]; A. Mazumdar, Nucl. Phys. B **597**, 561 (2001) [arXiv:hep-ph/0008087]; D. J. H. Chung, and T. Dent, Phys. Rev. D **66**, 023501 (2002) [arXiv:hep-ph/0112360]; T. Matsuda, Phys. Rev. D **65**, 103501 (2002) [arXiv:hep-ph/0202209]; A. Masiero, M. Peloso, L. Sorbo and R. Tabbash, Phys. Rev. D **62**, 063515 (2000) [arXiv:hep-ph/0003312]; In the context of MSSM, see G. Servant, JHEP **0201**, 044 (2002); [arXiv:hep-ph/0112209].
- [439] J. A. Frieman, G. B. Gelmini, M. Gleiser, and E. W. Kolb, Phys. Rev. Lett. **60**, 2101 (1988).
- [440] J. A. Frieman, A. V. Olinto, M. Gleiser, and C. Alcock, Phys. Rev. D **40**, 3241 (1989).
- [441] K. Griest, and E. Kolb, Phys. Rev. D **40**, 3231 (1989).
- [442] J. R. Ellis, K. Enqvist, D. V. Nanopoulos, K. A. Olive, Phys. Lett. B **225**, 313 (1989).
- [443] A. Kusenko, Phys. Lett.B **406**, 26 (1997) [arXiv:hep-ph/9705361].
- [444] M. Postma, Phys. Rev. D **65**, 085035 (2002) [arXiv:hep-ph/0110199].